CHEMISTRY FOR ENGINEERS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://chemforengr.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What bond forms by transfer of electrons between atoms?

A. Covalent Bond

B. Ionic Bond

C. Metallic Bond

D. Hydrogen Bond

2. For a nonvolatile solute, boiling point elevation ΔTb is given by which expression?

A. $\Delta T_b = i K_b m$

B. $\Delta T_b = i K_b$

C. $\Delta T_b = K_b m / i$

D. $\Delta T_b = m i / K_b$

3. What is the integrated rate law for a first-order reaction?

A. $[A] = [A]_0 e^{(kt)}$

B. $\ln[A] = \ln[A]_0 - kt$

C. $1/[A] = 1/[A]_0 + kt$

D. $[A] = [A]_0 - kt$

4. Which property describes the ability of an atom to attract electrons in a bond?

A. Electronegativity

B. Ionic Bond

C. Covalent Bond

D. Hydrogen Bond

5. Minimum energy required to start a reaction is called

A. Catalysis

B. Activation energy

C. Collision energy

D. Transition state energy

SAMPLE

6. Movement of particles from high to low concentration.

A. Osmosis

B. Diffusion

C. Dialysis

D. Filtration

7. If a reactant concentration is suddenly doubled and the system is allowed to re-establish equilibrium at the same temperature, what happens to the value of the equilibrium constant K?

A. K increases

B. K decreases

C. K remains the same

D. K becomes undefined

8. Which term is defined by the equation PV = nRT for gases?

A. STP

B. Arrhenius acid

C. Ideal Gas Law

D. Brønsted-Lowry base

9. In the osmotic pressure equation π = i M R T for dilute solutions, what does i represent?

A. The molar mass of solute.

B. The universal gas constant.

C. The van't Hoff factor, the effective number of particles in solution.

D. The temperature in Kelvin.

10. What is the significance of a "basis" in reactor calculations for a chemical process?

A. A basis defines the reactor's design pressure.

B. A basis fixes a reference flow rate or amount (such as 1 hour, 1 mole, or 100 mol) to express conversions and rates.

C. A basis determines the exact temperature of operation.

D. A basis is a fixed reference temperature for calculations.

SAMPLE

Answers

SAMPLE

1. B
2. A
3. B
4. A
5. B
6. B
7. C
8. C
9. C
10. D

SAMPLE

Explanations

SAMPLE

1. What bond forms by transfer of electrons between atoms?

A. Covalent Bond

B. Ionic Bond

C. Metallic Bond

D. Hydrogen Bond

When electrons are transferred from one atom to another, the atoms become oppositely charged ions that attract each other. This electrostatic attraction pulls the ions together, forming an ionic bond. It typically happens between a metal (which tends to lose electrons) and a nonmetal (which tends to gain electrons), creating a compound composed of a lattice of cations and anions. This is different from covalent bonds, where electrons are shared; or metallic bonds, where electrons are delocalized across a metal lattice; and from hydrogen bonds, which are intermolecular attractions rather than actual electron transfer.

2. For a nonvolatile solute, boiling point elevation ΔTb is given by which expression?

A. $\Delta T_b = i K_b m$

B. $\Delta T_b = i K_b$

C. $\Delta T_b = K_b m / i$

D. $\Delta T_b = m i / K_b$

Boiling point elevation is a colligative property, meaning it depends on how many dissolved particles the solute adds to the solvent, not on their identity. In dilute solutions, the rise in boiling temperature is proportional to the molality of the solute, the solvent's ebullioscopic constant, and the number of particles produced per formula unit of solute, captured by $\Delta T_b = i K_b m$. Here, i is the van't Hoff factor, which equals 1 for non-dissociating solutes and is greater than 1 for electrolytes that dissociate into multiple particles. K_b is a property of the solvent, and m is the molality. This is why the correct expression multiplies i, K_b, and m. The other forms either omit a factor or place i or K_b in the wrong position, which would not reflect the way the boiling point elevation scales with particle number, solvent characteristics, and concentration.

3. What is the integrated rate law for a first-order reaction?

A. $[A] = [A]_0 e^{(kt)}$

B. $\ln[A] = \ln[A]_0 - kt$

C. $1/[A] = 1/[A]_0 + kt$

D. $[A] = [A]_0 - kt$

A first-order reaction has a rate that depends on the concentration of A, described by $-d[A]/dt = k[A]$. Integrating this relationship gives $\ln[A] = \ln[A]_0 - kt$, which is the integrated rate law. This form shows the natural log of the concentration decreasing linearly with time, with slope $-k$, and it can also be written as $[A](t) = [A]_0 e^{(-kt)}$. This is the best answer because the logarithmic form directly reflects the exponential decay characteristic of a first-order process. The other forms correspond to different scenarios: $[A] = [A]_0 - kt$ would imply a constant rate (zero-order), $[A] = [A]_0 e^{(kt)}$ would imply growth rather than decay, and $1/[A] = 1/[A]_0 + kt$ describes a second-order process where $1/[A]$ grows linearly with time.

SAMPLE

4. Which property describes the ability of an atom to attract electrons in a bond?

A. Electronegativity

B. Ionic Bond

C. Covalent Bond

D. Hydrogen Bond

Electronegativity is the tendency of an atom to attract electrons toward itself when forming a chemical bond. This property explains how electron density is shared unequally in bonds, leading to bond polarity—for example, chlorine pulls electron density toward itself in H–Cl, creating a dipole. Ionic bonds describe a type of bond formed by transfer of electrons and electrostatic attraction between ions, not a single atom's pull on electrons. Covalent bonds describe sharing of electron pairs between atoms, which is influenced by electronegativity but is not itself the measure of an atom's pull. Hydrogen bonds are a specific intermolecular interaction that arises from polarity and hydrogen's attraction to electronegative atoms, not a description of an atom's inherent ability to attract electrons. Therefore, electronegativity best captures the concept of an atom attracting electrons in a bond.

5. Minimum energy required to start a reaction is called

A. Catalysis

B. Activation energy

C. Collision energy

D. Transition state energy

Activation energy is the minimum energy required to start a reaction. It represents the energy barrier that reactants must overcome to reach the transition state and proceed to products. On a potential energy diagram, this is the difference between the energy of the reactants and the highest-energy point along the reaction path—the transition state. Not every collision provides enough energy, and only those with energy at least equal to the activation energy can lead to reaction; temperature affects how many molecules meet this threshold, and this is captured by the Arrhenius relationship. Catalysts work by offering an alternative pathway with a lower transition-state energy, thereby lowering the activation energy and increasing the reaction rate. The energy of the transition state itself is higher than the reactants by the amount of the activation energy, which is why Ea is the key quantity describing the minimum energy needed to initiate the reaction.

SAMPLE

6. Movement of particles from high to low concentration.

A. Osmosis

B. Diffusion

C. Dialysis

D. Filtration

Diffusion is the net movement of particles from regions of higher concentration to regions of lower concentration, driven by the random thermal motion of molecules. This random motion causes particles to spread out until the concentrations balance, and it happens without needing external energy input. The rate of diffusion increases with higher temperature (more kinetic energy) and with smaller particles or a less viscous medium, while moving more slowly for larger molecules or through denser media. Osmosis is specifically about the movement of water across a selectively permeable membrane, typically from a region of lower solute concentration to higher solute concentration, so it's a solvent movement rather than a general solute diffusion. Dialysis is a practical technique that uses diffusion of small solutes across a semipermeable membrane to separate substances; it's diffusion in a membrane context, not the broad concept itself. Filtration relies on applying pressure to move a mixture through a barrier and separates components by size, not by a concentration gradient. So the broad, defining idea that matches "movement of particles from high to low concentration" is diffusion.

7. If a reactant concentration is suddenly doubled and the system is allowed to re-establish equilibrium at the same temperature, what happens to the value of the equilibrium constant K?

A. K increases

B. K decreases

C. K remains the same

D. K becomes undefined

The value of the equilibrium constant is determined by temperature, not by how much reactant you start with. At the same temperature, K is fixed for a given reaction. If you suddenly double the reactant concentration, the system shifts to re-establish equilibrium (Le Chatelier's principle), producing more products or adjusting the mix until the ratio of products to reactants, each raised to their stoichiometric powers, matches the same K as before. So, once equilibrium is re-established, K remains unchanged. If the temperature were changed, then K would change accordingly; at constant temperature, it stays the same.

SAMPLE

8. Which term is defined by the equation PV = nRT for gases?

A. STP

B. Arrhenius acid

C. Ideal Gas Law

D. Brønsted-Lowry base

This is about how a gas behaves when it follows a simple, idealized relationship between pressure, volume, amount, and temperature. PV = nRT ties these quantities together with the universal gas constant R, showing that pressure times volume depends on how much gas there is (n) and how hot it is (T). Because of this equation, increasing the temperature at constant n and P raises the volume, while compressing the gas (reducing V) at the same n and T raises the pressure. The constant R makes the units work out; with common units, R is 0.0821 L·atm·mol ¹·K ¹ or 8.314 J·mol ¹·K ¹, depending on whether you're using liters-atmospheres or joules for energy. For example, 1 mole of an ideal gas at 25°C (298 K) and 1 atm would occupy about 24.5 L, since V = nRT/P. At standard temperature and pressure (STP), the molar volume is about 22.4 L per mole. The other terms listed are about standard conditions or acid–base definitions (STP describes conditions; Arrhenius acid and Brønsted–Lowry base describe how substances behave in reactions), not a gas-law equation.

9. In the osmotic pressure equation π = i M R T for dilute solutions, what does i represent?

A. The molar mass of solute.

B. The universal gas constant.

C. The van't Hoff factor, the effective number of particles in solution.

D. The temperature in Kelvin.

Osmotic pressure depends on how many solute particles are present in the solution. The i in the equation is the van't Hoff factor, the effective number of particles produced when the solute dissolves. This factor adjusts the concentration term to reflect how the solute contributes to osmotic pressure. For a nonelectrolyte that does not ionize, i is about 1, so π scales with M, R, and T just as in the simple gas-like analogy. For electrolytes that dissociate into ions, i reflects how many particles appear after dissociation (for a fully dissociated salt that gives two particles per formula unit, i ≈ 2). In real solutions, i can be a bit less than the ideal integer due to incomplete dissociation or ion pairing, but it still represents the effective particle count driving osmotic pressure. The other quantities in the formula—M, R, and T—represent molarity, the gas constant, and temperature, respectively, and are not what i stands for.

SAMPLE

10. What is the significance of a "basis" in reactor calculations for a chemical process?

A. A basis defines the reactor's design pressure.

B. A basis fixes a reference flow rate or amount (such as 1 hour, 1 mole, or 100 mol) to express conversions and rates.

C. A basis determines the exact temperature of operation.

D. A basis is a fixed reference temperature for calculations.

In reactor calculations, the basis is a fixed reference quantity used to express conversions and reaction rates in a consistent, scalable way. By choosing a basis such as 1 hour of operation, 1 mole of feed, or 100 moles, all quantities—conversions, yields, and rates—are written relative to that reference, which makes mass balances and rate expressions unambiguous and easy to scale to actual conditions. This normalization has nothing to do with setting design pressure or the operating temperature—the basis is about how you standardize amounts and flows. For example, if you adopt a mole basis, a conversion of 0.8 means 0.8 mole of reactant has reacted per mole of feed (depending on the exact definition used), and you can translate that to a per-hour or per-second rate once you know the basis. The temperature or pressure you operate at can then be applied to the basis-normalized quantities to obtain real-world numbers.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://chemforengr.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE