
CJE MATERNITY 1 PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://cjematernity1.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. True or false: Accelerations in the fetal heart rate indicate a healthy fetal and placental exchange.

A. True

B. False

C. True only in the absence of contractions

D. False unless monitored continuously

2. What is the main purpose of monitoring for hypotension in a mother prior to epidural or spinal analgesia?

A. To ensure a comfortable position during labor

B. To prevent complications related to anesthesia

C. To assess fetal heart rate

D. To prepare for cesarean delivery

3. Which of the following indicates a serious side effect of magnesium sulfate?

A. Lethargy

B. Visual blurring

C. Headache

D. Weight gain

4. True or False: Fetal distress is a common complication of labor.

A. True

B. False

C. Only in late labor

D. Only in early labor

5. What are some adverse effects associated with oxytocin use?

A. Uterine Tachysystole

B. Increased contractions

C. Fetal hypoxia

D. Risk for uterine rupture

SAMPLE

6. Which of the following are signs of an infection in the belly button area?

A. Dryness and flakiness

B. Youthful appearance

C. Redness and swelling

D. Complaints of pain only

7. What does a score of 0 for activity in an Apgar score signify?

A. Active movement

B. Flexed arms and legs

C. Absent muscle tone

D. Minimal movement

8. What intervention should the nurse expect to implement for a hypoglycemic SGA newborn?

A. Administer glucose immediately regardless of feeding time.

B. Obtain a glucose level right before feeding.

C. Delay feeding until the next shift.

D. Monitor oxygen levels continuously.

9. What is the average gestational weight of a newborn born at 37 weeks?

A. 8 lbs. 6 ounces

B. 5 lbs. 13 ounces

C. 6 lbs. 1 ounce

D. 7 lbs. 3 ounces

10. True or false: It is normal for the umbilical cord stump to ooze a little before it falls off.

A. True

B. False

C. Only during the first day

D. Requires immediate attention

SAMPLE

Answers

SAMPLE

1. A
2. B
3. A
4. B
5. A
6. C
7. C
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. True or false: Accelerations in the fetal heart rate indicate a healthy fetal and placental exchange.

A. True

B. False

C. True only in the absence of contractions

D. False unless monitored continuously

Accelerations in the fetal heart rate are indeed a positive sign indicating healthy fetal and placental function. These accelerations reflect the fetus's ability to respond to stimuli, such as movement or uterine contractions, and are generally associated with adequate oxygenation and a well-functioning placental exchange. When accelerations occur, they typically indicate that the fetal central nervous system is developed and capable of regulating heart rate effectively. This is an encouraging indication, especially during labor, suggesting that the fetus is not in distress and is receiving sufficient blood flow and oxygen through the placenta. While the context of monitoring conditions like contractions or continuous fetal monitoring can provide additional information, the fundamental assertion that accelerations represent a healthy status stands true in general obstetric assessments of fetal well-being.

2. What is the main purpose of monitoring for hypotension in a mother prior to epidural or spinal analgesia?

A. To ensure a comfortable position during labor

B. To prevent complications related to anesthesia

C. To assess fetal heart rate

D. To prepare for cesarean delivery

Monitoring for hypotension in a mother prior to epidural or spinal analgesia is essential primarily to prevent complications related to anesthesia. Epidurals and spinals can significantly affect cardiovascular dynamics by causing vasodilation and reducing venous return, which can lead to a drop in maternal blood pressure. This hypotension can result in decreased uteroplacental perfusion, which may compromise fetal oxygen delivery and lead to fetal distress. By monitoring blood pressure closely before administering these types of analgesia, healthcare providers can identify potential risks early and take preventive measures, such as administering intravenous fluids or medications, to mitigate the chances of hypotensive episodes. This proactive approach is key to ensuring both maternal safety and optimal fetal health during labor. Other options, such as ensuring comfort or preparing for a cesarean delivery, do not capture the critical risk management aspect that hypotension monitoring represents in the context of anesthesia. Assessing fetal heart rate is important in its own right, but it is not the primary concern directly tied to the risks associated with administering epidural or spinal analgesia.

SAMPLE

3. Which of the following indicates a serious side effect of magnesium sulfate?

A. Lethargy

B. Visual blurring

C. Headache

D. Weight gain

Lethargy is indeed a serious side effect of magnesium sulfate. It can indicate that the serum magnesium levels are too high, potentially leading to magnesium toxicity. Symptoms such as lethargy suggest that the central nervous system is being affected, which is a concern in patients receiving this medication for conditions like preeclampsia or eclampsia. Monitoring for lethargy is crucial, as it may require prompt adjustment of the magnesium sulfate dosage or even cessation of treatment to protect the patient from further complications related to excess magnesium in the bloodstream. Visual blurring, headache, and weight gain can occur with magnesium sulfate use, but they are generally less indicative of serious side effects. Visual blurring may be related to changes in blood pressure or fluid balance rather than direct magnesium toxicity. Headaches may also arise from various causes that are not directly linked to the magnesium levels, and weight gain could result from fluid retention, which is common in pregnant patients and may not be a direct consequence of magnesium sulfate. Therefore, while they can be concerning, they do not signal the immediate danger associated with lethargy.

4. True or False: Fetal distress is a common complication of labor.

A. True

B. False

C. Only in late labor

D. Only in early labor

Fetal distress is actually not classified as a common complication of labor. The term refers to signs that a fetus is not well, often indicated by an abnormal heart rate. While it can occur, especially under certain risk factors or complications, the majority of labors progress without significant signs of fetal distress. The occurrence of fetal distress may depend on various factors, including but not limited to maternal health, the presence of certain complications, and fetal conditions. Therefore, stating that it is a common complication would misrepresent the incidence of this condition. Additionally, fetal distress can arise at different stages of labor, rather than being specific to early or late labor, which reinforces that the claim is not true. By understanding the context and nuances, we can see why the assertion that fetal distress is a common complication does not hold up, supporting the designation of the statement as false.

SAMPLE

5. What are some adverse effects associated with oxytocin use?

A. Uterine Tachysystole

B. Increased contractions

C. Fetal hypoxia

D. Risk for uterine rupture

Oxytocin is a medication commonly used to induce or augment labor by stimulating uterine contractions. One of the notable adverse effects associated with its use is uterine tachysystole, which refers to excessive uterine activity characterized by more than five contractions in a 10-minute period. This is significant because it can lead to increased uterine pressure, which in turn can have detrimental effects on fetal oxygenation. Increased contractions are a primary effect of oxytocin; therefore, while this characteristic is indeed influential in labor management, it does not represent an adverse effect. Similarly, fetal hypoxia can occur due to insufficient oxygen supply during intense contractions but is more of a consequence of tachysystole than a direct result of oxytocin itself. The risk for uterine rupture is a serious concern but is associated with longstanding use or overdistension of the uterus, particularly in women with a previous cesarean section, rather than being a direct result of oxytocin administration alone. By understanding that uterine tachysystole is a specific adverse effect of oxytocin usage, it highlights the importance of careful monitoring during labor induction to balance the benefits of labor stimulation with the potential risks posed by excessive uterine activity.

6. Which of the following are signs of an infection in the belly button area?

A. Dryness and flakiness

B. Youthful appearance

C. Redness and swelling

D. Complaints of pain only

Redness and swelling around the belly button area are significant indicators of an infection. Infections typically provoke an inflammatory response, leading to increased blood flow and fluid accumulation, which can result in the characteristic redness and swelling. This response is the body's way of trying to fight off harmful pathogens. In contrast, dryness and flakiness, typically associated with skin conditions such as eczema or dermatitis, are not direct signs of infection. A youthful appearance may denote healthy or uninfected skin but does not correlate with infection indicators. Complaints of pain alone, while they can be associated with various conditions, do not provide the visual cues necessary to diagnose an infection effectively. Therefore, the combination of redness and swelling is a strong clinical sign that an infection may be present.

7. What does a score of 0 for activity in an Apgar score signify?

A. Active movement

B. Flexed arms and legs

C. Absent muscle tone

D. Minimal movement

A score of 0 for activity in an Apgar score indicates absent muscle tone. This assessment is part of the broader evaluation of a newborn's physical condition shortly after birth. The Apgar score uses criteria that include appearance, pulse, grimace response, activity, and respiration, with activity specifically reflecting the muscle tone and movement of the infant. When a newborn receives a score of 0 for activity, it signifies that the baby shows no movement or muscle tone, which is a critical observation that can flag potential issues requiring immediate medical attention. Adequate muscle tone is essential for normal development and indicates that the nervous system is functioning properly. On the other hand, a score of 1 would typically reflect some flexed position of the arms and legs, while a higher score would correspond to active movement. Thus, a score of 0 distinctly highlights a concerning absence of muscle tone that is crucial for evaluating the newborn's health status.

SAMPLE

8. What intervention should the nurse expect to implement for a hypoglycemic SGA newborn?

A. Administer glucose immediately regardless of feeding time.

B. Obtain a glucose level right before feeding.

C. Delay feeding until the next shift.

D. Monitor oxygen levels continuously.

Obtaining a glucose level right before feeding is an appropriate intervention for a hypoglycemic small-for-gestational-age (SGA) newborn. This is crucial because it allows the healthcare provider to assess the newborn's blood glucose status and determine whether hypoglycemia is present before proceeding with feeding. Monitoring the glucose level helps guide clinical decisions regarding feeding, as the infant may require immediate intervention, such as a glucose supplement or formula, if levels are low. By checking the glucose level prior to feeding, the nurse can ensure that the infant is stable enough to feed safely and effectively, which reduces the risk of further hypoglycemic episodes. It's also important to note that frequent monitoring of blood glucose is essential for SGA newborns, who may have difficulty maintaining stable glucose levels due to potential metabolic issues. Immediate administration of glucose without assessing the blood sugar level could lead to unnecessary interventions or over-treatment if the glucose is within acceptable limits. Delaying feeding until the next shift is counterproductive, as timely nutritional support is critical for these vulnerable infants. Continuous monitoring of oxygen levels, while important, does not directly address the immediate risk posed by hypoglycemia. Thus, the correct strategy focuses on timely assessment and appropriate feeding interventions.

9. What is the average gestational weight of a newborn born at 37 weeks?

A. 8 lbs. 6 ounces

B. 5 lbs. 13 ounces

C. 6 lbs. 1 ounce

D. 7 lbs. 3 ounces

The average gestational weight of a newborn born at 37 weeks typically falls within the range of approximately 5 pounds 8 ounces to 6 pounds 12 ounces. This weight reflects the common growth patterns of a fetus that has reached late preterm status, which is considered to be between 34 and 36 weeks. While individual weights can vary due to multiple factors, including maternal health, nutrition, and genetics, newborns delivered at 37 weeks are generally on the lower end of the average birth weight scale. The weight of 5 pounds 13 ounces aligns closely with these statistical averages for infants born just before reaching full term, which is 39 to 40 weeks. In contrast, the other options represent birth weights that are either above or below this typical average for late preterm babies. Recognizing this growth pattern helps in understanding fetal development and potential health implications at delivery.

SAMPLE

10. True or false: It is normal for the umbilical cord stump to ooze a little before it falls off.

A. True

B. False

C. Only during the first day

D. Requires immediate attention

It is normal for the umbilical cord stump to ooze a little before it falls off. After birth, the stump typically goes through a process of drying and healing, which can involve some minor oozing as the area around the cord heals. This is part of the natural process, as the body is working to seal the site where the cord was attached, and a small amount of fluid can help promote healing. While a small amount of oozing can be expected, it is important for caregivers to monitor the stump for signs of infection or excessive bleeding. If there are concerns about the amount of ooze or if it is accompanied by other concerning symptoms, it may warrant a discussion with a healthcare provider.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://cjematernity1.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE