CLINICAL CHEMISTRY II PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://clinicalchem2.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. How is CSF glucose interpreted in CNS infections?

A. Reduced in bacterial meningitis relative to serum glucose; preserved in most viral infections

B. Elevated in bacterial meningitis

C. Unchanged in bacterial meningitis but reduced in viral infections

D. Not useful for differentiating infections

2. Which of the following is a negative acute-phase reactant?

A. CRP

B. Albumin

C. Fibrinogen

D. Ferritin

3. How does the urea breath test detect H. pylori infection?

A. The patient drinks urea labeled with carbon-13 and CO2 is measured in the breath

B. The patient consumes glucose and the breath is analyzed

C. The test measures urease activity in blood

D. Urea is not involved in the test

4. Describe the main difference between serum and plasma when interpreting some laboratory results.

A. Serum contains all clotting factors; plasma does not.

B. Serum and plasma are identical in composition.

C. Serum lacks clotting factors; plasma contains anticoagulant and coagulation factors; some analytes differ slightly due to fibrinogen and sample type.

D. Plasma is derived after clotting.

5. Colorectal cancer is associated with which dietary factors?

A. Colorectal cancer

B. Prostate cancer

C. Skin cancer

D. Leukemia

SAMPLE

6. H. pylori affects approximately what proportion of people, usually asymptomatic?

A. ~10%

B. ~50%

C. ~75%

D. ~90%

7. Which product is most directly produced by amylase action on carbohydrates?

A. Fructose

B. Galactose

C. Maltose

D. Glucose

8. In pancreatitis, which metabolic abnormality related to glucose may be observed?

A. Altered glucose regulation

B. Hyperglycemia always present

C. Hypoglycemia always present

D. Normal glucose metabolism

9. What is the effect of preanalytical hemolysis on potassium and LDH measurements?

A. Hemolysis falsely lowers potassium and LDH.

B. Hemolysis has no effect on potassium or LDH measurements.

C. Hemolysis releases intracellular potassium and LDH, falsely elevating them.

D. Hemolysis only affects LDH, not potassium.

10. Which serology tests look for autoantibodies in celiac disease?

A. Anti-endomysial antibody (EMA-IgA) and Anti-tTG-IgA/tTG-IgG

B. Anti-nuclear antibodies

C. Deamidated gliadin peptide IgG only

D. Celiac disease DNA test

SAMPLE

Answers

SAMPLE

1. A
2. B
3. A
4. D
5. A
6. B
7. D
8. A
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. How is CSF glucose interpreted in CNS infections?

A. Reduced in bacterial meningitis relative to serum glucose; preserved in most viral infections

B. Elevated in bacterial meningitis

C. Unchanged in bacterial meningitis but reduced in viral infections

D. Not useful for differentiating infections

CSF glucose patterns help distinguish bacterial from viral CNS infections. In bacterial meningitis, bacteria and inflammatory cells use a lot of glucose, and inflammation can impair glucose transport from the blood into the CSF, so the CSF glucose level falls relative to serum glucose. In most viral infections, glucose consumption is less and the transport barrier remains more intact, so CSF glucose stays normal or near normal compared with serum. That combination—low CSF glucose compared with serum in bacterial meningitis and preserved CSF glucose in most viral infections—is why this option is best. Clinically, a CSF/serum glucose ratio below about 0.4–0.5 supports bacterial meningitis, while a normal ratio favors viral etiologies.

2. Which of the following is a negative acute-phase reactant?

A. CRP

B. Albumin

C. Fibrinogen

D. Ferritin

During an acute inflammatory response, the liver shifts its protein production: some proteins rise to help fight infection (positive acute-phase proteins) while others fall (negative acute-phase proteins). Albumin decreases because inflammatory cytokines, especially IL-6, downregulate its synthesis in the liver and its distribution can shift into the interstitial space due to increased vascular permeability. The other listed proteins—CRP, fibrinogen, and ferritin—are positive acute-phase proteins and rise in inflammation to support defense, clotting, and iron handling. Thus, albumin is the negative acute-phase reactant.

3. How does the urea breath test detect H. pylori infection?

A. The patient drinks urea labeled with carbon-13 and CO_2 is measured in the breath

B. The patient consumes glucose and the breath is analyzed

C. The test measures urease activity in blood

D. Urea is not involved in the test

The test relies on the stomach's urease activity to reveal an active Helicobacter pylori infection. When urea labeled with carbon-13 is swallowed, if H. pylori is present, its urease rapidly splits the urea into ammonia and labeled carbon dioxide. The labeled CO_2 is carried in the bloodstream and exhaled in the breath, so a rise in the labeled CO_2 in breath samples indicates infection. Glucose isn't used to detect H. pylori here and wouldn't reflect urease activity, measuring urease in blood isn't a breath test, and urea is indeed involved in this test.

SAMPLE

4. Describe the main difference between serum and plasma when interpreting some laboratory results.

A. Serum contains all clotting factors; plasma does not.

B. Serum and plasma are identical in composition.

C. Serum lacks clotting factors; plasma contains anticoagulant and coagulation factors; some analytes differ slightly due to fibrinogen and sample type.

D. Plasma is derived after clotting.

The main idea is that whether the blood is allowed to clot or kept from clotting determines what the liquid portion contains. Serum is what you get after blood is drawn and allowed to clot, then the clot is removed; it lacks fibrinogen and other clotting factors that are consumed during clot formation. Plasma is obtained from blood drawn with an anticoagulant and then centrifuged before clotting occurs, so it still contains fibrinogen and the full set of clotting factors. Because of this difference, some analytes can differ slightly between serum and plasma, partly due to the presence of fibrinogen and other clotting proteins in plasma and the effects of the anticoagulant used. Coagulation tests (like PT and aPTT) specifically require plasma because they depend on those clotting factors. So the key distinction is that serum lacks clotting factors because clotting has occurred, while plasma contains anticoagulant-preserved clotting factors. The idea that plasma is derived after clotting is not correct.

5. Colorectal cancer is associated with which dietary factors?

A. Colorectal cancer

B. Prostate cancer

C. Skin cancer

D. Leukemia

Dietary factors influence the risk of colorectal cancer. A pattern high in red and processed meats and low in fiber from fruits, vegetables, and whole grains is associated with a higher risk, while a diet rich in fiber and plant-based foods is linked to lower risk. Other contributors include obesity, physical inactivity, and alcohol use; calcium and vitamin D may offer some protection. Among the options, colorectal cancer is the one known to be associated with dietary patterns, whereas the other cancers listed are not primarily driven by diet in the same way.

6. H. pylori affects approximately what proportion of people, usually asymptomatic?

A. ~10%

B. ~50%

C. ~75%

D. ~90%

Helicobacter pylori infection is very common worldwide, and most people who carry it don't develop symptoms. The typical global carriage rate is about half the population, often acquired in childhood and lasting for life unless treated. Because many individuals remain asymptomatic, the proportion without symptoms is around 50%. Prevalence does vary by region and living conditions, but 50% is the standard estimate used in teaching. The smaller figure would miss many carriers, while the much larger figures would overstate how common infection is in the general population.

SAMPLE

7. Which product is most directly produced by amylase action on carbohydrates?

A. Fructose

B. Galactose

C. Maltose

D. Glucose

Amylase works by cutting the alpha-1,4 glycosidic bonds in starch, breaking it down into smaller sugars. The immediate product of this enzymatic action is maltose, a disaccharide of two glucose units (with maltotriose also formed from some fragments). Glucose itself is not released directly by amylase; it appears only after maltose is further digested by maltase (a brush-border enzyme) and other dextrins are processed. Fructose and galactose aren't direct products of amylase acting on starch either. So the sugar most directly produced by amylase action is maltose.

8. In pancreatitis, which metabolic abnormality related to glucose may be observed?

A. Altered glucose regulation

B. Hyperglycemia always present

C. Hypoglycemia always present

D. Normal glucose metabolism

Pancreatitis can injure the pancreas's endocrine tissue, which regulates blood glucose through insulin and other hormones. When this part of the organ is damaged, glucose handling becomes disrupted, leading to altered glucose regulation. This can manifest as impaired glucose tolerance or diabetes, especially with chronic pancreatic damage, and glucose levels may be high or low depending on the situation and inflammatory stress. Therefore, altered glucose regulation is the appropriate general description of the metabolic abnormality that may be observed. The notions of hyperglycemia or hypoglycemia being always present, or glucose metabolism remaining normal, do not fit the variable ways pancreatitis can affect insulin production and glucose control.

9. What is the effect of preanalytical hemolysis on potassium and LDH measurements?

A. Hemolysis falsely lowers potassium and LDH.

B. Hemolysis has no effect on potassium or LDH measurements.

C. Hemolysis releases intracellular potassium and LDH, falsely elevating them.

D. Hemolysis only affects LDH, not potassium.

When red blood cells break open before analysis, their contents spill into the serum or plasma. Potassium and LDH are both highly abundant inside RBCs, so their release into the sample makes measured levels appear higher than they truly are in the patient's circulation. Potassium is kept at a high concentration inside red cells, while the plasma potassium normally sits around 4 to 5 mEq/L. Even a modest amount of hemolysis can significantly raise the measured potassium because the intracellular potassium from ruptured cells mixes with the plasma, leading to a falsely elevated value. LDH is another enzyme concentrated inside red cells. When hemolysis occurs, LDH leaks out into the plasma, boosting the overall LDH activity measured in the sample. This can mimic conditions that cause tissue damage or disease, but in reality, it's just the artifact of cellular rupture during sample handling. Because of this, preanalytical hemolysis is a common cause of spuriously high potassium and LDH results. If a sample is hemolyzed, the results are unreliable, and repeating the draw with a non-hemolyzed sample is usually indicated.

SAMPLE

10. Which serology tests look for autoantibodies in celiac disease?

A. Anti-endomysial antibody (EMA-IgA) and Anti-tTG-IgA/tTG-IgG

B. Anti-nuclear antibodies

C. Deamidated gliadin peptide IgG only

D. Celiac disease DNA test

Serology in celiac disease centers on detecting autoantibodies against components of the gluten-triggered immune response, mainly endomysial antigens and tissue transglutaminase. The most informative combination is anti-endomysial antibodies (EMA) of the IgA class and anti-tissue transglutaminase antibodies (tTG), with the tTG tests available as IgA or IgG. EMA-IgA is highly specific for celiac disease, and tTG-IgA offers high sensitivity and specificity for routine screening; tTG-IgG is particularly helpful when IgA deficiency makes IgA-based testing unreliable. Anti-nuclear antibodies are not specific to celiac disease. Deamidated gliadin peptide antibodies can be used, especially if IgA deficiency is present, but IgG-only DGP testing is less preferred as a sole screen. A celiac disease DNA test looks at genetic predisposition, not autoantibodies. Therefore, the serology that looks for autoantibodies in celiac disease is EMA-IgA and tTG-IgA/tTG-IgG.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://clinicalchem2.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE