


CMTBC REGISTRATION PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE

<https://cmtbcregistration.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. If AROM is strong and pain-free, what does this indicate?**
 - A. No contractile tissue lesion
 - B. Muscle strain
 - C. Tendinopathy
 - D. Joint instability
- 2. Does skeletal muscle contain gap junctions?**
 - A. Yes
 - B. No
 - C. Only in certain areas
 - D. It depends on the muscle type
- 3. Which range of motion assessment is vital for knee injuries?**
 - A. Flexion and extension
 - B. Abduction and adduction
 - C. Cervical rotation
 - D. Shoulder elevation
- 4. Where do lymphatic vessels empty into the venous system?**
 - A. At the level of the aorta and iliac veins
 - B. At the subclavian vein and internal jugular vein
 - C. At the femoral vein and external jugular vein
 - D. At the pulmonary veins and superior vena cava
- 5. What is the range for ulnar deviation at the radiocarpal joint?**
 - A. 30-45 degrees
 - B. 80-90 degrees
 - C. 70-90 degrees
 - D. 15 degrees
- 6. What is the most effective treatment for lateral epicondylitis?**
 - A. Stretching of the extensors
 - B. Reciprocal inhibition to flexors
 - C. Application of heat to the elbow
 - D. Strengthening of the biceps

- 7. Which cranial nerve is the only one with a dermatome?**
- A. Facial
 - B. Optic
 - C. Trigeminal
 - D. Hypoglossal
- 8. Which pathology is characterized by Trendelenburg gait, circumduction gait, and foot equinus?**
- A. Cerebral palsy
 - B. Multiple sclerosis
 - C. Parkinson's disease
 - D. Stroke
- 9. In what action is the depressor anguli oris muscle primarily involved?**
- A. Frowning
 - B. Smiling
 - C. Eating
 - D. Speaking
- 10. During supine assessment of a patient with low back pain, increased lumbar lordosis occurs when one knee is brought to the chest. Which condition contributes to these presentations?**
- A. Sacroiliac joint dysfunction
 - B. Weakness in lumbar extensor muscles
 - C. Tight gluteus maximus in the bent leg
 - D. Iliopsoas contracture in the straight leg

Answers

SAMPLE

1. A
2. B
3. A
4. B
5. A
6. B
7. C
8. A
9. A
10. D

SAMPLE

Explanations

SAMPLE

1. If AROM is strong and pain-free, what does this indicate?

A. No contractile tissue lesion

B. Muscle strain

C. Tendinopathy

D. Joint instability

When active range of motion (AROM) is both strong and pain-free, it indicates that there is likely no contractile tissue lesion present. In this context, strong AROM suggests that the muscles are functioning well and are capable of generating appropriate force without any signs of weakness or impairment. The absence of pain further reinforces the idea that the tissue is not compromised, as pain during movement typically indicates some form of injury or lesion in the contractile structures, such as muscles or tendons. The other options consider potential issues that would demonstrate symptoms during AROM. Muscle strains typically produce weakness and pain with movement. Tendinopathy, often marked by pain especially when the tendon is loaded or stretched, would likewise show discomfort during active motion. Joint instability can lead to a lack of strength and may also cause pain or give-way during movement. Therefore, the strong and pain-free performance during AROM is a clear indicator that the contractile structures are intact and functionally sound.

2. Does skeletal muscle contain gap junctions?

A. Yes

B. No

C. Only in certain areas

D. It depends on the muscle type

Skeletal muscle is characterized by its striated appearance and is primarily under voluntary control. A distinctive feature of skeletal muscle tissue is that it is composed of multinucleated fibers that do not typically exhibit gap junctions. Gap junctions are specialized intercellular connections that facilitate communication and transfer of ions and small molecules between adjacent cells. In the case of skeletal muscle, each muscle fiber operates independently of neighboring fibers, primarily relying on the neuromuscular junction for stimulation and communication from the nervous system. Unlike cardiac muscle, which utilizes gap junctions for coordinated contractions, skeletal muscle relies on a different mechanism for contraction, dependent on motor neuron activation. Thus, the absence of gap junctions in skeletal muscle cells reflects the unique functional requirements of this type of muscle, which necessitates direct and voluntary control rather than synchronized activity.

3. Which range of motion assessment is vital for knee injuries?

A. Flexion and extension

B. Abduction and adduction

C. Cervical rotation

D. Shoulder elevation

The assessment of flexion and extension is crucial for knee injuries because these movements directly relate to the primary functions of the knee joint. The knee is a hinge joint, primarily allowing for bending (flexion) and straightening (extension) motions. Evaluating these ranges of motion helps to identify impairments or limitations in the joint's functionality, which is typically a significant concern following a knee injury. Assessing flexion provides insights into the ability to bend the knee, which is vital for activities like walking, running, and sitting. Extension assessment reveals how well the knee can be straightened, which is essential for stability and support while standing or engaging in physical activities. Understanding the degree of flexion and extension can also indicate the severity of an injury, guide rehabilitation protocols, and determine readiness for return to activities. In contrast, the other options relate to different joints or movement patterns that are not as directly relevant to knee injuries.

4. Where do lymphatic vessels empty into the venous system?

- A. At the level of the aorta and iliac veins
- B. At the subclavian vein and internal jugular vein**
- C. At the femoral vein and external jugular vein
- D. At the pulmonary veins and superior vena cava

Lymphatic vessels play a crucial role in the body's immune system and fluid balance by transporting lymph, a fluid containing infection-fighting white blood cells, throughout the body. The lymphatic system ultimately drains into the venous system at specific junctions. The correct answer indicates that lymphatic vessels empty into the venous system at the subclavian vein and internal jugular vein. Specifically, lymph from the right side of the head, neck, arm, and upper torso typically drains into the right lymphatic duct, which then merges with the subclavian vein. Lymph from the rest of the body, including the left arm and side of the head and neck, drains through the thoracic duct, which empties into the left subclavian vein where it meets the internal jugular vein. This is the primary point of convergence for the lymphatic circulation into the systemic circulation, making the connection with the major veins critical for returning fluid to the bloodstream and maintaining homeostasis. Other options refer to locations that are not physiologically relevant points for lymphatic drainage. The aorta and iliac veins are arteries and veins located much further down in the circulatory system; they do not play a direct role in lymphatic drainage. Similarly, the fem

5. What is the range for ulnar deviation at the radiocarpal joint?

- A. 30-45 degrees**
- B. 80-90 degrees
- C. 70-90 degrees
- D. 15 degrees

The range for ulnar deviation at the radiocarpal joint is typically around 30 to 45 degrees. This movement refers to the wrist bending toward the ulnar side, which can be observed when the hand is positioned palm down and the little finger moves closer to the forearm. This range is an important aspect of wrist mobility and is clinically significant for assessing wrist function and diagnosing potential injuries or conditions that might restrict movement. It's crucial for healthcare professionals to know this range to properly evaluate a patient's wrist function, as deviations from this norm may indicate underlying issues such as ligament injury, arthritis, or other musculoskeletal conditions. The other ranges presented in the options are larger than what is typically expected for ulnar deviation at the radiocarpal joint. Recognizing the standard range helps ensure that proper assessments and interventions can be made based on accurate anatomical and physiological knowledge.

6. What is the most effective treatment for lateral epicondylitis?

- A. Stretching of the extensors
- B. Reciprocal inhibition to flexors**
- C. Application of heat to the elbow
- D. Strengthening of the biceps

Reciprocal inhibition is a technique that involves the relaxation of one muscle group when the opposing muscle group contracts. In the case of lateral epicondylitis, which is often associated with overuse of the extensor muscles of the forearm, this approach can be quite effective. By facilitating the relaxation of the flexor muscles while engaging the extensors, the treatment not only helps alleviate pain but also encourages balance between these muscle groups. This balance can lead to improved functional movement and reduced stress on the affected area, promoting healing. While stretching the extensors may provide some benefits by improving flexibility, it doesn't directly address the underlying issues of muscle imbalance or overactivity associated with lateral epicondylitis. Application of heat can be soothing and may help improve blood flow temporarily, but it doesn't actively resolve the muscle dysfunction or pain. Strengthening the biceps, while generally beneficial for overall upper extremity strength, is not specifically targeted to the condition affecting the extensors at the lateral elbow. Therefore, reciprocal inhibition for the flexors directly addresses the muscular dynamics involved in lateral epicondylitis, making it a more effective treatment choice.

7. Which cranial nerve is the only one with a dermatome?

- A. Facial
- B. Optic
- C. Trigeminal**
- D. Hypoglossal

The correct answer is the Trigeminal nerve. This cranial nerve is unique among the cranial nerves because it has a clear dermatome distribution. A dermatome is a specific area of skin that is mainly supplied by a single spinal nerve root. The Trigeminal nerve is divided into three main branches: the ophthalmic, maxillary, and mandibular branches, each being responsible for sensory innervation to different regions of the face. This sensory supply can be mapped, similar to how dermatomes from spinal nerves are mapped, which is why the Trigeminal nerve stands out in the context of dermatomes. The other cranial nerves mentioned do not have this dermatome-like distribution. The Facial nerve primarily serves motor functions to the muscles of facial expression and has limited sensory components focused on taste. The Optic nerve is primarily responsible for vision with no sensory distribution to skin areas. The Hypoglossal nerve is focused on the motor control of the tongue without any sensory function. Thus, the Trigeminal nerve is distinctive for its role in providing sensory innervation in a manner akin to dermatomes.

8. Which pathology is characterized by Trendelenburg gait, circumduction gait, and foot equinus?

- A. Cerebral palsy**
- B. Multiple sclerosis
- C. Parkinson's disease
- D. Stroke

The correct answer focuses on cerebral palsy, which is indeed characterized by a range of atypical movement patterns, including Trendelenburg gait, circumduction gait, and foot equinus. In cerebral palsy, individuals may exhibit muscle tone abnormalities, which can result in difficulties with movement control and coordination. The Trendelenburg gait occurs due to weakness in the hip abductors, particularly the gluteus medius, leading to a characteristic pelvic drop on the opposite side when walking. The circumduction gait is often adopted as a compensatory strategy, where the affected leg is swung outwards during the swing phase to clear the ground, typically seen in individuals with spasticity or weakness in the lower limbs. Foot equinus refers to a position where the toes point down, which can affect balance and propulsion in gait. These gait abnormalities are a result of the motor impairments associated with cerebral palsy, where the brain's control over muscles and movement is disrupted during development. This condition highlights the challenges that can arise in achieving normal gait mechanics and the adaptations that the body may develop to cope with those challenges.

9. In what action is the depressor anguli oris muscle primarily involved?

- A. Frowning**
- B. Smiling
- C. Eating
- D. Speaking

The depressor anguli oris muscle plays a key role in facial expressions, particularly in the action of frowning. This muscle is located at the corners of the mouth and is responsible for pulling the corners of the mouth downwards. When this muscle contracts, it creates a downward movement that typically indicates sadness, discontent, or displeasure, which is commonly associated with frowning. In contrast, the other actions listed rely on different muscles: smiling involves the zygomaticus major and other muscles that elevate the corners of the mouth, eating engages various muscles for chewing and positioning food, and speaking involves a coordinated effort of several facial muscles for the movement of the lips and mouth. Thus, the depressor anguli oris is specifically linked to the expression of a frown, making it the correct choice in this context.

10. During supine assessment of a patient with low back pain, increased lumbar lordosis occurs when one knee is brought to the chest. Which condition contributes to these presentations?

- A. Sacroiliac joint dysfunction
- B. Weakness in lumbar extensor muscles
- C. Tight gluteus maximus in the bent leg
- D. Iliopsoas contracture in the straight leg**

In the scenario presented, the increased lumbar lordosis when bringing one knee to the chest is primarily influenced by the condition of an iliopsoas contracture in the straight leg. The iliopsoas is a major hip flexor that plays a crucial role in stabilizing the pelvis and lumbar spine. When the iliopsoas is contracted or tight, it can prevent proper positioning of the pelvis and lumbar spine during movement and positioning. As one knee is drawn toward the chest, if the other leg remains straight and the iliopsoas is contracted, it can create a posterior pelvic tilt on that side, which subsequently increases the lumbar lordosis. This results in an anterior tilt of the pelvis and emphasizes the natural curve of the lumbar spine, leading to increased lordosis in the supine position. Understanding the mechanics of how the iliopsoas interacts with the lumbar spine is critical for assessing low back pain and determining the underlying contributors to altered spinal alignment during movement. This highlights the importance of evaluating muscle flexibility and strength in the context of low back pain assessment.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://cmtbcregistration.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE