COLLEGE MATH PLACEMENT PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://collegemath.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the solution for t in the inequality 0.1 − 2t ≥ 0.7?

A. t ≥ −0.4

B. t ≤ −0.4

C. t ≤ −0.3

D. t ≥ −0.3

2. What is the simplified form of the expression 4x(2y) + 3y(2−x)?

A. 5xy + 6y

B. 8xy + 6y − x

C. 11xy + 6y

D. 8xy + 6y − 3x

3. How is Tuesday's chicken production expressed if it is described as ten less than twice the eggs produced on Monday, where m represents Monday's eggs?

A. 2m − 10

B. 2 + m − 10

C. 2m + 10

D. 10 − 2m

4. In set notation, how do you define the empty set?

A. {}

B.

C. N/A

D. NULL

5. What is the derivative of f(x) = x^3?

A. 2x^2

B. 3x^2

C. x^2

D. 2x

SAMPLE

6. What is the graphical representation of y = mx + b?

A. Cubic equation

B. Exponential function

C. Linear equation

D. Quadratic equation

7. What is the result of evaluating the integral ∫(2x)dx?

A. $x + C$

B. $x^2 + C$

C. $2x^2 + C$

D. $3x^2 + C$

8. If sin(θ) = 0.5, what is the angle θ in degrees?

A. 15 degrees

B. 30 degrees

C. 45 degrees

D. 60 degrees

9. What is the derivative of f(x) = sin(x)?

A. sin(x)

B. cos(x)

C. -sin(x)

D. -cos(x)

10. What is the sum of the angles in a triangle?

A. 180 degrees

B. 90 degrees

C. 360 degrees

D. 270 degrees

SAMPLE

Answers

SAMPLE

1. C
2. A
3. A
4. B
5. B
6. C
7. B
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What is the solution for t in the inequality 0.1 − 2t ≥ 0.7?

A. t ≥ −0.4

B. t ≤ −0.4

C. t ≤ −0.3

D. t ≥ −0.3

To solve the inequality \(0.1 - 2t \geq 0.7\), we start by isolating the term with \(t\). First, we can subtract \(0.1\) from both sides: \[-2t \geq 0.7 - 0.1 \] This simplifies to: \[-2t \geq 0.6 \] Next, we divide both sides by \(-2\). Remember that dividing or multiplying both sides of an inequality by a negative number reverses the inequality sign: \[t \leq \frac{0.6}{-2} \] Calculating that gives: \[t \leq -0.3 \] Thus, the solution indicates that \(t\) must be less than or equal to \(-0.3\), confirming that the correct choice is one that states \(t\) is less than or equal to \(-0.3\). The reasoning leading to this conclusion reflects an accurate handling of inequalities, particularly the importance of reversing the inequality when dividing by a negative number.

2. What is the simplified form of the expression 4x(2y) + 3y(2−x)?

A. 5xy + 6y

B. 8xy + 6y − x

C. 11xy + 6y

D. 8xy + 6y − 3x

To simplify the expression 4x(2y) + 3y(2−x), we start by distributing the terms in each part of the expression. First, look at the term 4x(2y). When simplifying this, we multiply 4x by 2y, which gives us: 4x(2y) = 8xy. Next, we simplify the term 3y(2 - x). Here, we distribute 3y across both terms within the parentheses: 3y(2) - 3y(x) = 6y - 3xy. Now, we can combine the simplified results from both parts of the expression: 8xy + (6y - 3xy). When combining like terms, specifically the xy terms, we notice that: 8xy - 3xy = 5xy. Putting everything together gives us: 5xy + 6y. This matches option A perfectly. By breaking down the expression carefully and combining like terms, we see that the correct simplification leads us to the answer of 5xy + 6y.

3. How is Tuesday's chicken production expressed if it is described as ten less than twice the eggs produced on Monday, where m represents Monday's eggs?

A. 2m − 10

B. 2 + m − 10

C. 2m + 10

D. 10 − 2m

The correct representation of Tuesday's chicken production comes from translating the verbal description into a mathematical expression based on Monday's egg production, represented as \(m \). The term "twice the eggs produced on Monday" means you take \(m \) (the number of eggs from Monday) and multiply it by 2, which gives you \(2m \). The phrase "ten less than" indicates that we need to subtract 10 from the result we obtained from that multiplication. Therefore, we take \(2m \) and subtract 10 from it. Putting these pieces together, the expression for Tuesday's chicken production becomes: \[2m - 10 \] This aligns perfectly with the first choice. The other choices incorporate different arithmetic operations or do not adhere to the original description accurately, which can lead to confusion about the correct relationship between Monday's egg production and Tuesday's chicken production.

SAMPLE

4. In set notation, how do you define the empty set?

A. {}

B.

C. N/A

D. NULL

The empty set is often best represented by the symbol in set notation. This symbol clearly signifies a set that contains no elements at all. It is a universally accepted notation in mathematics, making it an efficient way to denote the absence of members in a set. While the curly braces {} can also signify an empty set, they are technically a notation that can imply an empty collection rather than specifically the set itself. The use of as a representation is more precise and widely recognized. The terms N/A and NULL do not correspond to standard mathematical terminology regarding sets; these terms may represent non-applicable states or null values in other contexts, but they do not denote the empty set in formal set theory. Thus, choosing clearly identifies the empty set in a way that is precise and consistent with mathematical conventions.

5. What is the derivative of f(x) = x^3?

A. 2x^2

B. 3x^2

C. x^2

D. 2x

The derivative of a function gives the rate at which the function's output changes with respect to its input. For the function \(f(x) = x^3 \), we can apply the power rule of differentiation. The power rule states that if \(f(x) = x^n \), then the derivative \(f'(x) \) is calculated as \(f'(x) = n \cdot x^{n-1} \). In this case, applying the power rule to \(f(x) = x^3 \): 1. Identify \(n \): Here, \(n = 3 \). 2. Apply the power rule: \(f'(x) = 3 \cdot x^{3-1} = 3 \cdot x^2 \). Thus, the derivative of \(f(x) = x^3 \) is \(3x^2 \). This is why the correct answer is \(3x^2 \). The other options do not align with the application of the power rule for this specific exponent.

6. What is the graphical representation of y = mx + b?

A. Cubic equation

B. Exponential function

C. Linear equation

D. Quadratic equation

The equation \(y = mx + b \) represents a linear equation, where \(m \) is the slope of the line and \(b \) is the y-intercept. In this format, any value of \(x \) can generate a corresponding \(y \) value, resulting in a straight line when graphed on a coordinate plane. The slope \(m \) determines the angle at which this line rises or falls, while the y-intercept \(b \) indicates the point where the line crosses the y-axis. This simplicity in structure and visual representation is what distinguishes linear equations from other types of equations, such as cubic, exponential, or quadratic equations, all of which have different characteristics and curves in their graphical representations. Thus, the correct identification of this equation as representing a linear relationship is based on its form and the nature of its graph.

SAMPLE

7. What is the result of evaluating the integral ∫(2x)dx?

A. x + C

B. $x^2 + C$

C. $2x^2 + C$

D. $3x^2 + C$

*To evaluate the integral ∫(2x)dx, we apply the power rule for integration. The power rule states that the integral of x raised to the power n is (x^(n+1))/(n+1) plus a constant of integration, C. In the case of the integrand 2x, we can consider this as 2 times x raised to the power of 1. According to the power rule: 1. Increase the exponent by one: from 1 to 2. 2. Divide by the new exponent: divide by 2. So, the integral can be computed as follows: ∫(2x)dx = 2 * (x^(1+1))/(1+1) + C = 2 * (x²/2) + C = x² + C. Thus, when you perform the integration correctly, you arrive at x² + C, which confirms that this option is indeed the correct result for the integral ∫(2x)dx. The constant C represents any constant value that could be added because the integral represents an entire family of functions.*

8. If sin(θ) = 0.5, what is the angle θ in degrees?

A. 15 degrees

B. 30 degrees

C. 45 degrees

D. 60 degrees

To determine the angle \(\theta \) when \(\sin(\theta) = 0.5 \), we can refer to the unit circle and the values of sine for commonly known angles. The sine function, \(\sin(\theta) \), represents the ratio of the opposite side to the hypotenuse in a right triangle. From the unit circle, we learn that \(\sin(30^\circ) \) is equal to \(0.5 \). Therefore, when \(\sin(\theta) = 0.5 \), one valid solution for \(\theta \) is indeed \(30^\circ \). Additionally, the sine function is positive in the first and second quadrants. Thus, beyond \(30^\circ \), there is also an angle in the second quadrant where \(\sin(180^\circ - 30^\circ) = 0.5\), which gives us \(150^\circ \) as another solution. However, if we are only focusing on angles from \(0^\circ \) to \(180^\circ \) for typical placements in such problems, the immediate and simplest answer remains \(30^\circ \). Hence, the value of

9. What is the derivative of f(x) = sin(x)?

A. sin(x)

B. cos(x)

C. -sin(x)

D. -cos(x)

The derivative of the function \(f(x) = \sin(x) \) is determined using the fundamental rules of differentiation in calculus. Specifically, it is known that the derivative of the sine function is the cosine function. This relationship comes from the limit definitions and the fundamental properties of trigonometric functions. When differentiating \(\sin(x) \) with respect to \(x \), you arrive at \(\cos(x) \). This reflects how the slope of the sine curve varies as \(x \) changes. At any point on the sine curve, the rate of change (or slope) is equal to the value of the cosine function at that point. Thus, for this particular question, the derivative of \(f(x) = \sin(x) \) confirms that the correct answer is indeed the cosine function, \(\cos(x) \).

SAMPLE

10. What is the sum of the angles in a triangle?

A. 180 degrees

B. 90 degrees

C. 360 degrees

D. 270 degrees

In any triangle, the sum of the interior angles is always equal to 180 degrees. This fundamental property is true for all types of triangles, whether they are scalene, isosceles, or equilateral. To visualize this, you can imagine a triangle drawn on a flat surface. If you measure the three angles formed at each vertex of the triangle and add them together, you will always find that they total 180 degrees. This property is a key aspect of Euclidean geometry and is foundational for many other concepts in geometry and trigonometry. The other choices represent incorrect sums for triangles. For instance, a right angle measures 90 degrees, which is insufficient for a three-angle system. A sum of 360 degrees applies to a full circle, while 270 degrees does not correspond to any basic properties of triangle geometry. Thus, the sum of the angles being 180 degrees is the only correct answer when discussing the angles in a triangle.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://collegemath.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE