DAVITA CERTIFIED CLINICAL HEMODIALYSIS TECHNICIAN (CCHT) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://davitaccht.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Acceptable interdialytic weight gains are:

A. 3 - 5% of EDW

B. 6 - 7% of EDW

C. 8 - 9% of EDW

D. 10 - 11% of EDW

2. What should a technician's immediate action be if foam is visible in the venous drip chamber during hemodialysis?

A. Assess needle placement

B. Clamp the venous line

C. Mute the alarm

D. Return the blood

3. Which factor is likely to contribute the least to post-transplant infections?

A. Immunosuppressive therapy

B. Poor nutrient absorption

C. Hygiene practices

D. Medical complications during surgery

4. What principle does the water softener operate on?

A. Osmosis

B. Filtration

C. Ion exchange

D. Electrodialysis

5. What is the most common complication in peritoneal dialysis?

A. Pain on inflow

B. Pericarditis

C. Peritonitis

D. Protein loss

SAMPLE

6. Which of the following counts as a systemic symptom of an allergic reaction?

A. Hives.

B. Chills.

C. Dizziness.

D. Nausea.

7. Why is it important to monitor bicarbonate levels in dialysate?

A. To prevent infection

B. To correct metabolic acidosis

C. To maintain osmotic balance

D. To ensure safe temperature regulation

8. What is essential for ensuring the accuracy of lab results?

A. Drawing anytime during treatment.

B. Drawing at the right time, site, and order.

C. Only before treatment.

D. Drawing after giving heparin.

9. How should you correct an incorrect entry on a patient's chart?

A. Circle it and write above

B. Draw one line through it and initial it

C. Draw one line, date it, sign it and note "error"

D. White out and write over the old entry

10. How does decreasing the patient's prescribed blood flow rate affect the rate of diffusion during dialysis?

A. It decreases diffusion.

B. It increases diffusion.

C. It has no effect on diffusion.

D. It enhances osmotic balance.

SAMPLE

Answers

SAMPLE

1. A
2. B
3. C
4. C
5. C
6. A
7. B
8. B
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. Acceptable interdialytic weight gains are:

A. 3 - 5% of EDW

B. 6 - 7% of EDW

C. 8 - 9% of EDW

D. 10 - 11% of EDW

The acceptable range for interdialytic weight gains is typically considered to be between 3% to 5% of the patient's estimated dry weight (EDW). This percentage is crucial for ensuring that fluid management during hemodialysis is effective, as excessive weight gain can indicate fluid overload, potentially leading to complications such as hypertension, cardiovascular issues, and difficulty in dialysis adequacy. Keeping weight gains within this range helps maintain optimal health and reduces the risk of complications related to dialysis treatment. Weight gains beyond this range, such as 6% to 7% or higher options, may not be ideal, as they can increase the burden on the heart and other organs when excess fluid is present during dialysis sessions. Therefore, adhering to the 3% to 5% limit is an important practice in the care of patients undergoing hemodialysis.

2. What should a technician's immediate action be if foam is visible in the venous drip chamber during hemodialysis?

A. Assess needle placement

B. Clamp the venous line

C. Mute the alarm

D. Return the blood

If foam is visible in the venous drip chamber during hemodialysis, clamping the venous line is the appropriate immediate action. The presence of foam can indicate a potential air bubble or a problem with the blood return process, which could lead to air embolism if not addressed quickly. Clamping the venous line stops blood flow and prevents any further air from entering the patient's bloodstream. This action ensures the patient's safety while further assessment can be conducted. Once the line is clamped, the technician can then proceed to investigate the cause of the foam, such as checking for air in the system, inspecting the connections, or evaluating the adequacy of the blood return. Other actions, such as assessing needle placement or returning the blood, might be relevant in different contexts but are not the immediate procedures in response to visible foam. Muting the alarm would also be inappropriate as it might prevent necessary alerts from being heard, which could jeopardize patient safety.

3. Which factor is likely to contribute the least to post-transplant infections?

A. Immunosuppressive therapy

B. Poor nutrient absorption

C. Hygiene practices

D. Medical complications during surgery

Hygiene practices significantly reduce the risk of infections in transplant patients. Proper hygiene protocols, such as hand washing, sterile techniques during procedures, and maintaining a clean environment, are fundamental measures in preventing the introduction of pathogens that could lead to infections. High standards of hygiene help protect immunocompromised patients, particularly those on immunosuppressive therapy, which means that good hygiene practices will likely contribute the least to the risk of post-transplant infections compared to other factors like immunosuppressive therapy, which directly weakens the immune system, or medical complications during surgery, which can introduce bacteria or result in wounds that are susceptible to infection. While poor nutrient absorption might play a role in overall health, it is hygiene that acts as a primary defense against the incidence of infections post-transplant.

EXAMPLE

4. What principle does the water softener operate on?

A. Osmosis

B. Filtration

C. Ion exchange

D. Electrodialysis

The water softener operates on the principle of ion exchange. In the context of water treatment, this process involves replacing hard ions, such as calcium and magnesium, which contribute to water hardness, with sodium ions. The softening process typically occurs in a resin bed within the softener unit, where hard water passes through and the hard ions are attracted to the resin and exchanged for sodium ions. Ion exchange is crucial because hard water can lead to scale buildup in pipes and appliances, reducing their efficiency and lifespan. By utilizing this method, water softeners effectively mitigate these issues, resulting in "softer" water that's better for household use and appliances. While options like osmosis, filtration, and electrodialysis are important processes in water treatment, they do not specifically address the unique function of removing hardness from water through the targeted exchange of ions.

5. What is the most common complication in peritoneal dialysis?

A. Pain on inflow

B. Pericarditis

C. Peritonitis

D. Protein loss

Peritonitis is recognized as the most common complication in peritoneal dialysis. This condition arises when the peritoneal cavity becomes infected, often due to the introduction of bacteria during the dialysis process. This infection can result from improper techniques during catheter insertion or care, or from contamination of the dialysis solution. When a patient is undergoing peritoneal dialysis, the peritoneal membrane acts as a filter for waste products and excess fluid. However, it is also susceptible to infections. Symptoms of peritonitis can include abdominal pain, fever, nausea, and changes in dialysis fluid clarity, which can indicate an ongoing infection requiring immediate medical attention. Timely recognition and treatment of peritonitis are crucial, as it can lead to serious complications if not addressed promptly. Regular monitoring and following strict hygiene protocols during the dialysis process are essential preventive measures to reduce the risk of developing this condition. Understanding the significance of peritonitis in the context of peritoneal dialysis allows healthcare providers to emphasize preventive strategies and provide proactive care for patients undergoing this therapy.

SAMPLE

6. Which of the following counts as a systemic symptom of an allergic reaction?

A. Hives.

B. Chills.

C. Dizziness.

D. Nausea.

Hives, also known as urticaria, are raised, itchy welts on the skin that occur as a result of allergic reactions. While they are often localized, they can indicate a systemic allergic response within the body. When an allergic reaction occurs, the immune system releases histamine and other chemicals, which can lead to the development of hives. This response can be part of a broader systemic reaction, especially if it progresses to more severe manifestations such as anaphylaxis, which affects multiple systems in the body. In the context of allergic reactions, systemic symptoms are those that indicate the involvement of multiple organ systems rather than just localized effects. Hives, while visible on the skin, serve as an indicator of this systemic involvement because they can show that the body is reacting to an allergen beyond just a localized response. Other options, like chills, dizziness, and nausea, can certainly arise from various causes, including allergic reactions, but they are not specific indicators of an allergy in the way that hives are. Each of these symptoms can be linked to different physiological or medical conditions, and their presence alone does not definitively confirm an allergic reaction. Instead, they are more generalized responses and might indicate reactions that are not primarily allergic in nature.

7. Why is it important to monitor bicarbonate levels in dialysate?

A. To prevent infection

B. To correct metabolic acidosis

C. To maintain osmotic balance

D. To ensure safe temperature regulation

Monitoring bicarbonate levels in dialysate is crucial for correcting metabolic acidosis, which is a common condition in patients with chronic kidney disease. As the kidneys lose their ability to excrete acid and retain bicarbonate, metabolic acidosis can develop, leading to various complications such as bone disease, muscle wasting, and cardiovascular problems. During hemodialysis, bicarbonate is typically added to the dialysate to help buffer the acidity of the blood, providing the necessary support to normalize pH levels. Keeping the bicarbonate concentration within appropriate limits ensures that the dialysis treatment effectively addresses the acid-base imbalance, thereby helping to improve the overall health and well-being of the patient. This focus on bicarbonate levels is not related to preventing infection, maintaining osmotic balance, or ensuring safe temperature regulation, which are also important factors in dialysis but do not directly address the issue of metabolic acidosis. Therefore, the emphasis on correcting metabolic acidosis highlights why monitoring bicarbonate levels is specifically critical in the treatment of patients undergoing hemodialysis.

SAMPLE

8. What is essential for ensuring the accuracy of lab results?

A. Drawing anytime during treatment.

B. Drawing at the right time, site, and order.

C. Only before treatment.

D. Drawing after giving heparin.

Ensuring the accuracy of laboratory results is crucial in any clinical setting, especially in hemodialysis. Drawing blood samples at the right time, from the correct site, and in the appropriate order is fundamental in obtaining reliable results. Timing is critical because certain laboratory values can fluctuate based on the patient's physiological state—considerations like the timing of medication administration or dietary intake can heavily influence results. The site is equally important; for example, venous blood samples should be taken from a site that is not altered by prior infusions or hemodialysis treatment to avoid contamination. Additionally, drawing samples in the correct order prevents cross-contamination of tubes that can result from drawing blood out of the appropriate sequence. This meticulous approach ultimately minimizes errors and provides clinicians with the accurate data needed to make informed decisions regarding patient care.

9. How should you correct an incorrect entry on a patient's chart?

A. Circle it and write above

B. Draw one line through it and initial it

C. Draw one line, date it, sign it and note "error"

D. White out and write over the old entry

When correcting an incorrect entry in a patient's chart, it's essential to maintain the integrity and accuracy of the medical record, ensuring that all corrections are clearly documented. The correct method involves drawing a single line through the erroneous entry, dating it, signing it, and adding a note that states "error." This approach preserves the original entry while providing a clear indication of the change made. It allows for transparency about the mistake and ensures that anyone reviewing the chart can see what was originally documented as well as the correction. This method aligns with best practices in medical record keeping, which emphasize maintaining an auditable record and preventing any potential confusion. By providing your signature and date, you are also affirming the validity of the correction, which is critical in clinical settings for accountability and compliance with regulatory standards.

10. How does decreasing the patient's prescribed blood flow rate affect the rate of diffusion during dialysis?

A. It decreases diffusion.

B. It increases diffusion.

C. It has no effect on diffusion.

D. It enhances osmotic balance.

Decreasing the patient's prescribed blood flow rate fundamentally affects the rate of diffusion during dialysis by reducing the movement of blood through the dialyzer. In dialysis, diffusion is the process through which solutes (like urea and creatinine) move from an area of higher concentration in the blood to an area of lower concentration in the dialysate. When the blood flow rate is lower, it results in a decreased transport of these solutes past the semipermeable membrane into the dialysate. The slower blood flow means that there is less opportunity for solutes to diffuse out of the blood, effectively resulting in a slower rate of diffusion. This can compromise the efficiency of the dialysis treatment and may lead to inadequate removal of toxins from the bloodstream. In contrast, a higher blood flow rate facilitates a greater gradient for diffusion because more solutes are presented to the membrane more quickly, allowing for more effective removal. Therefore, adjusting the blood flow rate is a critical factor in optimizing diffusion rates during hemodialysis.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://davitaccht.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE