DIFFERENTIAL DIAGNOSIS AND MANAGEMENT OF COMMON ACUTE EYE AND MUSCULOSKELETAL CONDITIONS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://
diffdiagmgmtacuteeyemskconditions.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What are the key symptoms of a hordeolum?

A. Red, painful bump along the eyelid margin, swelling, crusting, tearing, sensitivity to light.

B. Painless swelling of the eyelid.

C. Gradual vision loss.

D. Itchy but non-inflammatory sensation.

2. Which knee examination maneuver is used to detect meniscal tears?

A. Pivot-shift test

B. Lachman test

C. Apley compression test

D. Valgus stress test

3. In a patient with suspected orbital involvement after trauma, which finding would most strongly suggest orbital cellulitis rather than a periocular infection?

A. Pain with eye movements and restricted extraocular movements.

B. Crusting and itching of eyelid margins.

C. Mucopurulent discharge without pain on movement.

D. Normal vision with mild lid swelling.

4. What does an endocrine evaluation identify in hip conditions like SCFE?

A. Bacterial infection

B. Normal aging changes

C. Vascular problems

D. Underlying hormonal or metabolic abnormalities.

5. What is a hordeolum (stye)?

A. A chronic dry eye condition.

B. A allergic reaction of the eyelid.

C. A viral conjunctival infection.

D. An infection and inflammation of the eyelid, commonly referring to an infected eyelash follicle.

SAMPLE

6. What is the management step for retinal detachment?

A. Wait and watch

B. Emergency ophthalmology referral

C. Home rest and analgesia

D. Oral antibiotics only

7. What are the symptoms of iritis?

A. Absent vision and eye movement.

B. Redness, pain, photophobia, blurred vision.

C. Diffuse redness with discharge.

D. Severe itching without pain.

8. What is the typical examination finding for a chalazion?

A. Painless lump on eyelid that is fast-growing.

B. Nontender lump that is slow-growing.

C. Red, tender swelling with crusting.

D. It is typically associated with fever.

9. What are the symptoms of De Quervain's Tenosynovitis?

A. Painful swelling on the dorsal wrist

B. Inflammation of tendons on the side of the wrist at the base of the thumb.

C. Numbness in the little finger

D. Joint locking of the elbow

10. What are common risk factors for developing a chalazion?

A. Diabetes, hypertension, obesity.

B. Acne rosacea, chronic blepharitis, seborrhea, tuberculosis, viral infection.

C. Myopia and hyperopia.

D. Only aging-related changes.

SAMPLE

Answers

SAMPLE

1. A
2. C
3. A
4. D
5. D
6. B
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What are the key symptoms of a hordeolum?

A. Red, painful bump along the eyelid margin, swelling, crusting, tearing, sensitivity to light.

B. Painless swelling of the eyelid.

C. Gradual vision loss.

D. Itchy but non-inflammatory sensation.

An acute eyelid infection of the glands at the lash line presents as a red, painful lump along the eyelid margin with swelling, crusting, tearing, and sometimes light sensitivity. This combination of sudden onset, localized tenderness, erythema at the lid margin, crusting at the lashes, and tearing is classic for a hordeolum (stye). The other descriptions fit different conditions: a painless lid swelling usually points to a chalazion, a gradual vision loss suggests a problem farther back in the eye or optic pathway, and an itchy but non-inflammatory sensation aligns more with allergic conjunctivitis or dry eye rather than an acute bacterial eyelid infection.

2. Which knee examination maneuver is used to detect meniscal tears?

A. Pivot-shift test

B. Lachman test

C. Apley compression test

D. Valgus stress test

Meniscal tears respond to loading and twisting of the knee, so a test that specifically stresses the meniscus is most informative. The Apley compression test does this by flexing the knee to about 90 degrees, pressing the tibia downward through the heel, and rotating the tibia internally and externally. Reproducing pain, a joint-line click, or locking during the compression suggests a torn meniscus. If distraction of the tibia relieves pain, that points away from a meniscal tear and toward a ligamentous injury. Other maneuvers target ligaments—such as the pivot-shift and Lachman for the ACL, and the valgus stress test for the MCL—so they're less specific for meniscal pathology.

3. In a patient with suspected orbital involvement after trauma, which finding would most strongly suggest orbital cellulitis rather than a periocular infection?

A. Pain with eye movements and restricted extraocular movements.

B. Crusting and itching of eyelid margins.

C. Mucopurulent discharge without pain on movement.

D. Normal vision with mild lid swelling.

The distinguishing sign is involvement of the orbit itself, which makes eye movements painful and limited. When the infection extends behind the orbital septum to affect the extraocular muscles and orbital tissues, moving the eye becomes painful and the range of motion can be restricted. This contrasts with periocular (preseptal) infections that are confined to the eyelids and surrounding skin, where eye movements remain full and pain on movement is absent. In practice, this sign points to orbital cellulitis, a medical emergency requiring urgent imaging to assess extent and prompt IV antibiotics to prevent vision loss or intracranial spread. Crusting and itching of the eyelid margins point to blepharitis or conjunctival irritation rather than orbital involvement. Mucopurulent discharge without pain on movement suggests conjunctivitis rather than orbital cellulitis, and normal vision with mild lid swelling is more compatible with superficial eyelid/skin involvement or uncomplicated conjunctival irritation rather than orbital disease.

SAMPLE

4. What does an endocrine evaluation identify in hip conditions like SCFE?

A. Bacterial infection

B. Normal aging changes

C. Vascular problems

D. Underlying hormonal or metabolic abnormalities.

In SCFE, the slipping of the proximal femoral epiphysis often results from systemic factors that weaken the growth plate, especially hormonal or metabolic issues. An endocrine evaluation is specifically used to uncover underlying hormonal or metabolic abnormalities, such as thyroid dysfunction or growth hormone axis problems, that may predispose to slipping or affect healing and recurrence risk. Identifying these factors matters because it guides both the treatment of the slip and the management of the underlying condition, potentially altering prognosis. Bacterial infection, normal aging changes, or vascular problems are not the targets of an endocrine workup in this context.

5. What is a hordeolum (stye)?

A. A chronic dry eye condition.

B. A allergic reaction of the eyelid.

C. A viral conjunctival infection.

D. An infection and inflammation of the eyelid, commonly referring to an infected eyelash follicle.

A stye is an acute infection and inflammation of the eyelid margin, most often arising from an infected eyelash follicle (or sometimes a meibomian gland). This causes a painful, red, localized lump at the lid edge, often with purulent discharge. It's typically treated with warm compresses and eyelid hygiene, and may require topical antibiotics for external styes or antibiotics if the infection involves deeper glands. It's distinct from chronic dry eye (a tear-film issue, not an infection), an allergic eyelid reaction (usually itchy with swelling but no purulent lump), and a viral conjunctival infection (affects the conjunctiva with eye discharge, not a tender eyelid nodule).

6. What is the management step for retinal detachment?

A. Wait and watch

B. Emergency ophthalmology referral

C. Home rest and analgesia

D. Oral antibiotics only

Retinal detachment is a sight-threatening emergency that can progress rapidly, potentially causing permanent vision loss if not treated promptly. The best management step is an emergency referral to ophthalmology for urgent evaluation and possible surgical reattachment. Depending on the detachment's type and extent, treatment may involve laser therapy or cryopexy for small or peripheral detachments, with definitive repair typically by scleral buckle or pars plana vitrectomy. Waiting, resting at home, or using analgesics or antibiotics does not address the underlying detachments and allows progression, so they are not appropriate.

SAMPLE

7. What are the symptoms of iritis?

A. Absent vision and eye movement.

B. Redness, pain, photophobia, blurred vision.

C. Diffuse redness with discharge.

D. Severe itching without pain.

Iritis is inflammation of the iris within the anterior uveal tract. The hallmark symptoms include eye pain, sensitivity to light (photophobia), redness, and blurred vision. Pain from the inflamed iris is often aching and worsens with light exposure; photophobia occurs because the iris reacts to bright light. Redness is typically a ciliary flush—redness around the cornea—rather than a diffuse conjunctival injection. Blurred vision results from inflammatory cells and protein particles (flare) in the anterior chamber interfering with light passage. This combination—red eye with eye pain, photophobia, and decreased vision—fits iritis well. By contrast, absence of vision or eye movement would raise concern for deeper orbital or optic problems, diffuse discharge points toward conjunctivitis, and itching without pain is more typical of allergic conjunctivitis.

8. What is the typical examination finding for a chalazion?

A. Painless lump on eyelid that is fast-growing.

B. Nontender lump that is slow-growing.

C. Red, tender swelling with crusting.

D. It is typically associated with fever.

Chalazion is a chronic blockage of a meibomian gland that leads to a sterile granulomatous inflammation. On exam, the classic finding is a firm, localized eyelid lump that is painless (nontender) and grows slowly over weeks. It is not red or acutely inflamed, and fever is not typical. This slow, non-painful progression distinguishes it from a stye, which is painful and red, or from cellulitis, which often includes fever and more diffuse eyelid swelling. So a nontender lump on the eyelid that enlarges slowly best fits a chalazion.

9. What are the symptoms of De Quervain's Tenosynovitis?

A. Painful swelling on the dorsal wrist

B. Inflammation of tendons on the side of the wrist at the base of the thumb.

C. Numbness in the little finger

D. Joint locking of the elbow

De Quervain's tenosynovitis happens when the tendons that move the thumb—the abductor pollicis longus and extensor pollicis brevis—become irritated and swollen as they pass through the first dorsal wrist compartment. This results in pain and often swelling on the radial (thumb-side) aspect of the wrist right at the base of the thumb. The pain is usually triggered or worsened by thumb movement, gripping, pinching, or lifting, and tenderness is felt directly over that area. In practice, patients notice a painful, swollen spot at the base of the thumb and difficulty with tasks that involve thumb use. Numbness in the little finger would point toward a different nerve issue, and elbow locking or non-specific dorsal wrist swelling wouldn't match the tenderness and pain pattern at the base of the thumb seen with this condition.

SAMPLE

10. What are common risk factors for developing a chalazion?

A. Diabetes, hypertension, obesity.

B. Acne rosacea, chronic blepharitis, seborrhea, tuberculosis, viral infection.

C. Myopia and hyperopia.

D. Only aging-related changes.

Chalazion forms when the Meibomian glands in the eyelids become obstructed by inflammatory debris and lipid stasis, leading to a localized granulomatous reaction. Conditions that cause chronic eyelid margin inflammation and Meibomian gland dysfunction increase this risk. Acne rosacea and chronic blepharitis are classic drivers of lid-margin inflammation, and seborrhea reflects a chronic skin condition that often accompanies blepharitis and meibomian gland dysfunction. When the eyelids are chronically inflamed, gland openings clog more easily, lipids back up, and a chalazion can develop. Infections can contribute as well. Viral infections that inflame the eyelids or conjunctiva perpetuate lid-margin irritation and obstruction, and tuberculosis is a granulomatous process that can involve the eyelid and mimic or precipitate chalazion-like lesions, making it a recognized albeit less common associated factor in certain settings. These factors fit chalazion risk much more closely than systemic/metabolic conditions (like diabetes, hypertension, or obesity) or refractive errors, which do not directly cause the eyelid margin inflammation or Meibomian gland obstruction that underlie chalazion formation.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://diffdiagmgmtacuteeyemskconditions.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE