DR. HIGH YIELD SURGERY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://drhighyieldsurgery.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What is the suggested imaging for diagnosing posterior urethral valves in a neonate?**

   A. Ultrasound of the abdomen

   B. Voiding cystourethrogram

   C. MRI of the pelvis

   D. CT scan of the abdomen

2. **According to Light's criteria, what indicates a transudative pleural effusion?**

   A. Low lactate dehydrogenase

   B. High protein ratio

   C. Low pleural glucose

   D. Presence of lymphocytes

3. **What is the first-line treatment for pseudomembranous colitis?**

   A. Oral metronidazole

   B. Oral vancomycin

   C. Intravenous vancomycin

   D. Probiotics

4. **What is a potential risk of using barium for a Gastrografin swallow in a case of Boerhaave syndrome?**

   A. It can worsen the symptoms

   B. It is less informative than Gastrografin

   C. It may cause allergic reactions

   D. It can obstruct the esophagus

5. **What is the most common cause of Cauda Equina syndrome?**

   A. Trauma

   B. Disk herniation

   C. Spinal stenosis

   D. Neoplasm

SAMPLE

6. In trauma situations, presence of peritoneal signs commonly indicates what condition?

A. Herniation

B. Peritonitis

C. Pancreatitis

D. Pneumothorax

7. What condition can mimic hypercalcemia presentations?

A. Diabetes mellitus

B. Hyperthyroidism

C. Acromegaly

D. Milk-alkali syndrome

8. The presence of flank pain and hypotension after a blood transfusion is indicative of which reaction?

A. Febrile non-hemolytic reaction

B. Acute hemolytic reaction

C. Hyperacute reaction

D. Delayed hemolytic reaction

9. What condition is suggested by a baby boy who cannot urinate on the first day of life?

A. Hypospadias

B. Posterior urethral valve obstruction

C. Ureteral atresia

D. Prenatal hydronephrosis

10. Which of the following is a diagnostic test for acute mesenteric ischemia?

A. Chemistry panel

B. CT angiography of the abdomen

C. Upper endoscopy

D. Stool test

SAMPLE

Answers

SAMPLE

1. B
2. A
3. B
4. A
5. B
6. B
7. D
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is the suggested imaging for diagnosing posterior urethral valves in a neonate?

A. Ultrasound of the abdomen

B. Voiding cystourethrogram

C. MRI of the pelvis

D. CT scan of the abdomen

Voiding cystourethrogram (VCUG) is the preferred imaging modality for diagnosing posterior urethral valves in neonates because it provides critical information about the anatomy and function of the urinary tract, particularly the urethra and bladder. This specific imaging technique allows for the visualization of the urethra during the filling and voiding phases, which is essential for identifying any obstruction caused by posterior urethral valves. During the procedure, contrast material is introduced into the bladder, enabling the clinician to observe how the urine flows through the urethra. If posterior urethral valves are present, the VCUG may reveal signs such as a dilated bladder or reflux into the kidneys, providing evidence of obstruction. While an abdominal ultrasound can be useful for assessing hydronephrosis or other renal issues, it lacks the detail necessary to evaluate the urethra directly. Magnetic resonance imaging (MRI) and computed tomography (CT) scans provide significant structural information but are typically not the first-line imaging techniques for this specific condition due to factors such as cost, availability, and exposure to radiation (particularly in the case of CT scans). Therefore, VCUG stands out as the most effective and appropriate choice for diagnosing posterior urethral valves in neonates

2. According to Light's criteria, what indicates a transudative pleural effusion?

A. Low lactate dehydrogenase

B. High protein ratio

C. Low pleural glucose

D. Presence of lymphocytes

Light's criteria are used to distinguish between transudative and exudative pleural effusions based on the composition of pleural fluid. According to these criteria, a pleural effusion is considered transudative if it meets at least one of the following criteria: the pleural fluid protein to serum protein ratio is less than 0.5, the pleural fluid lactate dehydrogenase (LDH) to serum LDH ratio is less than 0.6, and the pleural fluid LDH is less than two-thirds the upper limit of normal for serum LDH. In this case, the indication of a transudative pleural effusion with low lactate dehydrogenase aligns with the second criterion in Light's criteria, suggesting that the effusion is likely caused by systemic factors such as heart failure, cirrhosis, or nephrotic syndrome rather than local inflammatory processes. The other choices present features more typical of exudative pleural effusions. A high protein ratio is indicative of an exudative process, as is a low pleural glucose level due to metabolic activity in malignancies or infections. The presence of lymphocytes can often be associated with tuberculosis or malignancy, which also

SAMPLE

3. What is the first-line treatment for pseudomembranous colitis?

A. Oral metronidazole

B. Oral vancomycin

C. Intravenous vancomycin

D. Probiotics

The first-line treatment for pseudomembranous colitis, particularly when caused by Clostridium difficile, is oral vancomycin. This antibiotic is specifically effective against C. difficile due to its ability to remain concentrated in the intestine, which is crucial for treating intestinal infections. Oral vancomycin is preferred over other treatments primarily because it has a high efficacy for this infection and a favorable safety profile. Oral metronidazole may be considered in certain cases, especially in mild infections, but current guidelines emphasize the use of oral vancomycin as the primary treatment, particularly for moderate to severe cases of pseudomembranous colitis. Intravenous vancomycin is not effective for treating C. difficile infections because it does not reach adequate concentrations in the gut when administered via this route. Probiotics may have a role in prevention or adjunct therapy, but they are not established as a primary treatment for the active infection.

4. What is a potential risk of using barium for a Gastrografin swallow in a case of Boerhaave syndrome?

A. It can worsen the symptoms

B. It is less informative than Gastrografin

C. It may cause allergic reactions

D. It can obstruct the esophagus

In the context of Boerhaave syndrome, which is a spontaneous rupture of the esophagus, using barium for a swallow study poses significant risks. When barium is introduced into the gastrointestinal tract in the presence of an esophageal tear, it can lead to exacerbation of the condition. Specifically, barium can escape into the mediastinum, resulting in serious complications such as mediastinitis or chemical pneumonitis, both of which can worsen the patient's symptoms and overall clinical status. This worsening effect is crucial when diagnosing and managing this life-threatening condition. While allergic reactions and obstruction are potential risks of various contrast materials, they are less relevant in this specific scenario compared to the severe repercussions that barium can cause in Boerhaave syndrome. Additionally, the concern about barium being less informative than Gastrografin does not pertain to the risks but rather speaks to differences in imaging quality or diagnostic utility, which would not impact the immediate risks associated with the condition.

5. What is the most common cause of Cauda Equina syndrome?

A. Trauma

B. Disk herniation

C. Spinal stenosis

D. Neoplasm

Cauda Equina syndrome is primarily caused by the compression of the cauda equina nerves, which is most commonly due to disk herniation. When a herniated intervertebral disk occurs, especially in the lumbar region, it can protrude and exert pressure on these nerves. This can lead to a variety of neurological symptoms such as low back pain, sciatica, saddle anesthesia, and motor weakness, among others. While trauma, spinal stenosis, and neoplasms can also cause cauda equina syndrome, herniated disks are considerably more frequent in clinical practice, making them the leading cause. The mechanical pressure that results from a herniated disk is a well-understood mechanism explaining the onset of symptoms associated with this syndrome. Understanding this relationship is crucial for timely diagnosis and intervention to prevent permanent neurological damage.

EXAMPLE

6. In trauma situations, presence of peritoneal signs commonly indicates what condition?

A. Herniation

B. Peritonitis

C. Pancreatitis

D. Pneumothorax

In trauma situations, the presence of peritoneal signs typically indicates peritonitis. Peritoneal signs, such as tenderness, rigidity, and rebound tenderness, suggest that the peritoneum has become inflamed. This inflammation can be caused by an irritant, such as blood or infectious material, entering the peritoneal cavity due to the trauma. In the context of trauma, peritonitis often arises from injuries to hollow or solid organs, leading to leakage of their contents into the peritoneal space. The clinical presentation of peritonitis can be acute and may require prompt surgical intervention to address the underlying cause, such as repairing an organ injury or managing an infection. While other conditions mentioned, such as herniation, pancreatitis, and pneumothorax, may present with their own unique symptoms, they do not typically cause the distinct peritoneal signs seen in peritonitis. Herniation usually involves issues like nerve compression or vascular impairment, pancreatitis primarily presents with abdominal pain without the classic peritoneal signs, and pneumothorax typically exhibits respiratory difficulties and chest pain rather than abdominal symptoms. Thus, peritoneal signs specifically point towards a diagnosis of peritonitis in trauma cases.

7. What condition can mimic hypercalcemia presentations?

A. Diabetes mellitus

B. Hyperthyroidism

C. Acromegaly

D. Milk-alkali syndrome

Milk-alkali syndrome is characterized by a triad of hypercalcemia, metabolic alkalosis, and renal insufficiency. This condition arises from excessive intake of calcium and absorbable alkali, often from the consumption of calcium-rich products combined with alkali substances like antacids. The elevation of calcium levels in the blood can manifest symptoms similar to those of hypercalcemia itself, such as nausea, vomiting, confusion, and fatigue. Understanding why milk-alkali syndrome closely mimics hypercalcemia is crucial. In this syndrome, the body's response to high calcium levels can lead to similar clinical presentations, making it difficult to distinguish between the two without further diagnostic evaluation. In contrast, the other conditions listed do not present with increased serum calcium as a primary feature. Diabetes mellitus primarily affects glucose metabolism and can lead to various complications, while hyperthyroidism is related to excessive thyroid hormone levels, influencing metabolism but not directly causing hypercalcemia. Acromegaly, a result of excess growth hormone, affects growth and development but does not cause elevated calcium levels in the same manner. Therefore, it is the unique mechanism of milk-alkali syndrome that allows it to mimic the presentations of hypercalcemia effectively.

SAMPLE

8. The presence of flank pain and hypotension after a blood transfusion is indicative of which reaction?

A. Febrile non-hemolytic reaction

B. Acute hemolytic reaction

C. Hyperacute reaction

D. Delayed hemolytic reaction

Flank pain and hypotension occurring after a blood transfusion are hallmark signs of an acute hemolytic reaction. This type of reaction often results from ABO blood group incompatibility, where the recipient's immune system responds to the transfused red blood cells as foreign. The rapid destruction of these cells leads to the release of hemoglobin and other intracellular components into the circulation. This can cause significant cardiovascular instability, resulting in hypotension. Flank pain often arises from kidney damage due to the hemolysis and the subsequent release of free hemoglobin, which can lead to renal tubular obstruction and acute kidney injury. In contrast, febrile non-hemolytic reactions are usually characterized by fever and chills without the severe systemic manifestations seen in acute hemolytic reactions. Hyperacute reactions, while also serious, are rare and typically occur almost immediately after transfusion due to pre-existing antibodies—usually in cases of organ transplantation rather than blood transfusion. Delayed hemolytic reactions can occur days to weeks after a transfusion and are less likely to present with acute flank pain and hypotension. Overall, the combination of flank pain and hypotension specifically points towards an acute hemolytic reaction as the cause, emphasizing the need for careful monitoring during and

9. What condition is suggested by a baby boy who cannot urinate on the first day of life?

A. Hypospadias

B. Posterior urethral valve obstruction

C. Ureteral atresia

D. Prenatal hydronephrosis

The situation described, where a baby boy cannot urinate within the first day of life, is highly suggestive of posterior urethral valve obstruction. This condition arises from the presence of abnormal flaps of tissue (valves) in the posterior urethra, leading to obstruction of urine flow. In male infants, this is the most common cause of urinary obstruction, and it can result in significant urinary retention. A key aspect of this diagnosis is the timing; inability to urinate shortly after birth indicates a considerable obstruction, as urine production begins shortly after delivery. If the obstruction is severe, it may result in a distended bladder and renal problems due to backpressure. Other conditions, such as hypospadias or ureteral atresia, while potentially causing urinary issues, would not typically present with an acute inability to urinate immediately after birth in the same way, placing posterior urethral valve obstruction at the forefront for consideration in this scenario.

SAMPLE

10. Which of the following is a diagnostic test for acute mesenteric ischemia?

A. Chemistry panel

B. CT angiography of the abdomen

C. Upper endoscopy

D. Stool test

The diagnostic test for acute mesenteric ischemia is CT angiography of the abdomen. This imaging modality is particularly effective in evaluating blood vessels and can reveal arterial occlusions or significant stenosis that may lead to mesenteric ischemia. CT angiography provides a rapid and non-invasive way to visualize the mesenteric circulation, showing not only the presence of emboli or thrombi in the mesenteric arteries but also assessing the viability of the bowel. Early diagnosis is crucial in acute mesenteric ischemia, as timely intervention can significantly affect patient outcomes. Other options are not specific for diagnosing acute mesenteric ischemia; for instance, a chemistry panel may provide metabolic information but lacks the specificity needed for vascular or perfusion issues. Upper endoscopy focuses on the upper gastrointestinal tract and does not visualize mesenteric vessels, hence it's not useful for diagnosing ischemia in the mesentery. A stool test may help identify gastrointestinal bleeding or infectious processes, but again, it does not directly assess vascular supply to the bowel. Thus, CT angiography is the most appropriate and specific diagnostic tool for this condition.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://drhighyieldsurgery.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE