ECCO CARING FOR PATIENTS WITH CARDIOVASCULAR DISORDERS PART 1 PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://eccocaringforcardiovasculardisorders1.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which complication can occur from myocardial infarction involving the papillary muscles?

A. Acute mitral regurgitation leading to pulmonary edema and shock

B. Atrial fibrillation

C. Ventricular septal defect

D. Pericarditis

2. Which combination of signs is associated with pericardial tamponade?

A. Hypertension, muffled heart sounds, elevated JVP; pulsus paradoxus.

B. Hypotension, muffled heart sounds, elevated JVP; not including pulsus paradoxus.

C. Chest pain with radiation to the left arm.

D. Hypotension, muffled heart sounds, elevated JVP; pulsus paradoxus.

3. How does anticoagulation differ between nonvalvular atrial fibrillation and valvular atrial fibrillation?

A. Nonvalvular AF is treated with DOACs or warfarin; valvular AF (with mechanical valves or rheumatic MR/AS) requires warfarin

B. Nonvalvular AF requires warfarin only; valvular AF can use DOACs

C. Nonvalvular AF uses aspirin; valvular AF uses warfarin only

D. Both can be managed with DOACs

4. What position is typically recommended to optimize ventilation in a patient with cardiogenic shock?

A. Trendelenburg

B. Semi-Fowler position

C. Prone

D. Supine flat

5. What are the main reasons nitroglycerin is given to a patient with ACS?

A. To increase afterload to improve perfusion

B. To reduce preload and dilate coronary arteries

C. To decrease heart rate significantly

D. To treat infection

SAMPLE

6. Which diagnostic test is primarily used to assess left ventricular ejection fraction in heart failure?

A. Echocardiography

B. Chest radiography

C. Electrocardiography

D. Cardiac MRI

7. Which Doppler measurements are used to assess diastolic function?

A. Systolic ejection fraction.

B. QRS duration.

C. E/A and E/e' ratios.

D. Pulmonary capillary wedge pressure.

8. What does a troponin elevation indicate in acute coronary syndrome?

A. Liver injury

B. Kidney injury

C. Pulmonary embolism

D. Myocardial injury

9. Which should the nurse include when educating a patient on reducing the risk of acute coronary syndrome?

A. Stop smoking

B. Ignore symptoms

C. Eat a diet high in saturated fat

D. Avoid physical activity

10. Troponin elevations in myocardial ischemia typically rise within hours after injury and persist for days. Which choice reflects this timing?

A. Rise within seconds

B. Rise within hours and persist for days

C. Do not rise in ACS

D. Rise within hours and persist for days

SAMPLE

Answers

SAMPLE

1. A
2. D
3. A
4. B
5. B
6. A
7. C
8. D
9. A
10. D

SAMPLE

Explanations

SAMPLE

1. Which complication can occur from myocardial infarction involving the papillary muscles?

A. Acute mitral regurgitation leading to pulmonary edema and shock

B. Atrial fibrillation

C. Ventricular septal defect

D. Pericarditis

Damage to the papillary muscles during a myocardial infarction disrupts the mitral valve apparatus. The papillary muscles anchor the chordae tendineae to the mitral leaflets, helping them close properly in systole. If these muscles are infarcted or rupture, the leaflets can't coapt, and mitral regurgitation occurs suddenly. The backflow into the left atrium raises left atrial and pulmonary venous pressures, leading to acute pulmonary edema and a drop in forward cardiac output, which can cause shock. Clinically, this presents as a new or rapidly worsening murmur with signs of heart failure after an MI. Other post-MI complications like atrial fibrillation, ventricular septal defect, or pericarditis can occur, but they are not specifically caused by papillary muscle injury. Atrial fibrillation is an arrhythmia, ventricular septal defect results from interventricular septum rupture, and pericarditis is inflammation around the heart; none arise directly from papillary muscle disruption in the way acute mitral regurgitation does.

2. Which combination of signs is associated with pericardial tamponade?

A. Hypertension, muffled heart sounds, elevated JVP; pulsus paradoxus.

B. Hypotension, muffled heart sounds, elevated JVP; not including pulsus paradoxus.

C. Chest pain with radiation to the left arm.

D. Hypotension, muffled heart sounds, elevated JVP; pulsus paradoxus.

Pericardial tamponade happens when fluid in the pericardial space raises pressure around the heart, making it hard for the heart to fill properly during diastole. That reduced filling lowers the stroke volume and cardiac output, so blood pressure tends to fall (hypotension). At the same time, blood backs up into the superior vena cava, elevating the jugular venous pressure. The heart is also cushioned by fluid, which dampens the sounds heard with a stethoscope (muffled heart sounds). A classic extra sign is pulsus paradoxus, where the systolic blood pressure drops more than about 10 mm Hg during inspiration because the heart's filling is severely limited by the surrounding fluid. So the best combination is hypotension with muffled heart sounds, elevated JVP, and pulsus paradoxus, all aligning with tamponade physiology. Chest pain radiating to the left arm points more toward myocardial ischemia, and hypertension contradicts the expected low filling pressures seen in tamponade. Not all patients show pulsus paradoxus, but its presence strongly supports tamponade when other signs are present.

SAMPLE

3. How does anticoagulation differ between nonvalvular atrial fibrillation and valvular atrial fibrillation?

A. Nonvalvular AF is treated with DOACs or warfarin; valvular AF (with mechanical valves or rheumatic MR/AS) requires warfarin

B. Nonvalvular AF requires warfarin only; valvular AF can use DOACs

C. Nonvalvular AF uses aspirin; valvular AF uses warfarin only

D. Both can be managed with DOACs

Anticoagulation choices hinge on whether atrial fibrillation is associated with a mechanical valve or rheumatic valve disease. In nonvalvular AF, DOACs are approved and commonly preferred because they provide similar protection against stroke as warfarin but with fixed dosing and less monitoring. In contrast, valvular AF that involves a mechanical heart valve or rheumatic mitral stenosis is not treated with DOACs; evidence shows warfarin remains necessary to prevent thromboembolism in these situations. Trials and guidelines have shown increased risk with DOACs in mechanical valves (and DOACs have not proven safe/effective in rheumatic mitral stenosis), so warfarin is the recommended anticoagulant for those patients. Thus, the statement that nonvalvular AF can be treated with either a DOAC or warfarin, while valvular AF with mechanical valves or rheumatic MR/AS requires warfarin, best captures the distinction.

4. What position is typically recommended to optimize ventilation in a patient with cardiogenic shock?

A. Trendelenburg

B. Semi-Fowler position

C. Prone

D. Supine flat

Elevating the head of the bed to a semi-Fowler position (about 30–45 degrees) helps ventilation in cardiogenic shock by improving lung mechanics and reducing pulmonary congestion. This position decreases venous return just enough to lessen left sided filling pressures and edema around the lungs, making it easier for the lungs to expand and for oxygen to transfer into the blood. It also allows the diaphragm to descend more effectively, improving airflow. Lying flat increases preload and can worsen dyspnea in pulmonary edema, while Trendelenburg raises venous return further and can worsen edema. Prone positioning is not routinely used for typical cardiogenic shock. So, semi-Fowler is the most beneficial choice for optimizing ventilation in this scenario.

5. What are the main reasons nitroglycerin is given to a patient with ACS?

A. To increase afterload to improve perfusion

B. To reduce preload and dilate coronary arteries

C. To decrease heart rate significantly

D. To treat infection

In ACS, the main goal of nitroglycerin is to lower the heart's oxygen demand and improve blood flow to the ischemic tissue. The primary way this happens is venodilation, which reduces preload (the volume entering the heart). With less volume, the ventricle works less hard, wall stress drops, and myocardial oxygen consumption decreases. Nitroglycerin can also dilate coronary arteries, helping improve perfusion to areas of ischemia. This combination relieves chest pain and helps balance the oxygen supply and demand in the heart. It's not used to increase afterload or to markedly slow the heart rate, and it's not a treatment for infection.

SAMPLE

6. Which diagnostic test is primarily used to assess left ventricular ejection fraction in heart failure?

A. Echocardiography

B. Chest radiography

C. Electrocardiography

D. Cardiac MRI

Assessing left ventricular ejection fraction in heart failure is best done with echocardiography because ultrasound imaging provides real-time visualization of the heart's chambers and contraction, allowing calculation of LV volumes and the fraction of blood ejected with each beat. By using methods like the biplane Simpson's rule, echo yields a numerical LVEF, which directly reflects systolic function and helps classify heart failure, guide treatment, and track changes over time. In addition to LVEF, echocardiography offers information about chamber size, wall motion, and valve function, all without radiation and with wide availability, often at the bedside. Chest X-ray can hint at congestive changes but does not quantify LVEF. ECG shows rhythm and conduction or ischemia but not the amount of blood the ventricle pumps. Cardiac MRI can measure LVEF with high accuracy and may be used when precision is needed or echo windows are poor, but it is less accessible, more costly, and less practical in many clinical settings, making echocardiography the primary test for this purpose.

7. Which Doppler measurements are used to assess diastolic function?

A. Systolic ejection fraction.

B. QRS duration.

C. E/A and E/e' ratios.

D. Pulmonary capillary wedge pressure.

Doppler assessment of diastolic function looks at how blood fills the left ventricle during diastole and how the myocardium relaxes. The mitral inflow pattern provides the E wave, which is the rapid early filling velocity, and the A wave, which comes from atrial contraction. The E/A ratio compares these two phases and reveals whether early filling is reduced or whether filling relies more on atrial contraction—a sign of different stages of diastolic dysfunction. To refine this, tissue Doppler measures e', the myocardial relaxation velocity at the mitral annulus. The E/e' ratio combines these signals to estimate left ventricular filling pressures noninvasively: a higher ratio suggests higher filling pressures and worse diastolic function. Other options aren't Doppler-based assessments of diastolic function. Systolic ejection fraction assesses systolic performance, not diastolic filling. QRS duration is an ECG interval reflecting ventricular depolarization, not diastolic filling. Pulmonary capillary wedge pressure is an invasive hemodynamic pressure measurement, not a Doppler-derived diastolic index.

8. What does a troponin elevation indicate in acute coronary syndrome?

A. Liver injury

B. Kidney injury

C. Pulmonary embolism

D. Myocardial injury

Troponin elevation signals myocardial injury from acute coronary syndrome. Troponin I and troponin T are cardiac-specific proteins that are released when heart muscle cells are damaged by ischemia from a ruptured plaque and coronary occlusion. Because troponin comes from heart muscle, its presence in the blood is a direct indicator of myocardial injury, and the pattern over time (rise and/or fall) helps confirm an ACS event and guides urgent treatment. While troponin can be elevated in other situations (rarely due to non-heart conditions), in the ACS setting it specifically points to damage of the heart muscle rather than liver or kidney injury or pulmonary embolism.

SAMPLE

9. Which should the nurse include when educating a patient on reducing the risk of acute coronary syndrome?

A. Stop smoking

B. Ignore symptoms

C. Eat a diet high in saturated fat

D. Avoid physical activity

Quitting smoking is a major, immediate step to lower the risk of acute coronary syndrome. Smoking causes or worsens several factors that drive heart events: it promotes atherosclerosis, raises heart rate and blood pressure, reduces oxygen delivery to heart tissue, and makes the blood more prone to clotting. When a person stops smoking, these harmful effects begin to reverse—heart rate and blood pressure improve, carbon monoxide levels fall, and endothelial function improves. The risk of a coronary event drops quickly after quitting and continues to decline over time, eventually approaching that of someone who has never smoked. So, in patient education, emphasize stopping smoking as the primary modifiable protective measure. Support the patient with practical quitting strategies, such as setting a quit date, using appropriate cessation aids, seeking behavioral counseling, avoiding smoking triggers, staying physically active, and managing weight.

10. Troponin elevations in myocardial ischemia typically rise within hours after injury and persist for days. Which choice reflects this timing?

A. Rise within seconds

B. Rise within hours and persist for days

C. Do not rise in ACS

D. Rise within hours and persist for days

Troponin is a protein released into the blood when heart muscle cells are damaged. In myocardial ischemia or infarction, it does not appear instantly; it typically becomes detectable after a few hours (roughly 3–4 hours) and then can rise to a peak over the next day. Because troponin clearance from the blood is slow, these levels stay elevated for days—often about a week or longer, depending on the assay and the patient. So the statement that troponin rises within hours after injury and remains elevated for days best captures its timing in ACS. Notable contrast: troponin is not typically detectable within seconds, and it does rise in ACS, so the other ideas don't fit the known pattern.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://eccocaringforcardiovasculardisorders1.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE