ECHOCARDIOGRAPHY TECHNIQUES III EXAM 1 PRACTICE (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://echocardiographytech3exam1.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What are the steps to calculate MR EROA using PISA?

A. Find the aliasing velocity; measure radius R at aliasing; Flow = 2πR^2 × aliasing velocity; EROA = Flow / Vmax MR; MR volume = EROA × VTI MR

B. Use Vmax MR alone to compute EROA

C. Use stroke volume difference between left heart and right heart

D. Use LVOT diameter alone

2. Which statement best describes the vena contracta?

A. Upstream of the narrowed orifice

B. The widest part of the jet

C. The narrowest part of a jet downstream from a narrowed orifice

D. At the region of jet impingement on the wall

3. What is the motion pattern of the anterior tricuspid leaflet in Ebstein's anomaly?

A. whip-like motion

B. normal motion

C. restricted motion

D. paradoxical motion

4. A 90-degree Doppler angle yields zero Doppler shift. Which option lists 90 degrees as the angle?

A. 0 degrees

B. 45 degrees

C. 180 degrees

D. 90 degrees

5. What is the primary clinical triad associated with severe aortic stenosis?

A. Chest pain only

B. Angina pectoris, syncope, and CHF

C. Shortness of breath exclusively

D. Hypertension and edema

SAMPLE

6. RAP estimation by echo often uses the collapsibility of which structure?

A. Aorta

B. Superior vena cava

C. Inferior vena cava

D. Pulmonary artery

7. Explain the significance of an aortic regurgitation jet timing in diastole.

A. All of the above

B. AR jets occur during diastole

C. Holodiastolic flow reversal indicates severity

D. Jet timing helps differentiate acute from chronic AR

8. What is the role of subcostal views in adult echocardiography?

A. All of the above

B. Provide alternative windows

C. Useful when parasternal windows are poor

D. Helpful for evaluating the atrial septum and certain congenital or diaphragmatic conditions

9. In calcific aortic stenosis with MAC, what should the sonographer assess and what condition often results in LVH?

A. LV wall thickness and systolic function

B. LV end-diastolic volume

C. Right atrial size

D. Pulmonary regurgitation velocity

10. Which scenario biases the PISA method when quantifying MR?

A. Eccentric jets, irregular orifices, or complex anatomy

B. Central jets with a symmetric orifice

C. When MR is trivial

D. When the echocardiographic window is perfect

SAMPLE

Answers

SAMPLE

1. A
2. C
3. A
4. D
5. B
6. C
7. A
8. A
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. What are the steps to calculate MR EROA using PISA?

A. Find the aliasing velocity; measure radius R at aliasing; Flow = 2πR^2 × aliasing velocity; EROA = Flow / Vmax MR; MR volume = EROA × VTI MR

B. Use Vmax MR alone to compute EROA

C. Use stroke volume difference between left heart and right heart

D. Use LVOT diameter alone

The main idea is to use the PISA method to turn the regurgitant flow into an effective orifice area. Start by setting the aliasing velocity with color Doppler and measure the radius of the hemispherical flow convergence at that aliasing velocity. The regurgitant flow rate is Flow = 2πR^2 × V_a. Then combine that flow with the peak MR velocity from spectral Doppler to get the effective regurgitant orifice area: EROA = Flow / V_max MR. To obtain the regurgitant volume, multiply EROA by the MR velocity-time integral (VTI MR). This sequence ties the hemispherical flow concept to an area and then to volume. The other options don't fit: V_max MR alone can't determine EROA without flow; using stroke-volume differences or LVOT diameter alone yields different measurements and isn't the PISA-derived EROA approach.

2. Which statement best describes the vena contracta?

A. Upstream of the narrowed orifice

B. The widest part of the jet

C. The narrowest part of a jet downstream from a narrowed orifice

D. At the region of jet impingement on the wall

The vena contracta is the narrowest portion of a jet that forms as fluid must pass through a restricted opening. It occurs immediately downstream of the narrowed orifice, where the flow accelerates and the jet cross-section becomes smallest. This region has the highest velocity and the smallest cross-sectional area before the jet later expands as it moves away from the orifice. That's why the best description is the narrowest part of a jet downstream from a narrowed orifice. Upstream would be before the restriction, the widest part is after the jet has expanded, and the jet impingement on the wall is a downstream effect that occurs after the vena contracta.

3. What is the motion pattern of the anterior tricuspid leaflet in Ebstein's anomaly?

A. whip-like motion

B. normal motion

C. restricted motion

D. paradoxical motion

In Ebstein's anomaly the septal and posterior leaflets are displaced toward the apex, creating an atrialized right ventricle and leaving the anterior leaflet as the dominant, highly mobile component. Its attachment and tethering produce a rapid, pendulous excursion of the anterior leaflet during systole, which appears as a whip-like motion on imaging. This motion is distinctive and contrasts with normal leaflet motion; it is not typically restricted or paradoxical in this condition.

SAMPLE

4. A 90-degree Doppler angle yields zero Doppler shift. Which option lists 90 degrees as the angle?

A. 0 degrees

B. 45 degrees

C. 180 degrees

D. 90 degrees

Doppler shift depends on the angle between the ultrasound beam and the direction of blood flow, through the cosine of that angle. When the beam is at 90 degrees to the flow, the component of velocity along the beam is zero, so the frequency shift vanishes. In other words, cos 90° equals zero, making the Doppler shift zero regardless of how fast the blood is moving. That's why this option, which lists 90 degrees, is the correct one. Remember, at 0° the shift is maximal in the positive direction, and at 180° it's maximal in the negative direction, which is why aligning the beam as parallel as possible to flow is important for Doppler measurements.

5. What is the primary clinical triad associated with severe aortic stenosis?

A. Chest pain only

B. Angina pectoris, syncope, and CHF

C. Shortness of breath exclusively

D. Hypertension and edema

Severe aortic stenosis creates a fixed obstruction to left ventricular outflow, so the heart cannot increase cardiac output with activity. This leads to a characteristic triad of symptoms: angina from increased LV wall stress and higher myocardial oxygen demand with relatively reduced coronary perfusion; syncope from transient cerebral hypoperfusion during exertion when cardiac output cannot rise; and congestive heart failure symptoms from chronic pressure overload causing diastolic dysfunction and pulmonary congestion. This combination captures the classic presentation, whereas chest pain alone, shortness of breath alone, or unrelated signs like hypertension don't fully describe the typical causing factors and sequence seen with severe AS.

6. RAP estimation by echo often uses the collapsibility of which structure?

A. Aorta

B. Superior vena cava

C. Inferior vena cava

D. Pulmonary artery

RAP estimation by echo relies on how much the inferior vena cava changes size with respiration. The IVC drains directly into the right atrium, so its caliber tracks the pressure in the right atrium. When the right atrial pressure is low, the IVC is smaller and collapses readily with inspiration. When the right atrial pressure is high, the IVC becomes dilated and shows little inspiratory collapse. By measuring the IVC diameter near its entry to the RA and assessing the percent collapse during respiration, clinicians infer the likely right atrial pressure. For example, a small IVC with substantial collapse tends to correspond to normal or low RAP, whereas a large IVC with minimal collapse suggests elevated RAP. This is why the inferior vena cava, rather than the aorta, superior vena cava, or pulmonary artery, is used for this estimation. Keep in mind conditions like obesity, lung disease, or mechanical ventilation can affect the measurement, so interpretation should consider the overall clinical context.

SAMPLE

7. Explain the significance of an aortic regurgitation jet timing in diastole.

A. All of the above

B. AR jets occur during diastole

C. Holodiastolic flow reversal indicates severity

D. Jet timing helps differentiate acute from chronic AR

The timing of the aortic regurgitation jet in diastole reflects how much and how long blood is flowing backward from the aorta into the LV when the aortic pressure exceeds LV diastolic pressure. Because regurgitation happens during diastole, you'll see the AR jet on Doppler as a diastolic flow from the aorta into the LV. If the regurgitant volume is large, the jet can persist throughout the entire diastolic period (holodiastolic), which is a sign of severe AR. Holodiastolic flow reversal in the descending aorta is another important clue: it indicates that enough regurgitant volume is returning during diastole to reverse flow in the systemic arterial tree, reinforcing the assessment of severity. The jet timing also helps distinguish acute from chronic AR because the hemodynamic and structural adaptations differ between them; chronic AR often shows more pronounced diastolic run-off and arterial changes over time, whereas acute AR presents with a rapid, abrupt diastolic jet and different flow patterns due to the lack of LV remodeling. Putting these together, the diastolic jet timing informs presence, severity, and chronicity of AR.

8. What is the role of subcostal views in adult echocardiography?

A. All of the above

B. Provide alternative windows

C. Useful when parasternal windows are poor

D. Helpful for evaluating the atrial septum and certain congenital or diaphragmatic conditions

Subcostal views provide a different acoustic window that complements the standard parasternal and apical views. This approach is especially valuable when the usual windows are limited by body habitus, lung disease, or postsurgical changes, because it often yields clearer visualization of the right heart and interatrial septum from below the sternum. From the subcostal position, the interatrial septum is seen well, making it a reliable window for evaluating atrial septal defects or other septal abnormalities in adults. It also supports assessment of congenitally related or diaphragmatic conditions and can be modified into a transhepatic view to visualize right heart structures and the IVC when chest windows are poor or when diaphragmatic motion or abdominal anatomy provides a better acoustic path. For these reasons, subcostal views serve as alternative windows, are particularly useful when parasternal windows are poor, and aid in evaluating the atrial septum and certain congenital or diaphragmatic conditions. All of these roles are encompassed by using the subcostal approach.

SAMPLE

9. In calcific aortic stenosis with MAC, what should the sonographer assess and what condition often results in LVH?

A. LV wall thickness and systolic function

B. LV end-diastolic volume

C. Right atrial size

D. Pulmonary regurgitation velocity

The left ventricle adapts to the increased afterload from calcific aortic stenosis (often with MAC) by developing concentric hypertrophy. The key thing a sonographer should assess is the left ventricular wall thickness and its systolic function. Thickened walls (increased LV wall thickness) reflect hypertrophy, and evaluating systolic function shows whether the chamber can still contract effectively, which is typically preserved early on. This chronic pressure overload is what commonly leads to LV hypertrophy. Other options don't directly reflect this adaptation: LV end-diastolic volume is more about volume status and eccentric remodeling, the right atrial size pertains to right-sided pressures, and pulmonary regurgitation velocity relates to pulmonary circulation rather than the LV's response to aortic valve disease.

10. Which scenario biases the PISA method when quantifying MR?

A. Eccentric jets, irregular orifices, or complex anatomy

B. Central jets with a symmetric orifice

C. When MR is trivial

D. When the echocardiographic window is perfect

PISA quantification of MR depends on a hemispherical flow convergence region and a circular regurgitant orifice that produces a single, central jet. When the jet is eccentric, the orifice is irregular, or the anatomy is complex, these assumptions break down, so the measured radius at the aliasing velocity does not accurately reflect the true flow. This distorts the calculated flow rate and thus biases the estimated effective regurgitant orifice area and regurgitant volume. If the jet is central with a symmetrical orifice, the hemispheric flow model holds better and the PISA calculation remains reliable. When MR is trivial, the signal is small and measurements become more susceptible to noise, but the geometry isn't inherently biased. A perfect echocardiographic window by itself doesn't introduce bias; the issue is the jet shape and orifice geometry that violate PISA assumptions.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://echocardiographytech3exam1.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE