EMERGENCY ENDOTRACHEAL INTUBATION (EEI) AIRWAY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://eeiairway.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What are the two main laryngoscope blade types, and how do they differ?

A. Macintosh (curved blade) and Miller (straight blade); different shapes to optimize glottic visualization.

B. Miller (curved blade) and Macintosh (straight blade); different shapes to optimize glottic visualization.

C. Both blades are curved; the difference is handle length only.

D. Laryngoscope blades differ only in material, not shape.

2. Alveolar ventilation is defined as which of the following?

A. The total air entering the lungs per minute

B. The volume of air that reaches the alveoli; determined by subtracting dead space air from tidal volume

C. The maximum volume of air exhaled in one breath

D. The oxygen content in the alveoli

3. Which monitoring tools are essential for EEI?

A. Temperature and urine output only

B. Capnography, pulse oximetry, ECG, blood pressure, and airway device status checks

C. Only capnography

D. Only pulse oximetry and heart rate

4. In addition to capnography, which imaging modality can verify endotracheal tube position when available?

A. Chest radiography can verify endotracheal tube position when available.

B. MRI is the standard method for initial confirmation.

C. Ultrasound is the first-line confirmatory test.

D. CT scan provides immediate tube position.

5. Which condition is characterized by wheezing on both inspiration and expiration and bronchospasm?

A. Asthma.

B. Anaphylaxis.

C. Bronchiolitis.

D. Congestive Heart Failure.

SAMPLE

6. What does a cuff leak around the endotracheal tube indicate?

A. A leak around the deflated cuff indicates lower airway edema and reduced risk of post-extubation stridor.

B. A leak indicates severe edema.

C. No leak means the airway is always ready.

D. A leak confirms tube placement.

7. Which condition is described as destruction of airways distal to the bronchioles, with involvement of the pulmonary capillary bed?

A. Emphysema

B. CHF

C. RSV

D. TB

8. Which condition is associated with shortness of breath that is mostly on exertion and progressively worsens?

A. COPD

B. CHF

C. Emphysema

D. Chronic Bronchitis

9. Ataxic respirations are defined as which pattern?

A. Regular and predictable breathing

B. Pattern of irregular and unpredictable breathing

C. Rapid and shallow breathing

D. No breathing

10. Which condition commonly presents with abdominal distension, dependent edema, and increased breathing rate?

A. CHF

B. COPD

C. Emphysema

D. Chronic Bronchitis

SAMPLE

Answers

SAMPLE

1. A
2. B
3. B
4. A
5. A
6. A
7. A
8. A
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What are the two main laryngoscope blade types, and how do they differ?

A. Macintosh (curved blade) and Miller (straight blade); different shapes to optimize glottic visualization.

B. Miller (curved blade) and Macintosh (straight blade); different shapes to optimize glottic visualization.

C. Both blades are curved; the difference is handle length only.

D. Laryngoscope blades differ only in material, not shape.

The main idea here is how blade shape changes the way you lift tissues to visualize the glottis. The curved blade, the Macintosh, fits along the tongue's curve. You insert the tip into the vallecula (the space between the base of the tongue and the epiglottis) and lift gently. That lifting acts indirectly on the epiglottis and opens the view of the vocal cords with typically less force and less tissue disruption. The straight blade, the Miller, has a straight edge that is aimed directly under the epiglottis to lift it up. This direct lift can provide a good view in certain anatomies or when you need more leverage, such as in some pediatric airways or when the epiglottis sits in a position that's easier to lift directly. So, the two main blade designs differ primarily in their shape and the way they achieve glottic visualization: a curved blade that lifts via the vallecula, and a straight blade that lifts the epiglottis directly. This is why the option describing a curved Macintosh and a straight Miller best captures the difference.

2. Alveolar ventilation is defined as which of the following?

A. The total air entering the lungs per minute

B. The volume of air that reaches the alveoli; determined by subtracting dead space air from tidal volume

C. The maximum volume of air exhaled in one breath

D. The oxygen content in the alveoli

Alveolar ventilation is the amount of air per unit time that actually reaches the gas-exchanging areas of the lungs. Per breath, this is the portion of tidal volume that makes it to the alveoli, which means subtracting the air that fills the conducting airways (dead space) from the total tidal volume. So the air reaching the alveoli per breath is tidal volume minus dead space, and to get the per-minute value you multiply by the breathing rate. This is why the best description is the volume of air that reaches the alveoli, determined by subtracting dead space air from tidal volume. If you know tidal volume and dead space, you can estimate alveolar ventilation, and you'd multiply by the respiratory rate to get per-minute alveolar ventilation (VT − VD) × f. Understanding helps you see why the other ideas aren't alveolar ventilation: total air entering the lungs per minute is minute ventilation, not all of which reaches the alveoli; the maximum volume exhaled in one breath describes a different lung volume measure; and oxygen content in the alveoli describes gas composition, not the amount of air reaching the alveoli for gas exchange.

SAMPLE

3. Which monitoring tools are essential for EEI?

A. Temperature and urine output only

B. Capnography, pulse oximetry, ECG, blood pressure, and airway device status checks

C. Only capnography

D. Only pulse oximetry and heart rate

During emergency endotracheal intubation, real-time monitoring across several systems is essential to detect problems early and guide rapid actions. The best monitoring set includes capnography to confirm tube placement and ventilation by showing a continuous end-tidal CO2 waveform, pulse oximetry to track oxygenation, ECG to observe heart rhythm and detect hypoxia-related changes or ischemia, noninvasive blood pressure to assess the patient's hemodynamic response and guide treatment, and ongoing airway device status checks to verify the tube remains in the right position, is patent, and is delivering ventilation effectively (watching for breath sounds, chest rise, and equal ventilation). This combination provides immediate feedback on ventilation, oxygen delivery, cardiovascular status, and device integrity, which are all critical during a high-stakes procedure. Relying on only temperature and urine output, or on a single monitoring modality, misses key aspects of safety during EEI. Capnography alone won't reveal oxygen saturation or blood pressure changes, and airway placement isn't confirmed without device checks and clinical ventilation evidence. Conversely, using only pulse oximetry or only one cardiovascular or respiratory tool fails to provide a complete, rapid picture of the patient's status during intubation.

4. In addition to capnography, which imaging modality can verify endotracheal tube position when available?

A. Chest radiography can verify endotracheal tube position when available.

B. MRI is the standard method for initial confirmation.

C. Ultrasound is the first-line confirmatory test.

D. CT scan provides immediate tube position.

Beyond capnography, chest radiography is used to verify endotracheal tube position. It provides a clear image showing the tube tip relative to the carina, confirming the tube is in the trachea and at a safe depth. In adults, aim for the tip about 2–3 cm above the carina (adjust for pediatric size). Portable chest X-ray is quick, widely available, and can also assess other thoracic lines and overall lung status. Other imaging options aren't as practical for immediate confirmation: MRI isn't fast enough or commonly used for this purpose, CT isn't immediate and isn't feasible in unstable patients, and ultrasound, while helpful in some settings, isn't the standard confirmatory test.

SAMPLE

5. Which condition is characterized by wheezing on both inspiration and expiration and bronchospasm?

A. Asthma.

B. Anaphylaxis.

C. Bronchiolitis.

D. Congestive Heart Failure.

Bronchospasm with wheezing in the setting of an obstructive, reactive airway disease is the hallmark of asthma. In asthma, the airway smooth muscle constricts, and there's mucosal swelling and increased mucus, narrowing the airways. This causes turbulent airflow and audible wheezing. While wheezing is most often heard during expiration because that phase is when air flow is most restricted, extensive or severe bronchospasm can produce wheeze during both inspiration and expiration. That combination—wheezing occurring with both phases of breathing due to reversible bronchoconstriction—fits asthma well, since the underlying problem is bronchospasm that responds to bronchodilators. Anaphylaxis can involve bronchospasm, but it's a systemic reaction with other signs like hypotension, rash, or airway edema. Bronchiolitis usually affects infants and presents with viral symptoms and diffuse wheeze/crackles rather than prominent, reversible bronchospasm. Congestive heart failure features fluid overload and crackles from edema, not bronchospastic wheezing.

6. What does a cuff leak around the endotracheal tube indicate?

A. A leak around the deflated cuff indicates lower airway edema and reduced risk of post-extubation stridor.

B. A leak indicates severe edema.

C. No leak means the airway is always ready.

D. A leak confirms tube placement.

A cuff leak around the endotracheal tube after deflating the cuff reflects that air can pass around the tube, meaning there isn't significant edema in the upper airway (laryngeal area). This patency around the tube is what lowers the risk of post-extubation stridor, because with less swelling there's less tendency for the airway to narrow once the tube is removed. Conversely, little or no leak suggests substantial laryngeal edema and a higher risk of extubation failure or stridor, prompting caution or delay and possibly treatments to reduce edema. So a leak around the cuff indicates a lower likelihood of problematic upper airway edema and a reduced risk of post-extubation stridor. It does not assess tube placement, and absence of a leak does not mean the airway is never ready for extubation.

SAMPLE

7. Which condition is described as destruction of airways distal to the bronchioles, with involvement of the pulmonary capillary bed?

A. Emphysema

B. CHF

C. RSV

D. TB

Destruction of the alveolar walls and the tiny capillary network around them is the hallmark here. Emphysema involves damage to structures distal to the bronchioles, specifically the alveoli, which also takes out much of the pulmonary capillary bed that lies within those walls. This collapse of the small airways and loss of surface area for gas exchange leads to air trapping, enlarged air spaces, and reduced diffusion capacity, all classic features of emphysema. The other conditions don't fit this pattern. Congestive heart failure mainly causes fluid buildup and edema in the lungs rather than destruction of alveolar walls. RSV is a viral infection that inflames and narrows small airways (bronchiolitis) but does not cause the structural destruction of alveolar walls and capillaries. TB causes granulomatous inflammation and may cavitate, but it isn't characterized by diffuse destruction of distal airways with involvement of the capillary bed. So the description points to emphysema.

8. Which condition is associated with shortness of breath that is mostly on exertion and progressively worsens?

A. COPD

B. CHF

C. Emphysema

D. Chronic Bronchitis

Exertional dyspnea that progressively worsens is a classic sign of chronic obstructive lung disease. COPD causes a fixed, nonreversible limitation of airflow, so as the disease advances, a person becomes increasingly short of breath with activity and loses exercise tolerance over time. This pattern fits COPD well because it underlies the gradual decline in lung function seen with emphysema and chronic bronchitis, the two main components of COPD, often linked to a smoking history or other chronic lung irritants. Heart failure can also cause shortness of breath on exertion, but its typical features include orthopnea, paroxysmal nocturnal dyspnea, and edema, which help differentiate it from COPD. Emphysema and chronic bronchitis are parts of COPD, so they share the same progressive, exertional dyspnea profile, but positioning COPD as the overall chronic obstructive process captures the pattern most consistently.

9. Ataxic respirations are defined as which pattern?

A. Regular and predictable breathing

B. Pattern of irregular and unpredictable breathing

C. Rapid and shallow breathing

D. No breathing

Ataxic respirations reflect a loss of the normal respiratory rhythm due to brainstem dysfunction, producing an irregular and unpredictable pattern with varying depth and rate. This unpredictable, nonuniform breathing is the hallmark being tested. It often signals severe neurologic compromise and requires urgent airway assessment and support. Regular and predictable breathing describes a normal rhythm, rapid and shallow breathing is tachypnea, and no breathing is apnea.

SAMPLE

10. Which condition commonly presents with abdominal distension, dependent edema, and increased breathing rate?

A. CHF

B. COPD

C. Emphysema

D. Chronic Bronchitis

Abdominal distension with dependent edema signals fluid overload and venous congestion. In congestive heart failure, increased venous pressure, often from right-sided or biventricular failure, drives fluid into the interstitial spaces and peritoneal cavity, causing ascites and swelling of the legs that is worse with gravity. The faster breathing rate comes from pulmonary congestion and reduced gas exchange as the left heart struggles, leading to dyspnea and tachypnea. The lung-disease options focus on airway obstruction or parenchymal damage and don't typically produce ascites and edema, making congestive heart failure the best fit for this combination of signs.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://eeiairway.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE