EMERGENCY MEDICINE DRUGS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://emergencymeddrugs.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which adverse event may occur with excessive Albuterol administration?

A. Hyperkalemia

B. Paradoxical bronchospasm

C. Chelation

D. Renal failure

2. What is the typical IV dose for morphine when used for chest pain in the ED?

A. 1–2 mg IV

B. 4–6 mg IV

C. 2–4 mg IV; may repeat

D. 5–8 mg IV

3. Oxygen is classified as what type of substance?

A. Gas

B. Liquid

C. Aerosol

D. Solid

4. What is the initial amiodarone dose for shockable VF/pVT in ACLS, and what is the second dose if persistent?

A. 150 mg IV push; second dose 300 mg if persistent

B. 300 mg IV push; second dose 150 mg if persistent

C. 450 mg IV push; no second dose

D. 100 mg IV push; second dose 200 mg

5. Midazolam is metabolized to an active metabolite by hepatic enzymes. What is the name of this active metabolite?

A. Desmethyldiazepam

B. 1-hydroxymidazolam

C. Oxazepam

D. Midazolam-3-carboxylic acid

SAMPLE

6. What is the pregnancy category for epinephrine 1:1000?

A. Category A

B. Category D

C. Category C

D. Category X

7. In which condition should aspirin use be considered with caution?

A. Asthma

B. Hypertension

C. Diabetes

D. Hypothyroidism

8. Onset of action for albuterol is typically how fast?

A. 30-60 minutes

B. 2-15 minutes

C. Immediate

D. 1-2 hours

9. What is the typical induction dose of propofol for rapid sequence intubation (RSI)?

A. 0.3 mg/kg IV

B. 0.5 mg/kg IV

C. 1–2 mg/kg IV

D. 3–4 mg/kg IV

10. Which statement about midazolam and memory is correct?

A. It causes retrograde amnesia

B. It permanently erases memory

C. It causes anterograde amnesia

D. It has no effect on memory

SAMPLE

Answers

SAMPLE

1. B
2. C
3. A
4. B
5. B
6. C
7. A
8. B
9. C
10. C

SAMPLE

Explanations

SAMPLE

1. Which adverse event may occur with excessive Albuterol administration?

A. Hyperkalemia

B. Paradoxical bronchospasm

C. Chelation

D. Renal failure

Paradoxical bronchospasm is the adverse event to recognize with excessive albuterol. Albuterol stimulates beta-2 receptors to relax bronchial smooth muscle and open airways, but when given in too high a dose or too frequently, it can irritate the airways or trigger a reflex constriction, causing wheezing and coughing to worsen instead of improve. This paradoxical response, though uncommon, is a known potential effect of overuse or improper delivery. Hyperkalemia wouldn't fit because beta-agonists tend to drive potassium into cells, often causing hypokalemia rather than hyperkalemia. Chelation and renal failure are not typical direct adverse effects of albuterol. In practice, expect tachycardia, tremor, and possible hypokalemia with excessive use, with paradoxical bronchospasm being the specific paradoxical reaction to watch for.

2. What is the typical IV dose for morphine when used for chest pain in the ED?

A. 1–2 mg IV

B. 4–6 mg IV

C. 2–4 mg IV; may repeat

D. 5–8 mg IV

The key idea is that IV morphine for chest pain is given in small, titratable doses to relieve pain while allowing rapid reassessment. The typical starting dose is 2–4 mg IV, and you administer it, then reassess the patient's pain and vitals. If pain remains, you can give additional small doses (often 2 mg) every few minutes until relief is achieved, without pushing a large single dose all at once. This approach minimizes the risk of oversedation, respiratory depression, or hypotension while you obtain adequate analgesia. Higher single doses, such as 4–6 mg or 5–8 mg, are less desirable as starting choices because they raise the likelihood of adverse effects and reduce your ability to titrate to effect. A 1–2 mg initial dose is often insufficient for significant chest pain in many patients, underscoring why the 2–4 mg range with repeat dosing best fits typical practice.

3. Oxygen is classified as what type of substance?

A. Gas

B. Liquid

C. Aerosol

D. Solid

State of matter at room temperature and pressure determines how oxygen is classified. Under these conditions, oxygen is a gas, which is how it is stored and delivered in medical settings (compressed gas in cylinders or delivered as a breathable gas). The diatomic molecule O2 remains gaseous at room temperature, with a boiling point far below it, so it is not a liquid or solid in everyday use. An aerosol refers to tiny droplets or particles suspended in air, which is not how pure oxygen is supplied for therapy.

SAMPLE

4. What is the initial amiodarone dose for shockable VF/pVT in ACLS, and what is the second dose if persistent?

A. 150 mg IV push; second dose 300 mg if persistent

B. 300 mg IV push; second dose 150 mg if persistent

C. 450 mg IV push; no second dose

D. 100 mg IV push; second dose 200 mg

When a shockable rhythm like VF or pulseless VT doesn't respond to defibrillation, amiodarone is used as an antiarrhythmic to help terminate the rhythm and prevent recurrence. The fastest way to achieve therapeutic levels is a rapid 300 mg IV push as the first dose. If the rhythm persists after that, a second, smaller 150 mg IV push is given to increase the chances of conversion without delaying further treatment. This approach—300 mg upfront, then 150 mg if persistence or recurrence occurs—is preferred because it delivers a substantial early loading dose to stabilize the myocardium, with a subsequent dose to sustain effect. After the arrest, an infusion may be started to maintain antiarrhythmic activity (commonly 1 mg/min for several hours, then tapering), but the initial and second bolus dosing is the key point for shockable VF/pVT. Dosing options that start with a smaller initial amount or swap the order of the doses do not align with these guidelines.

5. Midazolam is metabolized to an active metabolite by hepatic enzymes. What is the name of this active metabolite?

A. Desmethyldiazepam

B. 1-hydroxymidazolam

C. Oxazepam

D. Midazolam-3-carboxylic acid

Midazolam is processed in the liver by cytochrome P450 enzymes, mainly CYP3A4, which convert it to 1-hydroxymidazolam. This metabolite is pharmacologically active and contributes to the sedative effects, sometimes extending the duration of action, especially with repeated dosing or hepatic impairment. It's later conjugated and eliminated, but its activity helps explain why midazolam can have longer-lasting effects than expected. Other benzodiazepine metabolites listed are not derived from midazolam, and the midazolam-3-carboxylic acid form is inactive.

6. What is the pregnancy category for epinephrine 1:1000?

A. Category A

B. Category D

C. Category C

D. Category X

The main idea here is how the FDA pregnancy categories describe fetal risk. Category C means that animal studies have shown potential fetal harm and there aren't enough well-controlled human studies, but the drug may still be used if the potential benefits to the mother (and thus to the fetus) outweigh the risks. Epinephrine 1:1000 is used in life-threatening situations like anaphylaxis. In such emergencies, stabilizing the mother is the priority because maternal shock and poor perfusion pose a far greater threat to the fetus than the potential fetal risk from the drug. That makes epinephrine fall into Category C: not without risk, but justifiable and necessary when benefits exceed potential harms.

SAMPLE

7. In which condition should aspirin use be considered with caution?

A. Asthma

B. Hypertension

C. Diabetes

D. Hypothyroidism

Aspirin should be used with caution in asthma because some people with asthma have aspirin-exacerbated respiratory disease. When aspirin (and other NSAIDs) inhibit COX enzymes, arachidonic acid is shunted toward the leukotriene pathway, and leukotrienes are potent bronchoconstrictors. This can trigger wheezing, coughing, and shortness of breath, especially in individuals with asthma and nasal polyps. Hypertension, diabetes, and hypothyroidism don't carry this specific risk of bronchospasm from aspirin, so they don't inherently require caution for this reason. If analgesia is needed in someone with asthma, consider acetaminophen or discuss NSAID desensitization only if aspirin is truly necessary for another condition.

8. Onset of action for albuterol is typically how fast?

A. 30-60 minutes

B. 2-15 minutes

C. Immediate

D. 1-2 hours

The key point is how quickly albuterol produces bronchodilation after inhalation. Inhaled short-acting beta-agonists act directly on β2 receptors in the airway smooth muscle, delivering rapid relief with minimal delay. Because of this direct delivery, the onset is measured in minutes, not hours, and can be as fast as a couple of minutes to around a dozen minutes depending on the device and patient technique. That's why a window of about 2–15 minutes best fits the typical onset. Options suggesting an effect that begins in 30–60 minutes or 1–2 hours are too slow for acute relief. Describing it as immediate would be overstating the speed, since even rapid inhaled bronchodilators take a short moment to onset.

9. What is the typical induction dose of propofol for rapid sequence intubation (RSI)?

A. 0.3 mg/kg IV

B. 0.5 mg/kg IV

C. 1–2 mg/kg IV

D. 3–4 mg/kg IV

When preparing for rapid sequence intubation, you want a propofol dose that reliably produces loss of consciousness quickly without causing excessive blood pressure drop. The induction dose is typically 1–2 mg/kg given IV in adults. This range achieves rapid unconsciousness within seconds, facilitating a smooth intubation. Doses much lower than this are unlikely to deepen anesthesia adequately for RSI and may require additional agents, while doses higher than this raise the risk of significant hypotension, apnea, and potential airway complications. In practice, the exact amount can be adjusted based on the patient's hemodynamic status and comorbidities, and adjuncts like analgesia (for example fentanyl) may be used to blunt the hemodynamic response to laryngoscopy.

SAMPLE

10. Which statement about midazolam and memory is correct?

A. It causes retrograde amnesia

B. It permanently erases memory

C. It causes anterograde amnesia

D. It has no effect on memory

Midazolam's memory effect is primarily anterograde amnesia: during the time the drug is active, encoding of new information is impaired, so the patient forgets events that occur after administration. This happens because benzodiazepines enhance GABA activity, dampening hippocampal encoding of new memories. The impairment is transient—memories formed before administration are preserved, and once the drug wears off, the ability to form new memories returns. It does not permanently erase memory, and it does not typically cause true retrograde amnesia.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://emergencymeddrugs.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE