
EMERGENCY MEDICINE IN-TRAINING EXAMINATION (EM-ITE) PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://emergencymedicineite.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is one of the T's that can cause pulseless electrical activity (PEA)?

A. Tension pneumothorax

B. Traumatic cardiac arrest

C. Transfusion reaction

D. Thyroid storm

2. What is the recommended treatment for a patient diagnosed with Chlamydia Trachomatis and sterile pyuria?

A. Amoxicillin

B. Acyclovir

C. Azithromycin and Ceftriaxone

D. Metronidazole

3. What is the treatment of choice for antidromic WPW?

A. Beta-blockers

B. Adenosine

C. Procainamide

D. Calcium channel blockers

4. What is a key symptom of nitrogen dioxide pneumonitis?

A. Hemoptysis

B. Chest pain

C. Lightheadedness

D. All of the above

5. What is the first-line treatment for hyperemesis gravidarum?

A. Dexamethasone

B. Pyridoxine and doxylamine

C. Zofran

D. Metoclopramide

SAMPLE

6. What condition is indicated by elevated alkaline phosphatase and GGT post liver transplant with normal bilirubin and aminotransferases?

A. Preservation injury

B. Acute cellular injury

C. Bile duct obstruction

D. Cholestasis

7. What ultrasound finding is expected for a glass foreign body?

A. Hypoechoic object without shadowing

B. Hyperechoic object with shadowing

C. Complex cystic mass

D. Mosaic pattern

8. What are the criteria for diagnosing spontaneous bacterial peritonitis in patients undergoing peritoneal dialysis?

A. Abdominal pain, positive culture

B. Cloudy dialysate, abdominal pain

C. Peritoneal fluid with at least 100 WBCs/mm and positive culture

D. Hepatic encephalopathy, cloudy dialysate

9. What is a common symptom of salicylate toxicity?

A. Bradycardia

B. Hyperpnea

C. Hypotension

D. Seizures

10. Which acid-base disorder is typically seen in salicylate toxicity?

A. Metabolic alkalosis

B. Respiratory acidosis

C. Respiratory alkalosis

D. Metabolic acidosis

SAMPLE

Answers

SAMPLE

1. A
2. C
3. C
4. D
5. B
6. A
7. B
8. C
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What is one of the T's that can cause pulseless electrical activity (PEA)?

A. Tension pneumothorax

B. Traumatic cardiac arrest

C. Transfusion reaction

D. Thyroid storm

Tension pneumothorax is a critical condition that can lead to pulseless electrical activity (PEA). In a tension pneumothorax, air becomes trapped in the pleural space, creating increasing pressure that can collapse the lung on the affected side and significantly impede venous return to the heart. This increased intrathoracic pressure can compromise cardiovascular function, leading to reduced cardiac output and the inability to generate effective circulation, even when electrical activity is present on the ECG. Tension pneumothorax can manifest quickly, and if not recognized and treated promptly through decompression (e.g., needle thoracostomy), it can rapidly become life-threatening. The mechanism by which it causes PEA primarily revolves around the impaired filling of the heart due to reduced venous return and overall obstruction of the great vessels. In this context, while traumatic cardiac arrest can indeed lead to PEA, it is primarily due to associated conditions (like hypovolemia or massive hemorrhage) rather than a direct physiologic obstruction as seen with tension pneumothorax. Other choices, such as transfusion reactions and thyroid storm, do not directly lead to PEA in the acute setting and typically involve different pathophysiological processes.

2. What is the recommended treatment for a patient diagnosed with Chlamydia Trachomatis and sterile pyuria?

A. Amoxicillin

B. Acyclovir

C. Azithromycin and Ceftriaxone

D. Metronidazole

The recommended treatment for a patient diagnosed with Chlamydia trachomatis and sterile pyuria is a combination of azithromycin and ceftriaxone. This approach is appropriate because Chlamydia is a common cause of sterile pyuria, where the urine shows white blood cells but the culture for bacteria is negative, indicating an infection without the presence of traditional urinary pathogens. Azithromycin effectively treats Chlamydia trachomatis, providing good tissue penetration and a single-dose regimen that enhances adherence. Ceftriaxone is included as it covers potential concurrent gonococcal infection, which is clinically relevant since co-infection with Neisseria gonorrhoeae is common in patients with Chlamydial infections. This regimen addresses both the Chlamydia infection and provides a safety net for possible co-infections while ensuring an effective treatment outcome. The inclusion of both antibiotics accommodates the need to cover potential sexually transmitted infections comprehensively and aligns with recommended guidelines for treating such infections, particularly in sexually active individuals. In contrast, amoxicillin is not the first-line treatment for Chlamydia; acyclovir is used mainly for herpes simplex virus infections rather than Chlamydia; and metronidazole is effective for anaerobic bacteria and proto

SAMPLE

3. What is the treatment of choice for antidromic WPW?

A. Beta-blockers

B. Adenosine

C. Procainamide

D. Calcium channel blockers

In the context of treating antidromic Wolff-Parkinson-White (WPW) syndrome, the choice of Procainamide is well-supported due to its ability to effectively slow down conduction through accessory pathways. Antidromic WPW occurs when the electrical impulse flows from the ventricles back to the atria via the accessory pathway, which is the opposite of orthodromic conduction. Procainamide, being a Class IA antiarrhythmic, is particularly effective in managing this condition. It works by prolonging the action potential and refractory period in cardiac tissue, which decreases the conduction velocity in the accessory pathway. This can help in converting the arrhythmia back to normal sinus rhythm or at the very least, in controlling the rapid heart rate caused by antidromic conduction. Other options may not be appropriate for managing this specific type of arrhythmia. Beta-blockers can help in some supraventricular tachycardias but are generally avoided in WPW due to the risk of promoting rapid conduction through the accessory pathway. Adenosine is effective for terminating orthodromic WPW but is less effective in antidromic cases. Calcium channel blockers can also increase conduction through the accessory pathway in antidromic

4. What is a key symptom of nitrogen dioxide pneumonitis?

A. Hemoptysis

B. Chest pain

C. Lightheadedness

D. All of the above

Nitrogen dioxide pneumonitis is a lung injury caused by inhalation of nitrogen dioxide (NO_2), a gas that can be released during combustion processes such as those found in vehicle exhaust or certain industrial activities. The key symptoms associated with nitrogen dioxide exposure, particularly in cases of pneumonitis, can include hemoptysis, chest pain, and lightheadedness. Hemoptysis, or coughing up blood, can occur due to the irritation and inflammation of the airways and lung tissue caused by nitrogen dioxide exposure, leading to damage to the delicate blood vessels within the lungs. Chest pain may arise from inflammation or damage to the lung tissues and structures, resulting in pain that can mimic other respiratory and cardiovascular issues. This symptom is particularly notable as the lungs and pleura respond to irritants, causing discomfort and pain during respiratory efforts. Lightheadedness can result from a couple of mechanisms. It may be a direct consequence of respiratory distress or inadequate oxygen exchange due to impaired lung function after exposure to nitrogen dioxide. This reduced oxygenation can also result in cerebral hypoxia, leading to feelings of dizziness or lightheadedness. Recognizing that nitrogen dioxide pneumonitis can present with a combination of these symptoms is essential in clinical practice. Therefore, the

SAMPLE

5. What is the first-line treatment for hyperemesis gravidarum?

A. Dexamethasone

B. Pyridoxine and doxylamine

C. Zofran

D. Metoclopramide

The first-line treatment for hyperemesis gravidarum is a combination of pyridoxine (vitamin B6) and doxylamine. This approach is based on the fact that these medications have shown efficacy in alleviating nausea and vomiting during pregnancy, particularly in hyperemesis gravidarum, which is a severe form of morning sickness characterized by persistent vomiting, dehydration, and weight loss. Pyridoxine is a safe and well-tolerated vitamin that has been found to reduce nausea in pregnant individuals. Doxylamine, an antihistamine, can also help mitigate symptoms of nausea and vomiting. The combination of these two drugs is often effective, and it is recommended because of its safety profile for both the mother and the fetus. In comparison, other treatments may be utilized for hyperemesis gravidarum, but they are not considered first-line. Dexamethasone, while potentially effective for severe cases, is used more cautiously due to concerns about potential risks in pregnancy. Zofran (ondansetron) is often used for nausea and vomiting in various contexts, but it is typically not considered first-line for hyperemesis gravidarum, and there are concerns regarding its use in pregnancy. Metoc

6. What condition is indicated by elevated alkaline phosphatase and GGT post liver transplant with normal bilirubin and aminotransferases?

A. Preservation injury

B. Acute cellular injury

C. Bile duct obstruction

D. Cholestasis

The scenario of elevated alkaline phosphatase and GGT following a liver transplant, combined with normal bilirubin and aminotransferases, is indicative of an issue related to bile flow rather than liver parenchymal damage or acute cellular injury. In this case, elevated alkaline phosphatase and GGT suggest a cholestatic pattern, which points towards a disruption in bile flow. Bile duct obstruction is often characterized by elevated alkaline phosphatase but typically also presents with increased bilirubin levels due to accumulation of bile acids backing up into the liver and bloodstream. Normal bilirubin levels in this scenario suggest that while there may be cholestasis, it is not due to obstruction of the bile duct but rather a condition that is affecting the bile flow. Cholestasis can occur post-liver transplant as a result of preservation injury, particularly if there was an ischemic or a perfusion problem with the transplanted liver. Preservation injury occurs when there is damage due to factors surrounding the procurement and storage of the liver organ before transplantation, affecting the function of bile ducts and bile production. This context helps to clarify that while bile duct obstruction manifests similarly with elevated alkaline phosphatase, the presence of normal bilirubin combined with the other findings suggests that

SAMPLE

7. What ultrasound finding is expected for a glass foreign body?

A. Hypoechoic object without shadowing

B. Hyperechoic object with shadowing

C. Complex cystic mass

D. Mosaic pattern

The expected ultrasound finding for a glass foreign body is a hyperechoic object with shadowing. When glass is imaged using ultrasound, it appears as a bright white reflection due to its high acoustic impedance compared to surrounding soft tissues. This hyperechoic appearance indicates that the glass is dense, reflecting a significant amount of the ultrasound waves. Additionally, the presence of shadowing beneath the glass foreign body occurs because the dense material absorbs and reflects the sound waves, causing a lack of echogenicity in the deeper tissues behind it. This phenomenon helps differentiate glass from other objects, as many soft tissues do not produce such shadowing effects. Recognizing these ultrasound characteristics is critical in emergency medicine when diagnosing and managing patients with potential foreign body injuries. In contrast, other findings like a hypoechoic object without shadowing would not accurately represent the dense nature of glass, while a complex cystic mass and a mosaic pattern are indicative of entirely different conditions or materials, such as fluid collections or certain types of masses in the body.

8. What are the criteria for diagnosing spontaneous bacterial peritonitis in patients undergoing peritoneal dialysis?

A. Abdominal pain, positive culture

B. Cloudy dialysate, abdominal pain

C. Peritoneal fluid with at least 100 WBCs/mm and positive culture

D. Hepatic encephalopathy, cloudy dialysate

The diagnosis of spontaneous bacterial peritonitis (SBP) in patients undergoing peritoneal dialysis relies on specific laboratory and clinical findings in the peritoneal fluid. The correct criteria state that SBP is indicated by the presence of at least 100 white blood cells (WBCs) per millimeter cubed of peritoneal fluid, with a positive culture for bacteria. This is important because the elevation in WBC count signifies an inflammatory response to infection within the peritoneal cavity, while a positive culture confirms the presence of a bacterial pathogen. The threshold of 100 WBCs/mm³ is significant because it indicates a level of infection that warrants treatment, and monitoring for bacterial peritonitis is essential in patients on peritoneal dialysis due to their predisposition to infections. Other options may include symptoms or nonspecific findings, but they do not meet the diagnostic criteria necessary for confirming SBP. For instance, abdominal pain and cloudy dialysate can occur in multiple scenarios and aren't definitive alone without the quantitative and culture data. Thus, the precise combination of WBC count and culture positivity is crucial for an accurate diagnosis of SBP in this patient population.

SAMPLE

9. What is a common symptom of salicylate toxicity?

A. Bradycardia

B. Hyperpnea

C. Hypotension

D. Seizures

Salicylate toxicity is characterized by a range of symptoms, one of the most prominent being hyperpnea, which refers to an increased rate and depth of breathing. This respiratory response occurs as the body attempts to compensate for metabolic acidosis—a primary metabolic derangement seen in salicylate toxicity. Salicylates can stimulate the respiratory center in the brain, leading to hyperventilation in an effort to blow off carbon dioxide and correct the acid-base imbalance. Additionally, hyperpnea may be accompanied by other central nervous system effects, such as tinnitus and altered mental status, but the hyperventilation itself is a key feature that distinguishes salicylate toxicity from other forms of poisoning. Understanding hyperpnea in the context of salicylate toxicity is vital, as it helps clinicians identify and manage the condition effectively, taking appropriate steps to rectify the underlying metabolic issues while addressing the respiratory symptoms.

10. Which acid-base disorder is typically seen in salicylate toxicity?

A. Metabolic alkalosis

B. Respiratory acidosis

C. Respiratory alkalosis

D. Metabolic acidosis

In salicylate toxicity, respiratory alkalosis is typically observed due to overstimulation of the respiratory center in the brain by salicylates. This stimulation leads to hyperventilation, resulting in excessive elimination of carbon dioxide (CO_2) from the body, which elevates blood pH and creates an alkalotic state. As salicylate levels rise, they can cause both a primary respiratory alkalosis and a metabolic acidosis due to the accumulation of organic acids, such as lactate, which may contribute to the acid-base disturbances observed. However, the initial and predominant effect seen with salicylate overdose is respiratory alkalosis, arising from that hyperventilation response. This phenomenon is an important consideration in the management of patients with suspected salicylate poisoning, making it crucial for emergency providers to recognize these hallmark changes in order to properly assess and treat the condition.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://emergencymedicineite.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE