EMT OBSTETRICS AND NEONATAL CARE PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://emtobstetricsneonatalcare.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. If a pregnant woman experiences vaginal bleeding, what should be done?

A. Perform a vaginal exam immediately

B. Monitor and assess the patient's condition and transport to a hospital

C. Give the patient medication for pain relief

D. Advise the patient to stay home and rest

2. If a newborn is born with a limp, what should your first action be?

A. Provide warmth and assess the airway

B. Begin resuscitation immediately

C. Notify the neonatal team for immediate evaluation

D. Monitor the newborn for a few minutes

3. When should you provide oxygen to a newborn immediately after delivery?

A. if the newborn is apneic

B. if the newborn has a high heart rate

C. if the newborn is crying

D. if the newborn is of low birth weight

4. What role does oxytocin serve during labor?

A. It allows for fetal movement

B. It stimulates uterine contractions

C. It relaxes the cervix

D. It decreases pain perception

5. What type of assessments are typically carried out immediately after delivery?

A. Assessing the mother's mental health

B. Checking the newborn's APGAR score and vital signs

C. Determining the mother's food preferences

D. Estimating the delivery time

6. What is a common sign of labor approaching?

A. Frequent headaches

B. Increased fetal movement

C. Regular contractions and cervical dilation

D. A decrease in appetite

SAMPLE

7. How is the fundal height of an expecting mother measured?

A. By measuring from the pubic symphysis to the top of the uterus

B. By measuring from the navel to the top of the uterus

C. By measuring the circumference of the abdomen

D. By assessing the distance from the rib cage to the pubic bone

8. Which is the appropriate technique for opening the airway of an infant with no suspected neck injury?

A. Lift the chin and hyperextend the neck

B. Tilt the head back without hyperextending the neck

C. Gently lift the chin while maintaining slight flexion

D. Perform as you would for an older child or adult

9. What is a common fetal position that may require intervention during delivery?

A. Anterior

B. Transverse

C. Occipito-posterior

D. Breech

10. Which condition is most often monitored for in a high-risk pregnancy?

A. Gestational diabetes

B. Normal fetal position

C. Postpartum bleeding

D. Preeclampsia

SAMPLE

Answers

SAMPLE

1. B
2. A
3. A
4. B
5. B
6. C
7. A
8. B
9. C
10. D

SAMPLE

Explanations

SAMPLE

1. If a pregnant woman experiences vaginal bleeding, what should be done?

A. Perform a vaginal exam immediately

B. Monitor and assess the patient's condition and transport to a hospital

C. Give the patient medication for pain relief

D. Advise the patient to stay home and rest

Monitoring and assessing the patient's condition while preparing for transportation to a hospital is crucial in the case of vaginal bleeding during pregnancy. Vaginal bleeding can signify a range of potential complications, including miscarriage, placental abruption, or placenta previa, all of which require immediate medical attention. By monitoring vital signs and understanding the extent and nature of the bleeding, responders can gather important information that can assist professionals at the hospital. Transporting the patient to a medical facility ensures that she receives the appropriate evaluation and treatment, which can be vital for her health and the health of the fetus. Performing a vaginal exam immediately may introduce risk and is typically not advised in a prehospital setting without specific training or indications, as it could worsen any existing conditions. Providing pain relief medication could mask important symptoms and is not standard practice without professional guidance. Advising the patient to rest at home overlooks the urgency of addressing potential complications and could lead to serious outcomes for both the mother and baby. Thus, the most prudent course of action involves careful assessment and timely transport to ensure that the patient receives necessary medical care.

2. If a newborn is born with a limp, what should your first action be?

A. Provide warmth and assess the airway

B. Begin resuscitation immediately

C. Notify the neonatal team for immediate evaluation

D. Monitor the newborn for a few minutes

When a newborn is born with a limp, the most immediate and crucial response involves providing warmth and assessing the airway. Newborns are particularly vulnerable to hypothermia, so ensuring they maintain a stable body temperature is essential. Additionally, a limp presentation may indicate a compromised airway, which could hinder their ability to breathe effectively. By assessing the airway first, you can ensure that there are no obstructions or other issues that could further impair their respiratory function. Establishing warmth and securing a patent airway are foundational steps that can stabilize the newborn before more advanced interventions are considered. If the newborn shows signs of severe distress or unresponsiveness during these initial actions, then further steps such as resuscitation may become necessary, but stabilization through warmth and airway assessment remains the priority.

SAMPLE

3. When should you provide oxygen to a newborn immediately after delivery?

A. if the newborn is apneic

B. if the newborn has a high heart rate

C. if the newborn is crying

D. if the newborn is of low birth weight

Providing oxygen to a newborn immediately after delivery is crucial in situations where the infant is apneic, meaning they are not breathing spontaneously. In these cases, immediate intervention is essential to ensure adequate oxygenation to the brain and other vital organs, as the newborn is at high risk for hypoxia and related complications. The lack of breathing can lead to a critical situation requiring supplemental oxygen or positive pressure ventilation to establish effective breathing and support the newborn's transition to life outside the womb. In contrast, a newborn with a high heart rate typically indicates that they are responding well and may not require immediate oxygen. Likewise, if the newborn is crying, it is a positive sign of effective respiratory effort, suggesting that oxygen is being adequately delivered naturally. Additionally, low birth weight alone does not warrant immediate oxygen administration unless accompanied by other concerning signs such as apnea or poor respiratory effort. Thus, the correct scenario for providing supplemental oxygen immediately after delivery focuses on the need to address apnea.

4. What role does oxytocin serve during labor?

A. It allows for fetal movement

B. It stimulates uterine contractions

C. It relaxes the cervix

D. It decreases pain perception

Oxytocin plays a crucial role during labor, primarily by stimulating uterine contractions. This hormone is released from the posterior pituitary gland and is essential for the progression of labor. When labor begins, oxytocin levels rise, causing the uterus to contract more forcefully and rhythmically. These contractions help to dilate the cervix, which is vital for the descent of the fetus through the birth canal. Additionally, the release of oxytocin increases in response to the stretching of the cervix and vaginal walls during labor, creating a positive feedback loop that intensifies contractions as labor progresses. This mechanism is fundamental for facilitating childbirth and ensuring that the baby is delivered effectively and safely. Therefore, its role in stimulating uterine contractions is critical for the labor process.

SAMPLE

5. What type of assessments are typically carried out immediately after delivery?

A. Assessing the mother's mental health

B. Checking the newborn's APGAR score and vital signs

C. Determining the mother's food preferences

D. Estimating the delivery time

Checking the newborn's APGAR score and vital signs is essential immediately after delivery as it provides critical information about the infant's immediate health status. The APGAR score is a quick assessment performed one and five minutes after birth, evaluating five areas: appearance (skin color), pulse (heart rate), grimace response (reflexes), activity (muscle tone), and respiratory effort. Each category is scored from 0 to 2, with a total score ranging from 0 to 10, which helps determine if the newborn needs immediate medical attention. In addition to the APGAR score, monitoring vital signs such as heart rate, respiratory rate, and temperature is crucial to ensure the newborn is transitioning well to life outside the womb. Any abnormal readings can indicate the need for further intervention or assessment. Collectively, these assessments help healthcare providers gauge the newborn's overall well-being and establish a baseline for further monitoring. The other options, while important in their context, do not have the same immediate relevance as evaluating the newborn's health right after delivery. Assessing the mother's mental health is essential but typically occurs later during postpartum care. Determining the mother's food preferences and estimating the delivery time are also not immediate priorities when the primary focus should be on the

6. What is a common sign of labor approaching?

A. Frequent headaches

B. Increased fetal movement

C. Regular contractions and cervical dilation

D. A decrease in appetite

Regular contractions and cervical dilation are common signs that labor is approaching. As the body prepares for childbirth, the uterus begins to contract in a regular pattern, which helps to thin and open the cervix (a process known as cervical dilation). These contractions can vary in intensity and duration but typically become more frequent and closer together as labor progresses. Cervical changes, such as effacement and dilation, are essential indicators that the body is getting ready for delivery. In contrast, frequent headaches, increased fetal movement, and a decrease in appetite are not direct or reliable indicators of labor nearing. While some women might experience headaches or changes in appetite as a result of hormonal fluctuations or discomfort during pregnancy, these symptoms do not signify that labor is imminent. Increased fetal movement is usually normal throughout pregnancy and does not specifically indicate labor; it may vary from day to day and can be influenced by numerous factors, including the mother's activity levels and the baby's position.

SAMPLE

7. How is the fundal height of an expecting mother measured?

A. By measuring from the pubic symphysis to the top of the uterus

B. By measuring from the navel to the top of the uterus

C. By measuring the circumference of the abdomen

D. By assessing the distance from the rib cage to the pubic bone

The fundal height is measured by assessing the distance from the pubic symphysis to the top of the uterus, also known as the fundus. This measurement is important during pregnancy as it provides healthcare providers with information about fetal growth and the gestational age of the baby. Typically, fundal height is expected to increase in a predictable manner as the pregnancy progresses; this helps in monitoring the health and development of the fetus. Measuring from the pubic symphysis to the top of the uterus is an established practice because it reflects the growth of the uterus, which rises up into the abdominal cavity as the pregnancy advances. Each centimeter of fundal height generally correlates with weeks of gestation after the 20th week, making it a practical tool for estimating the progress of the pregnancy. Other methods of measurement, while they may provide some data about the abdomen, do not offer the specific insights that fundal height does regarding fetal growth and gestational age. Therefore, the chosen method of measuring fundal height is crucial in prenatal care.

8. Which is the appropriate technique for opening the airway of an infant with no suspected neck injury?

A. Lift the chin and hyperextend the neck

B. Tilt the head back without hyperextending the neck

C. Gently lift the chin while maintaining slight flexion

D. Perform as you would for an older child or adult

The appropriate technique for opening the airway of an infant with no suspected neck injury involves tilting the head back without hyperextending the neck. In infants, the airway is naturally positioned more anteriorly (forward), and their anatomy requires a more gentle approach to avoid obstruction and to ensure proper ventilation. Tilting the head back allows for the alignment of the airway structures to facilitate airflow but does not involve hyperextension, which can actually lead to airway compression due to the flexibility of an infant's trachea. Maintaining a neutral position or a slight tilting aligns the airway effectively while also being mindful that infants have more short necks and larger heads relative to their bodies, making them more susceptible to airway obstruction if positioned incorrectly. In contrast, lifting the chin and hyperextending the neck, or performing techniques suitable for older children or adults, does not account for the unique anatomical differences in infants. Gently lifting the chin while maintaining slight flexion is close but still would not be the most effective approach compared to the option of tilting the head back without hyperextension. Thus, the chosen method ensures both safety and effectiveness in managing an infant's airway.

SAMPLE

9. What is a common fetal position that may require intervention during delivery?

A. Anterior

B. Transverse

C. Occipito-posterior

D. Breech

The occipito-posterior position is a common fetal position that can necessitate intervention during delivery due to complications associated with this orientation. In this position, the baby's back is facing the mother's back, which can lead to a longer and more difficult labor. This positioning may result in increased pain for the mother and a higher chance of fetal distress, particularly if the labor is prolonged. Additionally, the occipito-posterior position can lead to a greater likelihood of delivery interventions such as the use of forceps or vacuum extraction, and in some cases, a cesarean section may be necessary if the baby does not rotate to a more favorable position. The goal of managing this position typically involves monitoring labor progress closely and potentially utilizing specific maneuvers to encourage the baby to turn to a more favorable anterior position, which is ideal for delivery. Other positions, such as breech, are also common fetal orientations that can lead to intervention, but the occipito-posterior position specifically illustrates the challenges it presents during labor and delivery, highlighting the need for timely assessment and action by obstetric caregivers.

10. Which condition is most often monitored for in a high-risk pregnancy?

A. Gestational diabetes

B. Normal fetal position

C. Postpartum bleeding

D. Preeclampsia

In a high-risk pregnancy, preeclampsia is a condition that is most often monitored due to its potential to lead to serious complications for both the mother and the fetus. Preeclampsia is characterized by high blood pressure and often accompanied by protein in the urine, and it can develop after the 20th week of pregnancy. If left untreated, it can progress to severe complications, including eclampsia, which involves seizures, and can also result in placental abruption and fetal growth restriction. Monitoring for preeclampsia is critical because early detection and management can significantly reduce the risk of complications. Regular checks for blood pressure, urine protein levels, and overall maternal and fetal well-being are fundamental components of prenatal care in high-risk situations. While gestational diabetes is also a concern and requires monitoring, as does the fetal position, they are typically not monitored as frequently or with the same level of urgency as preeclampsia. Postpartum bleeding, while vital to address after delivery, is not a condition monitored throughout the pregnancy itself, which makes it less relevant in the context of high-risk pregnancy evaluations.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://emtobstetricsneonatalcare.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE