EXAM MASTER NEURO PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://exammasterneuro.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is a complication that occurs during the acute phase of Chagas disease?

A. Meningoencephalitis

B. Chronic cardiomyopathy

C. Megaesophagus

D. Thromboembolism

2. After a generalized tonic-clonic seizure, what should be done next if the patient does not regain consciousness within 30 minutes?

A. Monitor for an additional hour

B. Administer IV fluids

C. Start Lorazepam

D. Conduct a CT scan

3. What physiological change is most often associated with the use of Valproic acid for seizures?

A. Accelerated neuronal firing

B. Inhibition of GABA metabolism

C. Increased glutamate release

D. Increased sodium channel activity

4. Which diagnostic sign is most likely to be positive in a patient with signs of meningeal irritation?

A. Kernig's sign

B. Babinski's sign

C. Chvostek's sign

D. Quincke's sign

5. What is the typical presentation of a tension-type headache?

A. Unilateral pain, often with nausea

B. Bilateral, "tight" feeling without nausea

C. Rapid onset with intense pain and photophobia

D. Severe, stabbing pain around the eye

SAMPLE

6. Which drug is likely to be used in the Tensilon test for diagnosing myasthenia gravis?

A. Edrophonium

B. Aminoglycosides

C. Corticosteroids

D. Neostigmine

7. Which area of the brain is most likely affected in a patient with dysarthria, tongue deviation to the right, and right arm weakness?

A. Right brain stem

B. Left cerebral cortex

C. Left brain stem

D. Right cerebellum

8. What is the first-line treatment for a patient diagnosed with complex regional pain syndrome?

A. Corticosteroids

B. Physical therapy

C. Naprosyn (NSAIDs)

D. Gabapentin

9. What cerebrospinal fluid (CSF) finding is most commonly associated with Guillain-Barré syndrome?

A. Decreased CSF glucose

B. Elevated CSF protein content

C. Normal white blood cell count

D. Presence of bacterial infection markers

10. In the case of suspected viral encephalitis, what is the first imaging study recommended before lumbar puncture?

A. MRI of the brain

B. CT scan of the brain

C. PET scan

D. X-ray of the skull

SAMPLE

Answers

SAMPLE

1. A
2. C
3. B
4. A
5. B
6. A
7. C
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is a complication that occurs during the acute phase of Chagas disease?

A. Meningoencephalitis

B. Chronic cardiomyopathy

C. Megaesophagus

D. Thromboembolism

In the context of Chagas disease, the acute phase is characterized by a variety of symptoms and complications, one of which can be meningoencephalitis. This complication arises due to the central nervous system involvement by the Trypanosoma cruzi parasite, leading to inflammation of the brain and its protective membranes. During the acute phase, the immune response is activated, and the presence of the parasite can cause neurological symptoms, and in some cases, a severe inflammatory response can culminate in meningoencephalitis. While other complications such as chronic cardiomyopathy and megaesophagus are significant manifestations of the disease, they tend to occur later in the chronic phase. Thromboembolism can occur due to various factors but is not specifically a primary complication of the acute phase of Chagas disease. Therefore, during the acute phase, meningoencephalitis is the notable complication linked to direct parasitic involvement in the nervous system.

2. After a generalized tonic-clonic seizure, what should be done next if the patient does not regain consciousness within 30 minutes?

A. Monitor for an additional hour

B. Administer IV fluids

C. Start Lorazepam

D. Conduct a CT scan

In the scenario where a patient does not regain consciousness after 30 minutes following a generalized tonic-clonic seizure, administering Lorazepam is crucial. This medication belongs to the benzodiazepine class, which is effective for terminating active seizures and is commonly used in cases of status epilepticus—defined as continuous or repeated seizures without recovery. After a prolonged seizure, the risk of progressing to status epilepticus increases, so prompt administration of Lorazepam helps to quickly control seizure activity and protect the patient from further neurological impairment or complications. The management protocol emphasizes quick reversal of seizure activity, and Lorazepam's rapid onset makes it the preferred choice in this situation. Other interventions like monitoring, providing IV fluids, or conducting a CT scan would follow after initial seizure control has been achieved, but they do not address the immediate concern of ongoing seizure activity and the patient's lack of consciousness. Therefore, utilizing Lorazepam effectively prioritizes the patient's safety and neurological recovery.

SAMPLE

3. What physiological change is most often associated with the use of Valproic acid for seizures?

A. Accelerated neuronal firing

B. Inhibition of GABA metabolism

C. Increased glutamate release

D. Increased sodium channel activity

Valproic acid, also known as valproate, is primarily used as an anticonvulsant to manage seizures. One of its key mechanisms of action is the inhibition of gamma-aminobutyric acid (GABA) metabolism. GABA is the main inhibitory neurotransmitter in the central nervous system, and increasing its levels enhances inhibitory signaling in the brain, thereby stabilizing neuronal activity and preventing seizures. By inhibiting the metabolism of GABA, valproic acid causes an increase in GABA concentrations. This increased availability promotes greater inhibitory effects on neuronal firing, which is crucial since excessive neuronal excitability can lead to seizures. This action contrasts with other options; for instance, accelerated neuronal firing, increased glutamate release, and increased sodium channel activity would all contribute to enhanced excitability or neuronal firing rather than inhibition, which is contrary to the therapeutic goal of treating seizures. Thus, the role of valproic acid in GABA metabolism is central to its effectiveness in seizure management.

4. Which diagnostic sign is most likely to be positive in a patient with signs of meningeal irritation?

A. Kernig's sign

B. Babinski's sign

C. Chvostek's sign

D. Quincke's sign

When evaluating a patient with signs of meningeal irritation, Kernig's sign is the most relevant diagnostic sign likely to be positive. Kernig's sign is assessed by flexing the hip and knee and then attempting to extend the knee while the hip is flexed. Positive findings occur when the patient experiences resistance or pain during knee extension due to irritation of the meninges, which is commonly seen in conditions such as meningitis. This sign correlates well with the presence of meningeal irritation as it demonstrates a reaction of the nervous system to the inflammation that accompanies such conditions. Overall, the presence of Kernig's sign serves as an important clinical finding in the diagnosis of meningeal irritation, guiding further diagnostic and treatment steps. In contrast, Babinski's sign evaluates the integrity of the corticospinal tract and is not indicative of meningeal irritation. Chvostek's sign is related to neuromuscular excitability often associated with hypocalcemia and does not pertain to meningeal pathology. Quincke's sign, although historically referenced, is less commonly used today and primarily refers to a specific finding in patients with certain vascular conditions rather than directly pointing to meningeal irritation.

SAMPLE

5. What is the typical presentation of a tension-type headache?

A. Unilateral pain, often with nausea

B. Bilateral, "tight" feeling without nausea

C. Rapid onset with intense pain and photophobia

D. Severe, stabbing pain around the eye

Tension-type headache is characterized by a bilateral, pressing or tightening sensation that often feels like a band around the head. This type of headache generally does not present with significant accompanying symptoms such as nausea or vomiting, distinguishing it from other headache types like migraines. Patients typically describe the pain as mild to moderate and can persist for varying durations, but the key features remain the bilateral nature and the sensation of tightness rather than sharp or throbbing pain. Presenting with associated symptoms like nausea or being unilateral would suggest other headache types, such as migraines or cluster headaches. Understanding these characteristics is essential for differentiating tension-type headache from other types of headaches for accurate diagnosis and management.

6. Which drug is likely to be used in the Tensilon test for diagnosing myasthenia gravis?

A. Edrophonium

B. Aminoglycosides

C. Corticosteroids

D. Neostigmine

The Tensilon test is a diagnostic procedure specifically used to assess the presence of myasthenia gravis, an autoimmune disorder characterized by weakness and fatigue of skeletal muscles. The test involves the administration of a drug that temporarily increases the availability of acetylcholine at the neuromuscular junction, thereby improving muscle strength. Edrophonium, a short-acting acetylcholinesterase inhibitor, is the drug utilized in the Tensilon test. By inhibiting the enzyme acetylcholinesterase, edrophonium prevents the breakdown of acetylcholine, leading to an increase in its concentration available to bind to receptors on the muscle side of the neuromuscular junction. This results in a rapid increase in muscle strength, observable shortly after administration, and can help confirm a diagnosis of myasthenia gravis. Other options, such as aminoglycosides and corticosteroids, are not suitable for the Tensilon test. Aminoglycosides can actually exacerbate muscle weakness in myasthenia gravis due to their effects on the neuromuscular junction. Corticosteroids are used for long-term management of myasthenia gravis but do not serve as a diagnostic tool in this context. Neostigmine is another

SAMPLE

7. Which area of the brain is most likely affected in a patient with dysarthria, tongue deviation to the right, and right arm weakness?

A. Right brain stem

B. Left cerebral cortex

C. Left brain stem

D. Right cerebellum

A patient presenting with dysarthria, tongue deviation to the right, and right arm weakness suggests a specific pattern of neurological impairment. In this case, the combination of dysarthria and unilateral weakness can indicate that the area of the brainstem responsible for controlling motor functions has been affected. Dysarthria, which involves difficulty in articulating speech due to neurological causes, often arises from damage to areas that control speech muscles. The deviation of the tongue to the right indicates that the muscles of the left side of the tongue are impaired, which suggests weakness or paralysis on that side. This could result from damage to the lower motor neurons that innervate the tongue, likely located in the brainstem. The right arm weakness indicates that there may be an upper motor neuron lesion affecting the right side of the body. If the left side of the brain controls the right side of the body, an injury to the left brainstem could cause motor deficits on the right. However, in this scenario, the question suggests the right arm weakness along with the specific dysarthria and left tongue deviation, aligning more closely with an issue in the right brain stem. This is because both cranial nerves and motor pathways controlling the tongue and arm cross over, meaning that damage

8. What is the first-line treatment for a patient diagnosed with complex regional pain syndrome?

A. Corticosteroids

B. Physical therapy

C. Naprosyn (NSAIDs)

D. Gabapentin

For the management of complex regional pain syndrome (CRPS), the primary approach typically involves physical therapy. This treatment is essential because it helps to improve mobility, reduces pain, and aids in the recovery of function in the affected limb. Physical therapy focuses on desensitization, increasing range of motion, and enhancing strength, which are all crucial for patients with CRPS to regain normal function. While corticosteroids, NSAIDs such as Naprosyn, and medications like Gabapentin may play supportive roles in managing inflammation and neuropathic pain, they do not address the underlying issues of movement and function the way physical therapy does. In fact, early intervention with physical therapy can prevent long-term complications and improve outcomes in patients suffering from CRPS, making it the cornerstone of first-line treatment.

SAMPLE

9. What cerebrospinal fluid (CSF) finding is most commonly associated with Guillain-Barré syndrome?

A. Decreased CSF glucose

B. Elevated CSF protein content

C. Normal white blood cell count

D. Presence of bacterial infection markers

Elevated cerebrospinal fluid (CSF) protein content is a hallmark finding in Guillain-Barré syndrome, particularly during the course of the disease. This increase in protein, often referred to as albuminocytologic dissociation, occurs without a corresponding increase in white blood cell count, which retains a normal level in the early stages of the syndrome. This discrepancy is characteristic of Guillain-Barré syndrome and reflects the underlying pathology where nerve damage leads to a breakdown of the blood-nerve barrier, allowing proteins to leak into the CSF while the inflammatory response may not yet be detectable through increased white blood cells. In Guillain-Barré syndrome, the clinical presentation is often preceded by an infection, frequently of a viral or bacterial origin, but the specific CSF findings are crucial in distinguishing it from other neuropathies. The elevation in protein levels provides important diagnostic information, thereby supporting the diagnosis when combined with the typical clinical manifestations, such as rapid onset of muscle weakness and sensory changes.

10. In the case of suspected viral encephalitis, what is the first imaging study recommended before lumbar puncture?

A. MRI of the brain

B. CT scan of the brain

C. PET scan

D. X-ray of the skull

In cases of suspected viral encephalitis, a CT scan of the brain is typically recommended before performing a lumbar puncture. This is primarily due to the potential for increased intracranial pressure or other intracranial pathologies that could lead to complications during the lumbar puncture procedure. A CT scan can help identify any mass lesions, such as abscesses or tumors, and can also rule out conditions like hemorrhagic strokes that may require immediate intervention. The CT scan provides quick and efficient imaging, which is crucial in an emergency setting where time is of the essence. While an MRI of the brain offers more detailed imaging and is indeed valuable in diagnosing encephalitis, it is not the first imaging study typically performed in acute settings. PET scans are more specialized and generally not used initially in this context, while a skull X-ray would not provide adequate information to assess for the complications related to suspected encephalitis. Therefore, the CT scan serves as a safe and effective first step in the evaluation process.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://exammasterneuro.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE