FETAL HEALTH SURVEILLANCE PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://fetalhealthsurvaillance.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the normal pH range for venous blood in term newborns?

A. 7.25-7.35

B. 7.28-7.40

C. 7.40-7.50

D. 7.10-7.30

2. What happens to fetal cardiac output when heart rate decreases due to increased blood pressure?

A. It increases

B. It decreases

C. It remains the same

D. It fluctuates significantly

3. What defines recurrent decelerations during fetal monitoring?

A. Decelerations that occur with less than 25% of contractions

B. Decelerations that occur with 50% or fewer contractions in a 20 minute period

C. Decelerations that occur with 50% or more contractions in a 20 minute period

D. Decelerations that are intermittent and non-repetitive

4. What laboratory values indicate mixed acidosis?

A. Normal pH, low HCO_3

B. Low pH, high pCO_2

C. High HCO_3, normal pCO_2

D. Normal pH, low base excess

5. Which of the following is a primary factor that regulates fetal heart rate?

A. Blood pressure

B. Medulla oblongata activity

C. Amniotic fluid composition

D. Cardiac contractility

SAMPLE

6. What is a common limitation in obtaining adequate volume for fetal scalp blood sampling?

A. Difficulty in accessing the fetal heart rate

B. Insufficient operator training

C. Difficulty in obtaining the necessary volume

D. Response time delay

7. Which of the following is not a criterion to define an acute intrapartum event causing cerebral palsy?

A. Evidence of metabolic acidosis in umbilical cord blood

B. Cerebral palsy of the spastic quadriplegic type

C. Presence of infection during labor

D. Early onset of severe neonatal encephalopathy

8. How long must a change in baseline FHR be maintained to be considered the new baseline?

A. 5 minutes

B. 10 minutes

C. 15 minutes

D. 20 minutes

9. What is the health range for base excess/deficit?

A. -5.5 to 0.1 (meq/L)

B. 0.2 to 5.0 (meq/L)

C. -10 to -5 (meq/L)

D. 0.0 to 0.5 (meq/L)

10. What is a reassuring indicator regarding fetal health monitoring?

A. Spontaneous accelerations

B. Late decelerations

C. Prolonged variable decelerations

D. Minimal baseline variability

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. B
5. B
6. C
7. C
8. B
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. What is the normal pH range for venous blood in term newborns?

A. 7.25-7.35

B. 7.28-7.40

C. 7.40-7.50

D. 7.10-7.30

The normal pH range for venous blood in term newborns is indeed between 7.28 and 7.40. This range is important for maintaining homeostasis, and it reflects the internal environment of the body, ensuring optimal conditions for cellular functions. In term newborns, venous blood pH can vary slightly due to factors like transition from fetal to neonatal circulation, respiratory adjustments post-birth, and metabolic activity. The selected range allows for some variability while still being within the parameters that indicate a normally functioning biological system. A pH below this range could indicate acidosis, while a pH above might suggest alkalosis, both of which could prompt further investigation or intervention in a clinical setting. Understanding these normal values aids healthcare professionals in recognizing potential complications and managing the neonate's health effectively.

2. What happens to fetal cardiac output when heart rate decreases due to increased blood pressure?

A. It increases

B. It decreases

C. It remains the same

D. It fluctuates significantly

When there is an increase in blood pressure, the body initiates various physiological responses to maintain homeostasis and ensure adequate blood flow and oxygenation to vital organs, including the fetus. A decrease in fetal heart rate, in the context of increased blood pressure, typically indicates that the body may be experiencing a compensatory mechanism where cardiac output is reduced. Cardiac output, which is the product of heart rate and stroke volume, can be affected by these changes. A decrease in heart rate generally leads to a reduction in cardiac output if stroke volume does not significantly increase to compensate. In this situation, because the heart is beating less frequently, there is a lower volume of blood being pumped out over time. This relationship is particularly crucial in fetal monitoring and health assessments, where maintaining a balanced oxygen delivery is essential for fetal well-being. Thus, in scenarios where blood pressure rises and causes a decrease in heart rate, the overall result would be a decrease in fetal cardiac output due to the reduction in the frequency of heart contractions, aligning with the answer choice.

3. What defines recurrent decelerations during fetal monitoring?

A. Decelerations that occur with less than 25% of contractions

B. Decelerations that occur with 50% or fewer contractions in a 20 minute period

C. Decelerations that occur with 50% or more contractions in a 20 minute period

D. Decelerations that are intermittent and non-repetitive

Recurrent decelerations during fetal monitoring are defined by their consistent occurrence in relation to contractions. Specifically, these decelerations appear with 50% or more of the contractions observed within a designated time frame, such as 20 minutes. This pattern is crucial as it may indicate a potential issue with uteroplacental perfusion, which can affect fetal oxygenation and overall well-being. The assessment of decelerations is important in monitoring fetal health, as it helps healthcare providers identify potential distress or compromise. When decelerations are recurrent, it raises the concern for possible fetal hypoxia and justifies further evaluation and intervention to ensure the well-being of the baby. Recognizing these patterns is integral for appropriate clinical decision-making during labor.

SAMPLE

4. What laboratory values indicate mixed acidosis?

A. Normal pH, low HCO3

B. Low pH, high pCO2

C. High HCO3, normal pCO2

D. Normal pH, low base excess

Mixed acidosis occurs when there are simultaneous metabolic and respiratory components affecting the acid-base balance in the body. In this context, the identification of a low pH is crucial, as it indicates acidemia. High pCO2 suggests a respiratory contributor to the acidemia, often resulting from conditions like respiratory failure, while low HCO3 points to metabolic acidosis, which can arise from a variety of metabolic disturbances. When evaluating these lab values, a low pH paired with a high pCO2 confirms that there is both a respiratory and a metabolic acidosis present. The high pCO2 reflects insufficient ventilation to eliminate CO2, leading to a retention of carbonic acid, while the low HCO3 indicates that the buffering capacity of the body is diminished due to metabolic disturbances. Therefore, the combination of low pH and high pCO2 serves as a hallmark of mixed acidosis. In contrast, the other values do not appropriately reflect mixed acidosis. Normal pH with low HCO3 indicates a compensated metabolic acidosis without significant respiratory issue. High HCO3 with normal pCO2 typically indicates metabolic alkalosis or compensation for respiratory acidosis rather than mixed acidosis. Normal pH with low base excess may suggest a metabolic disturbance

5. Which of the following is a primary factor that regulates fetal heart rate?

A. Blood pressure

B. Medulla oblongata activity

C. Amniotic fluid composition

D. Cardiac contractility

The regulation of fetal heart rate is primarily managed by the autonomic nervous system, which is influenced by the activity of the medulla oblongata. This part of the brainstem has vital roles in controlling vital involuntary functions, including heart rate. It receives input from various sensory pathways and helps integrate these signals to maintain homeostasis, particularly during fetal development. The medulla oblongata sends signals to the heart through the autonomic nervous system, adjusting the heart rate according to the metabolic needs of the fetus. A healthy interaction between sympathetic and parasympathetic activity, which originates in the medulla oblongata, is crucial for appropriate fetal heart rate responses to various stimuli, such as maternal activity and environmental changes. While blood pressure, amniotic fluid composition, and cardiac contractility may play roles in fetal health and outcomes, they do not primarily dictate the regulation of the fetal heart rate as do the mechanisms driven by the medulla oblongata.

SAMPLE

6. What is a common limitation in obtaining adequate volume for fetal scalp blood sampling?

A. Difficulty in accessing the fetal heart rate

B. Insufficient operator training

C. Difficulty in obtaining the necessary volume

D. Response time delay

A common limitation in obtaining adequate volume for fetal scalp blood sampling lies in the difficulty of obtaining the necessary volume of blood from the fetus. This procedure, which is typically performed to assess fetal well-being by measuring the pH and other blood gases, requires a sufficient volume to produce accurate and reliable results. Multiple factors contribute to this challenge. The anatomy of the fetal scalp can sometimes make it difficult to retrieve an adequate sample, particularly if the fetus is in an unfavorable position or if the sampling device does not function effectively. Additionally, maternal and fetal factors, such as the presence of oligohydramnios (low amniotic fluid) or fetal compromise, can further complicate the sampling process, limiting the volume that can be obtained. This necessity for an adequate volume underscores the importance of careful technique and sometimes may lead to the decision to forego the procedure altogether if the operator feels that the chances of obtaining a usable sample are low, thereby ensuring the safety and health of the fetus.

7. Which of the following is not a criterion to define an acute intrapartum event causing cerebral palsy?

A. Evidence of metabolic acidosis in umbilical cord blood

B. Cerebral palsy of the spastic quadriplegic type

C. Presence of infection during labor

D. Early onset of severe neonatal encephalopathy

The correct choice highlights that the presence of infection during labor is not considered a defining criterion for an acute intrapartum event that causes cerebral palsy. In defining the criteria for acute intrapartum events leading to cerebral palsy, the focus is on immediate and significant physiological changes that occur during labor and delivery, such as fetal distress as evidenced by metabolic disturbances in umbilical cord blood, the onset and severity of neonatal encephalopathy, and the specific clinical presentation of cerebral palsy types. Metabolic acidosis in cord blood often signifies inadequate oxygenation or perfusion during labor, which is closely related to acute intrapartum events. Similarly, early onset of severe neonatal encephalopathy indicates an immediate and serious injury to the brain that could result from hypoxic-ischemic insults occurring during labor. The classification of cerebral palsy also considers types, with spastic quadriplegia being one of the common forms resulting from acute brain injury. Therefore, both the presence of metabolic acidosis and the specific types of cerebral palsy are deemed critical indicators related to acute events during labor. In contrast, while infections during labor can negatively impact fetal health, they are not classified as direct acute events related specifically to the immediate causation of cerebral pals

SAMPLE

8. How long must a change in baseline FHR be maintained to be considered the new baseline?

A. 5 minutes

B. 10 minutes

C. 15 minutes

D. 20 minutes

A change in baseline fetal heart rate (FHR) must be maintained for at least 10 minutes to be considered a new baseline. This duration allows for a sufficient assessment period to determine whether the FHR has truly shifted from its previous baseline rather than being a temporary fluctuation due to short-term factors, such as fetal movements or contractions. In clinical practice, monitoring the baseline FHR is essential for assessing fetal well-being. If the heart rate remains altered for a full 10 minutes following the initial change, it indicates a more persistent shift in the fetal condition, warranting further observation or intervention as necessary. This standard is established to ensure accuracy in interpreting fetal heart rate changes and making informed clinical decisions.

9. What is the health range for base excess/deficit?

A. -5.5 to 0.1 (meq/L)

B. 0.2 to 5.0 (meq/L)

C. -10 to -5 (meq/L)

D. 0.0 to 0.5 (meq/L)

The correct health range for base excess/deficit is from -5.5 to 0.1 milliequivalents per liter (meq/L). This range is important in assessing a patient's acid-base status. Base excess refers to the amount of excess or insufficient base in the bloodstream and is used to evaluate metabolic acidosis or alkalosis. A base excess within this range indicates a relatively normal metabolic state, where values towards the lower end (-5.5) may indicate a slight metabolic acidosis and values closer to 0.1 indicate a more balanced metabolic condition. Values that fall outside of this range can signal significant metabolic disturbances. For instance, a base excess lower than -5.5 may indicate severe acidosis, which could require immediate clinical intervention. Understanding this range is crucial for healthcare providers in monitoring the metabolic state of their patients and making appropriate medical decisions.

10. What is a reassuring indicator regarding fetal health monitoring?

A. Spontaneous accelerations

B. Late decelerations

C. Prolonged variable decelerations

D. Minimal baseline variability

Spontaneous accelerations are a reassuring indicator in fetal health monitoring because they demonstrate a well-functioning autonomic nervous system and indicate that the fetus is responding appropriately to the environment, which includes changes in activity and oxygen levels. These accelerations are temporary increases in the fetal heart rate that are often seen when the fetus is active or experiencing movement, reflecting good oxygenation and overall well-being. The presence of spontaneous accelerations suggests that the fetal heart rate is showing variability and responsiveness, both of which are positive signs during monitoring. In contrast, late decelerations and prolonged variable decelerations signify potential distress or concerns regarding fetal oxygen levels and overall health. Late decelerations typically indicate issues such as uteroplacental insufficiency, while prolonged variable decelerations can suggest umbilical cord compression, both of which may require further medical evaluation and intervention. Minimal baseline variability, on the other hand, suggests diminished responsiveness of the fetal heart rate, which can be a sign of fetal hypoxia or other underlying problems. Therefore, monitoring for spontaneous accelerations is a vital part of fetal health assessments as they provide reassurance regarding the fetus's condition.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://fetalhealthsurvaillance.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE