

FGDP SELECTION CRITERIA FOR DENTAL RADIOGRAPHY PRACTICE TEST (SAMPLE) STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://fgdpselectioncriteriafordentalrad.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. Why is the practice of using 'routine' bitewings for caries detection in children not supported when based solely on time since the last examination?**
 - A. Because radiographic equipment degrades over time
 - B. Because individuals can move in and out of caries risk categories over time
 - C. Because age is the only factor
 - D. Because bitewings harm the patient
- 2. Which of the following is an example of a deterministic radiation effect?**
 - A. Cancer
 - B. Hair loss
 - C. Genetic mutation
 - D. Cataracts
- 3. What additional information may you want to see on a pre-diagnostic radiograph for surgical root canal treatment?**
 - A. Final length of root and periodontal support
 - B. Evidence of caries
 - C. Presence of occlusal wear
 - D. Patient age
- 4. What does limited-volume CBCT refer to in implant imaging?**
 - A. A CBCT scan with a restricted field of view focusing on area of interest
 - B. A CBCT scan with full skull coverage
 - C. A 2D radiograph only
 - D. A CT scan with full head coverage
- 5. Which statement correctly justifies radiographic imaging in a patient?**
 - A. Imaging should be justified by clinical benefit and risk
 - B. Image whenever possible
 - C. Not necessary
 - D. Use imaging to maximize revenue

6. In dental radiography safety, in DDH, how far should a person stand from all machines?
- A. 1.5 m
 - B. 2.0 m
 - C. 2.5 m
 - D. 3.0 m
7. For identifying an unerupted maxillary canine, which radiographic views are appropriate?
- A. DPT & AOM vertical parallax
 - B. Panoramic radiograph
 - C. 2 periapicals with horizontal parallax
 - D. Lateral cephalogram
8. What are common artifacts on dental radiographs, and how can they be prevented or corrected?
- A. Motion blur, processing artifacts, lead foil artifact, sensor blemishes, anatomy misinterpretations
 - B. Only motion blur
 - C. Color artifacts
 - D. Dust on the sensor
9. Are there dose limits for patients in medical radiography?
- A. Yes, 1 mSv per year
 - B. No, there are no dose limits for patients
 - C. Yes, 0.1 mSv per year
 - D. Yes, 20 mSv per year
10. What are the functions of post-operative radiographs following root canal treatment?
- A. Assessment of quality of root canal filling
 - B. Baseline for review
 - C. Indication of possible complications or errors
 - D. All of the above

Answers

SAMPLE

1. B
2. B
3. A
4. A
5. A
6. B
7. A
8. A
9. B
10. D

SAMPLE

Explanations

SAMPLE

1. Why is the practice of using 'routine' bitewings for caries detection in children not supported when based solely on time since the last examination?

- A. Because radiographic equipment degrades over time
- B. Because individuals can move in and out of caries risk categories over time**
- C. Because age is the only factor
- D. Because bitewings harm the patient

The key idea is that decisions about taking bitewing radiographs for caries detection should be guided by a child's current caries risk and clinical findings, not by a fixed time since the last exam. Caries risk is dynamic: a child's likelihood of developing or progressing lesions can change as factors like diet, fluoride exposure, oral hygiene, saliva flow, eruption status, and prior caries experience fluctuate. Because of this, the probability of finding new or advancing interproximal caries changes over time, so a rigid interval may miss new problems in higher-risk children or lead to unnecessary imaging in lower-risk ones. That's why the best answer is that individuals can move in and out of caries risk categories over time. This dynamic risk means radiographic decisions should be based on the present risk assessment and any clinical signs, rather than simply on how much time has passed since the last radiograph. The other statements don't fit as the primary reason. Equipment wear is not the driving factor for whether to use routine bitewings. Age alone isn't the sole determinant of how often imaging is needed. And while minimizing radiation is important, the claim that bitewings inherently harm the patient ignores that imaging is typically justified by current risk and benefit, not blanket prohibition.

2. Which of the following is an example of a deterministic radiation effect?

- A. Cancer
- B. Hair loss**
- C. Genetic mutation
- D. Cataracts

Deterministic effects occur only after a threshold dose and become more severe as the dose increases. Hair loss fits this pattern: once the exposure passes a certain level, hair follicles are damaged and shedding occurs, with greater severity at higher doses. In contrast, cancer and genetic mutations are stochastic—they can occur at any dose and the probability rises with dose but without a clear threshold. Cataracts are also deterministic, but hair loss is the classic, straightforward example used to illustrate dose-dependent, threshold-based tissue reactions in radiobiology.

3. What additional information may you want to see on a pre-diagnostic radiograph for surgical root canal treatment?

- A. Final length of root and periodontal support**
- B. Evidence of caries
- C. Presence of occlusal wear
- D. Patient age

For planning surgical root canal treatment, you need radiographic information that directly guides the procedure. The most important additional detail is the final length of the root and the status of the surrounding periodontal support. Knowing the root length helps set the correct working length for instrumentation and informs whether any apical access or root-end preparation is needed during surgery. Assessing periodontal support on the image reveals the health of the periradicular bone around the apex and the likely prognosis, indicating how feasible the surgical approach will be and whether healing can be expected. Evidence of caries can influence restoration after endodontic therapy, and occlusal wear or patient age may affect overall treatment planning, but they do not provide the critical planning data for the surgical procedure itself.

4. What does limited-volume CBCT refer to in implant imaging?

- A. A CBCT scan with a restricted field of view focusing on area of interest**
- B. A CBCT scan with full skull coverage
- C. A 2D radiograph only
- D. A CT scan with full head coverage

Limited-volume CBCT is a cone-beam CT scan that confines imaging to a small, defined region around the planned implant site. This focused field of view provides a high-resolution, three-dimensional view of the local anatomy, allowing precise assessment of bone height, width, and quality and the relationship to critical structures such as the inferior alveolar canal or maxillary sinus. Because you image only the area of interest, the radiation dose is lower and the scan time shorter, while still giving detailed information necessary for accurate implant planning, including position, angulation, and depth. The other scenarios describe larger fields of view or different imaging modalities, which would expose more tissue or fail to provide the 3D detail essential for implant planning.

5. Which statement correctly justifies radiographic imaging in a patient?

- A. Imaging should be justified by clinical benefit and risk**
- B. Image whenever possible
- C. Not necessary
- D. Use imaging to maximize revenue

The main idea is that radiographic imaging should be justified by a clear clinical benefit relative to the radiation risk. In dental practice, this means each imaging exam is chosen because the information it will provide is likely to influence diagnosis, treatment planning, or prognosis. You assess the patient's symptoms, history, clinical exam findings, and whether the imaging answer will change how you manage care. If the expected diagnostic gain outweighs the radiation exposure, imaging is warranted. This also ties to keeping exposure as low as reasonably achievable (ALARA) by choosing the appropriate modality, limiting the field of view, and using the minimum number of images needed. This is why imaging should not be done "whenever possible" or as a routine without a clinical question, and why it is not appropriate to image to increase revenue. Instead, imaging should be used only when it will meaningfully impact patient care.

6. In dental radiography safety, in DDH, how far should a person stand from all machines?

- A. 1.5 m
- B. 2.0 m**
- C. 2.5 m
- D. 3.0 m

Distance from the X-ray source determines how much radiation you receive—the farther you are, the lower the dose due to the inverse square relationship. For dental radiography safety, the practical minimum stand-off distance from the machine is two metres. This provides a meaningful reduction in scattered radiation while keeping the workflow in a typical clinic workable. If you stand closer, exposure rises above recommended levels; standing farther than two metres offers extra protection but is not necessary as a minimum and may be impractical in small rooms.

7. For identifying an unerupted maxillary canine, which radiographic views are appropriate?

- A. DPT & AOM vertical parallax**
- B. Panoramic radiograph
- C. 2 periapicals with horizontal parallax
- D. Lateral cephalogram

Gaining a precise location of an unerupted maxillary canine relies on using two radiographs taken from different angles so you can apply parallax and judge depth. A panoramic (DPT) gives the overall layout of the dentition, but because it's a single projection, structures overlap and depth isn't revealed. When you add a vertical parallax view (the AOM vertical parallax), you change the angle in the vertical plane and observe how the tooth image shifts relative to adjacent teeth. The way it moves tells you whether the tooth is buccal or palatal and how deep it sits, which is essential for planning treatment. If you relied on a panoramic view alone, you wouldn't get the depth information needed. Two periapicals with horizontal parallax can provide some buccolingual clues, but without a vertical parallax component the localization isn't as reliable for an upper canine. A lateral cephalogram isn't suitable for localizing an unerupted tooth because it's not designed for three-dimensional localization of a single tooth and offers limited parallax information.

8. What are common artifacts on dental radiographs, and how can they be prevented or corrected?

- A. Motion blur, processing artifacts, lead foil artifact, sensor blemishes, anatomy misinterpretations**
- B. Only motion blur
- C. Color artifacts
- D. Dust on the sensor

The main idea being tested is understanding what kinds of features on dental radiographs are considered artifacts and how to prevent or correct them. Motion blur shows up when the patient or equipment moves during exposure, producing a smeared image that can obscure details. It's best tackled with clear patient instructions, proper stabilization, and, when possible, shorter exposure times or faster imaging techniques so the patient can stay still. Processing artifacts come from problems in the film development or processing steps, such as uneven development, incorrect timing, temperature, or chemical contamination. These produce streaks, shading, or foggy areas that don't reflect the patient's anatomy. Prevention involves regular maintenance of the processor, using fresh chemicals, keeping temperatures and timers correct, and handling films carefully to avoid processing errors. Lead foil artifact arises from the lead foil backing used in traditional film packets. If the foil is damaged, creased, or placed incorrectly, it can cast shadows or create a recognizable pattern on the radiograph. Prevention is straightforward: ensure the foil backing is intact, properly oriented, and not folded or bruised during packet preparation and handling. Sensor blemishes are issues in digital radiography from dead pixels, scratches, or surface irregularities on the sensor. They appear as fixed spots or irregularities on images. Prevention focuses on gentle handling of the sensor, keeping it clean, and performing regular quality checks; software tools and calibration can help correct or flag these areas when they occur. Anatomy misinterpretations aren't true image "mistakes" in the sense of a device problem, but they are artifacts created by how the anatomy is projected or overlapped in a given view. Misinterpretations can happen if projection geometry is off or if the view doesn't clearly separate structures. Prevention comes from using proper angulation and positioning, obtaining appropriate additional views when needed, and developing a solid understanding of normal anatomical relationships so features aren't mistaken for pathology or artifacts. So the best answer incorporates all the common sources—movement, processing, lead foil issues, sensor blemishes, and anatomical misinterpretations—together with practical prevention and correction strategies.

9. Are there dose limits for patients in medical radiography?

- A. Yes, 1 mSv per year
- B. No, there are no dose limits for patients**
- C. Yes, 0.1 mSv per year
- D. Yes, 20 mSv per year

In medical radiography there are no fixed dose limits for patients. Each imaging exam should be justified by clinical need, and the technique should be optimized so the patient receives the minimum dose that still yields a clinically useful image (the ALARA principle). Dose Reference Levels (DRLs) exist as guidance to help identify unusually high doses and prompt optimization, but they are not hard limits. By contrast, dose limits do apply to workers and the public, not to patient exposures.

10. What are the functions of post-operative radiographs following root canal treatment?

- A. Assessment of quality of root canal filling
- B. Baseline for review
- C. Indication of possible complications or errors
- D. All of the above**

Post-operative radiographs after root canal treatment serve multiple important roles. They let you check the quality of the obturation—whether the filling reaches the correct length, fills densely without voids, and avoids over- or under-extension beyond the apex. They also establish a baseline image that you can compare with future radiographs to monitor healing and detect any changes in the periapical area. Additionally, the image can reveal iatrogenic issues or errors from the procedure, such as instrument separation, perforations, ledging, missed canals, or other anomalies that could affect prognosis. Because these three functions together address the outcome, the post-operative radiograph is considered to serve all these purposes. Remember, while valuable, the radiograph is a two-dimensional view, so some details may require supplemental imaging if needed.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://fgdpselectioncriteriafordentalrad.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE