FISDAP MEDICAL PARAMEDIC PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://fisdapmedicalparamedic.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is a common sign of hypoxia?

A. Cyanosis of the skin
B. Increased appetite
C. Elevated blood pressure
D. Severe headache

2. When should nitroglycerin be administered to a patient with chest pain?

A. Only if the patient has a previous prescription
B. When the patient is experiencing shortness of breath
C. After checking the patient's blood pressure
D. Only in the presence of cardiac arrest

3. What is a common side effect of opioid overdose?

A. Increased alertness and energy
B. Respiratory depression
C. Severe headaches and nausea
D. High blood pressure

4. What symptom is most indicative of a rupture of varices in a patient with liver disease?

A. Severe dysphagia
B. Abdominal pain
C. Vomiting bright red blood
D. Elevated blood pressure

5. Which medication is commonly administered for status epilepticus?

A. Diazepam
B. Midazolam
C. Phenobarbital
D. Lorazepam

6. Which symptom is common in a patient with pneumonia?

A. Colicky abdominal pain
B. Enlarged lymph nodes
C. Difficulty breathing
D. Joint pain

SAMPLE

7. When assessing a trauma patient, what is the priority intervention?

A. Perform a head-to-toe assessment

B. Establish and maintain an airway

C. Control any bleeding

D. Take vital signs

8. Which algorithm should be used for treating a cocaine overdose?

A. Cardiac Arrest Algorithm

B. Acute Coronary Algorithm

C. Stroke Algorithm

D. Pediatric Emergency Algorithm

9. What is one of the most dangerous complications of an untreated drug overdose?

A. Kidney failure

B. Cardiac arrest

C. Stomach bleeding

D. Seizures

10. Which vital sign is least likely affected by shock?

A. Heart rate

B. Blood pressure

C. Respiratory rate

D. Temperature

SAMPLE

Answers

SAMPLE

1. A
2. C
3. B
4. C
5. B
6. C
7. B
8. B
9. B
10. D

SAMPLE

Explanations

SAMPLE

1. What is a common sign of hypoxia?

A. Cyanosis of the skin

B. Increased appetite

C. Elevated blood pressure

D. Severe headache

Cyanosis of the skin is a common sign of hypoxia because it indicates that there is inadequate oxygenation of the blood. When the body's tissues do not receive enough oxygen, hemoglobin, which normally carries oxygen in the blood, can become less saturated. This lack of oxygen can cause the blood to take on a bluish tint, particularly noticeable in areas with high capillary networks like the lips, fingertips, and nail beds. This visible change in skin color is a physical manifestation of hypoxia and serves as a critical indicator for medical professionals when assessing a patient's oxygenation status. In contrast, other options such as increased appetite, elevated blood pressure, and a severe headache are not directly associated with hypoxia. Increased appetite typically does not correlate with low oxygen levels, and while elevated blood pressure may occur due to various physiological stressors, it is not an exclusive sign of hypoxia. A severe headache could be associated with other conditions, including migraines or tension headaches, and is not a direct indicator of hypoxia.

2. When should nitroglycerin be administered to a patient with chest pain?

A. Only if the patient has a previous prescription

B. When the patient is experiencing shortness of breath

C. After checking the patient's blood pressure

D. Only in the presence of cardiac arrest

Administering nitroglycerin to a patient with chest pain should always involve an assessment of the patient's blood pressure prior to giving the medication. This is crucial because nitroglycerin acts as a vasodilator, which can significantly lower blood pressure. If the patient's blood pressure is too low, administering nitroglycerin could lead to severe hypotension, possibly resulting in adverse cardiovascular events. Monitoring the patient's blood pressure ensures that the paramedic can confirm that the medication is safe to give. Generally, nitroglycerin is indicated for patients experiencing angina or presumed cardiac-related chest pain, as long as their blood pressure is within a safe range. This precaution helps ensure effective and safe treatment for the patient presenting with chest pain. Considering the context of other options, administering nitroglycerin solely based on a previous prescription or in the presence of shortness of breath or cardiac arrest does not cover the safety assessment required in emergency situations. A thorough evaluation, including blood pressure measurement, remains the priority in determining the appropriateness of nitroglycerin administration.

SAMPLE

3. What is a common side effect of opioid overdose?

A. Increased alertness and energy

B. Respiratory depression

C. Severe headaches and nausea

D. High blood pressure

Respiratory depression is a hallmark effect of opioid overdose due to the way opioids interact with the central nervous system (CNS). Opioids bind to specific receptors in the brain and spinal cord that not only help modulate pain but also regulate automatic functions such as breathing. In an overdose situation, these respiratory centers can become significantly suppressed, leading to shallow or slowed breathing, and in severe cases, respiratory arrest. This is a critical concern for emergency responders and healthcare providers, as impaired breathing can quickly result in hypoxia—an insufficient oxygen supply to the body's tissues, which can lead to further complications, including brain damage or death if not promptly addressed. Recognizing respiratory depression as a common indicator of opioid overdose is essential for making timely and effective treatment decisions, such as administering naloxone, an opioid antagonist that can reverse the effects of opioid overdose and restore normal respiratory function.

4. What symptom is most indicative of a rupture of varices in a patient with liver disease?

A. Severe dysphagia

B. Abdominal pain

C. Vomiting bright red blood

D. Elevated blood pressure

Vomiting bright red blood is the most indicative symptom of a rupture of varices in a patient with liver disease. This occurs due to increased pressure in the portal vein, a condition known as portal hypertension, which can lead to the swelling and ultimate rupture of varices, typically in the esophagus or stomach. When these varices rupture, they can bleed profusely, and the blood is typically fresh due to its proximity to the point of rupture in the upper gastrointestinal tract. This results in the characteristic symptom of hematemesis, or vomiting of bright red blood, which is a critical and urgent sign that requires immediate medical attention. Other symptoms, while they may be present in patients with liver disease or gastrointestinal bleeding, do not specifically indicate a rupture of varices. For example, severe dysphagia (difficulty swallowing) and abdominal pain are not directly related to variceal rupture and can be caused by a variety of other conditions. Elevated blood pressure is not typically associated with variceal bleeding but may instead reflect the underlying liver disease and portal hypertension without directly indicating an active bleed. Thus, the symptom of vomiting bright red blood is crucial in identifying a rupture of varices and guiding emergency treatment.

SAMPLE

5. Which medication is commonly administered for status epilepticus?

A. Diazepam

B. Midazolam

C. Phenobarbital

D. Lorazepam

Status epilepticus is a medical emergency characterized by prolonged seizures or a series of seizures without full recovery in between. The choice of medication in these situations is critical for seizure control and the prevention of long-term neurological damage. Midazolam is commonly chosen due to its rapid onset of action when administered intravenously, intramuscularly, or intranasally. It has a short half-life, allowing for quick titration and adjustment based on the patient's response. Its effectiveness in terminating seizures promptly makes it a preferred option in the acute management of status epilepticus, particularly in prehospital and emergency settings. While other medications like diazepam, phenobarbital, and lorazepam can also be used for managing seizures, midazolam's favorable pharmacokinetic properties make it especially effective in urgent scenarios where swift action is needed. Its versatility in routes of administration adds to its practicality, making it an optimal choice for clinicians faced with patients experiencing status epilepticus.

6. Which symptom is common in a patient with pneumonia?

A. Colicky abdominal pain

B. Enlarged lymph nodes

C. Difficulty breathing

D. Joint pain

Difficulty breathing is a common symptom in patients with pneumonia due to inflammation and infection in the lung tissues, which can lead to impaired gas exchange. The inflammation causes the lung air sacs (alveoli) to fill with fluid or pus, resulting in reduced oxygen saturation and making it harder for the patient to breathe effectively. This symptom may present as shortness of breath, increased respiratory rate, or use of accessory muscles during breathing. In the context of pneumonia, other symptoms may accompany difficulty breathing, such as cough, fever, and chest pain, but the fundamental issue lies with the respiratory system's ability to function properly. Knowing this can help differentiate pneumonia from other conditions that may manifest different symptom profiles, such as gastrointestinal issues or joint disorders.

7. When assessing a trauma patient, what is the priority intervention?

A. Perform a head-to-toe assessment

B. Establish and maintain an airway

C. Control any bleeding

D. Take vital signs

In trauma cases, establishing and maintaining an airway is considered the priority intervention due to the critical importance of oxygenation in preventing morbidity and mortality. A compromised airway can lead to insufficient oxygen delivery to vital organs, which can quickly become life-threatening. This is particularly vital in trauma scenarios where patients may have sustained injuries that compromise their ability to breathe effectively or where they may be at risk of airway obstruction due to blood, secretions, or body positioning. While head-to-toe assessments, controlling bleeding, and taking vital signs are all essential components of trauma care, they come after ensuring the patient has a patent airway. Without a secured airway, subsequent interventions may not be effective, and the patient's condition could rapidly deteriorate. Therefore, the airway is prioritized to ensure that the patient can be properly oxygenated and to stabilize their condition as the other assessments and treatments are carried out.

SAMPLE

8. Which algorithm should be used for treating a cocaine overdose?

A. Cardiac Arrest Algorithm

B. Acute Coronary Algorithm

C. Stroke Algorithm

D. Pediatric Emergency Algorithm

For treating a cocaine overdose, the Acute Coronary Algorithm is the appropriate choice due to the cardiovascular effects that cocaine can have on the body. Cocaine use can lead to a range of serious cardiovascular issues, such as increased heart rate, hypertension, and the potential for acute coronary syndrome (ACS), which includes conditions like myocardial infarction. The Acute Coronary Algorithm provides specific interventions aimed at managing these cardiac complications. This includes monitoring the patient's vital signs, administering oxygen if necessary, providing aspirin to inhibit further clot formation, and preparing for possible nitroglycerin administration to relieve chest pain, provided there are no contraindications such as hypotension. In contrast, other algorithms such as those addressing cardiac arrest or pediatric emergencies are focused on entirely different scenarios where the immediate priorities and interventions differ significantly from those required in a case of cocaine overdose. Thus, the recommendation to utilize the Acute Coronary Algorithm is based on its alignment with the direct complications and management protocols associated with cocaine toxicity and its effects on the heart.

9. What is one of the most dangerous complications of an untreated drug overdose?

A. Kidney failure

B. Cardiac arrest

C. Stomach bleeding

D. Seizures

Cardiac arrest is one of the most dangerous complications of an untreated drug overdose due to the profound effects certain drugs can have on the cardiovascular system. Many overdose situations involve substances that depress or alter the heart's rhythm, leading to significant disturbances such as bradycardia, tachycardia, or complete cardiac failure. Opioids, for instance, can significantly slow down the respiratory rate and heart function, resulting in hypoxia and decreased blood circulation, which can culminate in cardiac arrest. Other complications can arise from an overdose, such as kidney failure from prolonged hypoperfusion and potential seizures caused by certain stimulants or withdrawal effects. While stomach bleeding can occur in cases of specific overdoses, it is not as immediate or life-threatening as the risk of cardiac arrest. Therefore, understanding the critical impact that drug overdoses can have on heart function underscores the importance of rapid and effective intervention in these medical emergencies.

SAMPLE

10. Which vital sign is least likely affected by shock?

A. Heart rate

B. Blood pressure

C. Respiratory rate

D. Temperature

When assessing vital signs in the context of shock, temperature is often the least likely to be impacted. While heart rate, blood pressure, and respiratory rate typically change in response to shock—often increasing as the body attempts to compensate for decreased perfusion—temperature tends to remain relatively stable in the early stages of shock. In many shock states, such as hypovolemic or cardiogenic shock, the body's compensatory mechanisms prioritize maintaining blood flow to vital organs by adjusting heart rate and respiratory rate and by constricting blood vessels, which can affect blood pressure significantly. However, these changes do not always lead to immediate fluctuations in body temperature. In fact, temperature may actually drop in certain types of shock, such as septic shock, when the body is unable to mount an adequate immune response, but it is not typically a direct indicator of shock severity. Thus, temperature is considered the least likely of the vital signs to be directly influenced by shock, whereas the others are more readily altered as the body responds to the physiological stresses of shock.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://fisdapmedicalparamedic.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE