FISDAP PARAMEDIC TRAUMA PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://fisdapparamedictrauma.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is a significant risk factor for cardiac tamponade?

A. High blood pressure

B. A tension pneumothorax

C. Cerebral hemorrhage

D. Respiratory failure

2. Where is Hydrofluoric Acid commonly found?

A. In household cleaners

B. In drain cleaner and industrial settings

C. In food preservation

D. In cosmetic products

3. What is 'Cushing's triad' indicative of?

A. Increased intracranial pressure (ICP)

B. Hypovolemic shock

C. Septic shock

D. Cardiogenic shock

4. What injury might occur if a deformed dashboard is noted during assessment after a motor vehicle collision?

A. Liver rupture

B. Sternal fracture

C. Aortic tear

D. Head laceration

5. What is the most common cause of death in pediatric trauma?

A. Chest trauma

B. Hemorrhagic shock

C. Head trauma

D. Spinal injury

6. When is it deemed necessary to perform an emergency tracheostomy?

A. In cases of allergic reaction

B. When the patient is unconscious

C. In cases of airway obstruction where standard measures fail

D. During routine airway management

SAMPLE

7. What is the primary concern when treating suspected fractures?

A. Pain management

B. Stabilization to prevent further injury

C. Checking for neurovascular status

D. Preventing infection

8. In trauma care, the term 'hypotensive' refers to what condition?

A. Lower than normal blood pressure

B. Increased heart rate

C. Elevated blood sugar levels

D. Pale, clammy skin

9. What should be done for chemical burns?

A. Apply ice to the area immediately

B. Flush the area with water and remove clothing

C. Cover the burn with ointment

D. Leave the burn area covered until evaluation

10. How should you manage a patient experiencing a suspected chemical burn?

A. Apply ice to the affected area

B. Flush the affected area with copious amounts of water and remove clothing

C. Cover the burn with a sterile dressing

D. Apply topical ointments to the burn site

SAMPLE

Answers

SAMPLE

1. B
2. B
3. A
4. A
5. C
6. C
7. B
8. A
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is a significant risk factor for cardiac tamponade?

A. High blood pressure

B. A tension pneumothorax

C. Cerebral hemorrhage

D. Respiratory failure

Cardiac tamponade occurs when fluid accumulates in the pericardial space, exerting pressure on the heart and impairing its ability to pump effectively. One significant risk factor for developing cardiac tamponade is a tension pneumothorax. In a tension pneumothorax, air enters the pleural space and creates increasing pressure that can shift mediastinal structures and obstruct venous return to the heart. This abnormal pressure can lead to compromised circulation and, in some cases, can exacerbate or contribute to the development of cardiac tamponade, particularly if there is a concurrent injury involving the thorax or pericardium. The other options do not directly contribute to the risk of cardiac tamponade. High blood pressure (hypertension) is not a mechanism that leads to fluid accumulation in the pericardial space. Cerebral hemorrhage primarily affects the brain, with no direct impact on cardiac function in relation to pericardial fluid. Similarly, respiratory failure can result from various conditions, but it does not inherently cause cardiac tamponade.

2. Where is Hydrofluoric Acid commonly found?

A. In household cleaners

B. In drain cleaner and industrial settings

C. In food preservation

D. In cosmetic products

Hydrofluoric acid is prominently utilized in various industrial applications, which makes the second choice the most accurate. It is primarily found in settings such as metal etching, glass etching, and electronics manufacturing, where it plays a crucial role in processes that require the removal of oxides from metal surfaces or the etching of silicon wafers. In addition to its industrial uses, hydrofluoric acid is also found in certain types of drain cleaners; however, it's important to note that its presence in household products is limited and generally more regulated due to its highly corrosive nature and the associated health risks. The other options are less relevant because hydrofluoric acid is not typically associated with household cleaners, food preservation, or cosmetic products. While some cleaning agents may contain other acids or harsh chemicals, hydrofluoric acid is specifically used in more specialized and controlled environments due to its potential for causing severe chemical burns and systemic toxicity. Understanding the specific applications and handling precautions for hydrofluoric acid is vital for safety in both industrial and emergency response contexts.

SAMPLE

3. What is 'Cushing's triad' indicative of?

A. Increased intracranial pressure (ICP)

B. Hypovolemic shock

C. Septic shock

D. Cardiogenic shock

Cushing's triad is indicative of increased intracranial pressure (ICP) and is characterized by a specific set of vital sign changes: hypertension (high blood pressure), irregular or abnormal respirations, and bradycardia (slow heart rate). These signs occur as the body responds to pressure on the brain and the subsequent reduction in perfusion, leading to a potentially life-threatening condition. When ICP rises, it affects the brain's ability to function properly, which may manifest in these classic symptoms due to the body's attempts to maintain cerebral perfusion. Hypertension may develop as the body increases blood flow to the brain in response to the pressure, while the bradycardia may be a result of increased vagal tone or directly related to the brain's response to rising pressure. Additionally, the respiratory irregularities can stem from the brainstem being affected by this increased pressure. In contrast, the other options relate to different physiological conditions: hypovolemic shock is usually indicated by hypotension and tachycardia, septic shock is characterized by fever, tachycardia, and often a wide pulse pressure, while cardiogenic shock typically involves signs of heart failure, such as jugular venous distension and pulmonary congestion. Each of these conditions

4. What injury might occur if a deformed dashboard is noted during assessment after a motor vehicle collision?

A. Liver rupture

B. Sternal fracture

C. Aortic tear

D. Head laceration

The presence of a deformed dashboard during the assessment of a patient after a motor vehicle collision suggests that the patient experienced significant forces during the impact. In such cases, one potential injury that may occur is a liver rupture. This can happen if the patient's abdomen strikes the steering wheel or dashboard with enough force, causing the liver, which is located in the upper right quadrant of the abdomen, to be compressed against the bony structures of the rib cage. Liver contusions or lacerations are common in trauma cases where the abdomen is subjected to rapid deceleration and blunt force. The dashboard's deformation indicates that the passenger may have been thrown forward into it, increasing the likelihood of solid organ injuries, particularly to the liver. In contrast, while a sternal fracture, aortic tear, and head laceration are all potential injuries that can occur in motor vehicle collisions, they are less directly associated with a deformed dashboard. Sternal fractures may result from direct impact to the chest but are not as specific to dashboard crush injuries. An aortic tear is typically associated with severe deceleration and can occur with significant blunt trauma but is also not as closely linked to dashboard contact. Similarly, head lacerations may result from various types of impacts but

SAMPLE

5. What is the most common cause of death in pediatric trauma?

A. Chest trauma

B. Hemorrhagic shock

C. Head trauma

D. Spinal injury

The most common cause of death in pediatric trauma is head trauma. This assertion is supported by numerous studies indicating that among children, particularly those involved in blunt force injuries, falls, and motor vehicle collisions, head injuries account for a significant percentage of mortality. The pediatric population has a larger head-to-body ratio, which can lead to more severe outcomes from traumatic brain injuries compared to adults. Additionally, the developing brain of a child is particularly susceptible to injury, making it critical for clinicians to prioritize head trauma in trauma assessments and management. In pediatric trauma cases, while other forms of injury can certainly lead to severe outcomes, head trauma consistently emerges as the leading cause of fatalities. Effective prevention strategies and timely interventions are essential in addressing this significant risk factor.

6. When is it deemed necessary to perform an emergency tracheostomy?

A. In cases of allergic reaction

B. When the patient is unconscious

C. In cases of airway obstruction where standard measures fail

D. During routine airway management

An emergency tracheostomy is indicated in situations where there is a significant airway obstruction that cannot be resolved through standard measures such as intubation or other non-invasive techniques. This may occur due to swelling from trauma, severe allergic reactions, foreign bodies, or other conditions that obstruct the airway. When conventional methods fail to secure the airway and the patient demonstrates signs of respiratory distress or impending respiratory failure, establishing an airway through a tracheostomy can provide immediate access to the airway and allow for ventilation. In scenarios involving routine airway management or general unconsciousness, other interventions may be appropriate before considering an emergency tracheostomy. While an unconscious patient may require airway stabilization, they typically can still be managed with basic airway adjuncts or intubation if needed, as long as the obstruction is not absolute. Emergency tracheostomy is reserved for critical scenarios, emphasizing the urgency and necessity of the procedure when other options are inadequate.

7. What is the primary concern when treating suspected fractures?

A. Pain management

B. Stabilization to prevent further injury

C. Checking for neurovascular status

D. Preventing infection

The primary concern when treating suspected fractures is stabilization to prevent further injury. When a fracture occurs, there is a risk of additional damage to surrounding tissues, nerves, and blood vessels if the fractured bone moves or is improperly handled. Stabilizing the fracture minimizes movement, which helps reduce the risk of complications such as soft tissue injury or increased pain. This step is crucial for ensuring that the patient is not exacerbating their condition while awaiting further evaluation and treatment. While pain management, checking for neurovascular status, and preventing infection are also important considerations in the overall management of bone injuries, the immediate priority in the case of suspected fractures is to stabilize the affected area. This approach ensures that any maneuvers to assess neurovascular status or to manage pain do not compromise the injury further.

SAMPLE

8. In trauma care, the term 'hypotensive' refers to what condition?

A. Lower than normal blood pressure

B. Increased heart rate

C. Elevated blood sugar levels

D. Pale, clammy skin

The term 'hypotensive' specifically refers to a condition where blood pressure is lower than what is considered normal for an individual. This decreased blood pressure can lead to inadequate blood flow to organs and tissues, potentially resulting in shock or other serious complications. Understanding this is crucial in trauma care, as hypotension can be an indicator of severe bleeding or other life-threatening conditions. When assessing a trauma patient, recognizing hypotension is vital, as it often requires immediate intervention to restore adequate blood volume and perfusion. This may involve fluid resuscitation or other lifesaving measures. The other choices do not accurately define hypotension; increased heart rate, elevated blood sugar levels, and pale, clammy skin are signs that might accompany hypotension but do not define the condition itself.

9. What should be done for chemical burns?

A. Apply ice to the area immediately

B. Flush the area with water and remove clothing

C. Cover the burn with ointment

D. Leave the burn area covered until evaluation

For chemical burns, flushing the area with water and removing clothing is the appropriate response. The primary goal in managing chemical burns is to dilute and wash away the chemical agent to minimize tissue damage. Water is effective in removing many types of chemical agents and should be applied for an adequate duration, typically 15-30 minutes, to ensure thorough decontamination. Removing clothing is also crucial because it may harbor the chemical and continue to irritate the skin. It is essential to be cautious and avoid direct contact with the chemical or contaminated clothing when performing this step. In contrast, applying ice can further damage the skin and worsen the injury, as it causes constriction of blood vessels. Covering a burn with ointment may trap the chemical and is not advisable until the area is thoroughly flushed. Leaving the burn area covered without evaluation can delay necessary treatment and may allow the chemical to continue harming the tissue. Immediate decontamination is a critical first step in the management of chemical burns.

SAMPLE

10. How should you manage a patient experiencing a suspected chemical burn?

A. Apply ice to the affected area

B. Flush the affected area with copious amounts of water and remove clothing

C. Cover the burn with a sterile dressing

D. Apply topical ointments to the burn site

Managing a chemical burn requires immediate and appropriate action to minimize tissue damage and potential complications. Flushing the affected area with copious amounts of water is critical because it helps to dilute and remove the chemical agent that caused the burn. The longer the chemical remains in contact with the skin, the greater the risk of severe injury. Water is typically the safest and most effective way to decontaminate the area, as it not only helps to wash away the substance but also cools the burn site. In addition, removing clothing that may have come into contact with the chemical is essential to prevent further exposure. This can significantly reduce the severity of the burn and aid in proper treatment. After thoroughly flushing the area and ensuring that any residual chemical has been removed, further treatment can then be administered, such as covering the burn with a sterile dressing or seeking medical attention. Other options, such as applying ice, can cause vasoconstriction and worsen the injury, while applying topical ointments can trap the chemical in the skin and delay necessary decontamination. Covering the burn with a sterile dressing is appropriate after initial management, but it is not a substitute for the immediate flushing required in the case of a chemical burn. Thus, the recommended and correct approach is

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://fisdapparamedictrauma.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE