
FITZGERALD NURSE PRACTITIONER (NP) PRE-TEST PRACTICE (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://fitzgeraldnppretest.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which statin exhibits no risk for drug-food interaction with grapefruit juice?

A. Simvastatin

B. Rosuvastatin

C. Atorvastatin

D. Lovastatin

2. What is one recommended lifestyle modification for patients with hypertension?

A. Increase dietary sodium intake

B. Engage in regular physical activity

C. Reduce physical activity

D. Always avoid medication

3. Before starting isotretinoin, what is essential counseling for a female patient?

A. Treatment duration will be 1 month

B. Two forms of birth control are required before and during treatment

C. Isotretinoin has no serious side effects

D. It is safe to become pregnant during treatment

4. What heart sound is often associated with heart failure?

A. S1

B. S2

C. S3 gallop

D. Friction rub

5. For which condition is the use of oral decongestants least appropriate?

A. Allergic rhinitis

B. Common cold

C. Hypertension

D. Eustachian tube dysfunction

SAMPLE

6. What should a woman do if she forgot to take her combined oral contraceptive pill yesterday?

A. Take today's pill immediately and use back-up contraception for the next 7 days

B. Take the missed pill ASAP and today's pill at the usual time and no back-up contraception is needed

C. Take today's and yesterday's pills immediately and use back-up contraception until the start of menses

D. Take today's and yesterday's pills and encourage emergency contraception

7. Which statement about medication use during breastfeeding is accurate?

A. Only a few drugs are contraindicated in breastfeeding mothers

B. Up to 10% of the maternal dose can be passed through breastmilk

C. The "pump-and-dump" technique can eliminate medication exposure to a nursing infant

D. Breastfeeding should wait at least 2 hours following any medication dose

8. What should be done if a patient's TSH level remains elevated after 8 weeks of levothyroxine treatment?

A. Return in 4 weeks as more time is needed

B. Increase the dose to 100 mcg/day

C. Decrease the dose to 50 mcg/day

D. Switch to every other day dosing

9. Which of the following indicates well-controlled asthma in a patient?

A. FEV1 of 75% predicted

B. Nighttime awakening occurring about once per week

C. Asthma symptoms occurring about once or twice per week

D. SABA use about every other day

10. For a child with increasing symptoms of acute bacterial rhinosinusitis, what is the recommended approach?

A. Sinus puncture and culture

B. Gram stain of sinus discharge

C. Head MRI

D. No additional diagnostic testing is needed as this is largely a clinical diagnosis

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. C
5. C
6. B
7. A
8. B
9. C
10. D

SAMPLE

Explanations

SAMPLE

1. Which statin exhibits no risk for drug-food interaction with grapefruit juice?

A. Simvastatin

B. Rosuvastatin

C. Atorvastatin

D. Lovastatin

Rosuvastatin is the statin that exhibits no significant risk for drug-food interaction with grapefruit juice. This is primarily due to its pharmacokinetic profile. Unlike other statins, rosuvastatin is not significantly metabolized by the cytochrome P450 3A4 enzyme system, which is the pathway through which many statins, such as atorvastatin, simvastatin, and lovastatin, are affected by grapefruit juice. Grapefruit juice can inhibit this enzyme, potentially leading to higher concentrations of those statins in the bloodstream, which increases the risk of side effects, particularly muscle-related issues and liver enzyme elevation. Therefore, rosuvastatin can be considered safe for patients who consume grapefruit juice, making it a suitable option when dietary restrictions are a concern.

2. What is one recommended lifestyle modification for patients with hypertension?

A. Increase dietary sodium intake

B. Engage in regular physical activity

C. Reduce physical activity

D. Always avoid medication

Engaging in regular physical activity is a highly recommended lifestyle modification for patients with hypertension. Regular exercise can help lower blood pressure, improve overall cardiovascular health, and enhance the effectiveness of medications when they are needed. It is advised that adults aim for at least 150 minutes of moderate-intensity aerobic activity each week, such as walking, cycling, or swimming. Exercise not only contributes to weight management but also promotes better blood circulation and reduces stress, all of which are beneficial in managing hypertension. By contrast, increasing dietary sodium intake or reducing physical activity would likely exacerbate hypertension rather than help manage it. Sodium has a well-documented role in raising blood pressure, so its increase is typically discouraged for those with hypertension. Avoiding medication entirely is not advisable as many patients may require pharmacological treatment alongside lifestyle modifications to effectively manage their blood pressure.

SAMPLE

3. Before starting isotretinoin, what is essential counseling for a female patient?

A. Treatment duration will be 1 month

B. Two forms of birth control are required before and during treatment

C. Isotretinoin has no serious side effects

D. It is safe to become pregnant during treatment

Two forms of birth control are required before and during isotretinoin treatment because the medication is highly teratogenic, meaning it can cause severe birth defects if taken during pregnancy. This requirement is integral to minimizing the risk of fetal exposure to the drug. Both the U.S. Food and Drug Administration (FDA) and the iPLEDGE program emphasize the necessity for women of childbearing potential to use two reliable forms of contraception due to isotretinoin's potential adverse effects on a developing fetus. In providing this counseling, the healthcare provider underscores the seriousness of adhering to strict birth control measures, stresses the need for regular pregnancy testing, and discusses the implications of not following these protocols, which may include the risk of serious health consequences for the unborn child. Educating female patients about the potential risks clarifies the importance of this aspect of treatment while optimizing the medication's benefits in managing conditions like severe acne.

4. What heart sound is often associated with heart failure?

A. S1

B. S2

C. S3 gallop

D. Friction rub

The S3 gallop is often associated with heart failure, particularly in cases of volume overload or decreased ventricular compliance. When the heart fails to pump effectively, as seen in heart failure, there can be an increase in blood volume within the left ventricle during diastole. This results in a third heart sound (S3) due to the rapid filling of the ventricle. The sound is produced when blood from the atria enters an already filled ventricle, and it is typically best heard in the left lateral position using a stethoscope. In adults, the presence of an S3 is strongly indicative of heart failure and is often described as a low-pitched, galloping sound. S1 and S2 are heart sounds that reflect the closure of heart valves during the cardiac cycle; S1 marks the onset of ventricular systole, while S2 marks the end of systole. Though essential for assessing cardiac function, they are not specifically associated with heart failure. A friction rub, on the other hand, is typically related to pericarditis and not heart failure, indicating inflammation of the pericardium, the fibrous sac surrounding the heart. Therefore, the S3 gallop sound is the most relevant indicator

SAMPLE

5. For which condition is the use of oral decongestants least appropriate?

A. Allergic rhinitis

B. Common cold

C. Hypertension

D. Eustachian tube dysfunction

Oral decongestants, such as pseudoephedrine, are effective in relieving nasal congestion by constricting blood vessels in the nasal passages. However, in patients with hypertension, the use of these medications is least appropriate due to their potential to elevate blood pressure. The mechanism by which oral decongestants exert their effect can lead to increased sympathetic nervous system activity, resulting in vasoconstriction that may exacerbate existing hypertension or complicate its management. With hypertension as a primary condition, practitioners often seek alternative treatments for nasal congestion that do not increase blood pressure. Other options for treating conditions like allergic rhinitis, the common cold, or Eustachian tube dysfunction may not pose the same risk and can safely utilize oral decongestants when warranted. Therefore, the association between oral decongestants and potential adverse cardiovascular effects makes them least suitable for individuals with hypertension.

6. What should a woman do if she forgot to take her combined oral contraceptive pill yesterday?

A. Take today's pill immediately and use back-up contraception for the next 7 days

B. Take the missed pill ASAP and today's pill at the usual time and no back-up contraception is needed

C. Take today's and yesterday's pills immediately and use back-up contraception until the start of menses

D. Take today's and yesterday's pills and encourage emergency contraception

Taking the missed combined oral contraceptive pill as soon as the woman remembers, followed by taking today's pill at the usual time, is the recommended course of action. This approach allows the woman to maintain her contraceptive regimen without the need for additional back-up contraception, provided that the missed pill was remembered within a 24-hour window. Combined oral contraceptives are designed to offer flexibility in such situations, reinforcing the importance of continuing the current cycle effectively. If a woman misses a pill, she should not take more than one additional pill at a time to avoid exceeding the prescribed dosage. Additionally, if she has missed multiple pills or is unsure about the timing of her last pill, she should consult the instructions provided with her contraceptive or seek professional medical advice for clarity on her specific situation.

SAMPLE

7. Which statement about medication use during breastfeeding is accurate?

A. Only a few drugs are contraindicated in breastfeeding mothers

B. Up to 10% of the maternal dose can be passed through breastmilk

C. The "pump-and-dump" technique can eliminate medication exposure to a nursing infant

D. Breastfeeding should wait at least 2 hours following any medication dose

The statement regarding medication use during breastfeeding that is accurate is that only a few drugs are contraindicated in breastfeeding mothers. The vast majority of medications are considered safe for use while breastfeeding, and many medications can be transferred to breast milk in small amounts without posing a risk to the infant. The considerations for medication during lactation include the specific drug's safety profile, the timing of administration, and the potential effects on the infant. It's important for healthcare providers to assess the risks and benefits of any medication for breastfeeding mothers, but overall, most medications are compatible with breastfeeding. In contrast, the other statements may not accurately reflect current knowledge or practices. While some medications may pass into breastmilk, the percentage can vary widely based on multiple factors, thus the claim about a specific percentage is not universally applicable. The "pump-and-dump" technique has limited effectiveness in eliminating medication exposure, as many drugs are rapidly cleared from milk after administration; this practice is not typically recommended as a primary approach to manage medication transfer. Lastly, the recommendation to wait for a specific period post-medication before breastfeeding may depend on the particular medication and its pharmacokinetics rather than a blanket two-hour rule.

8. What should be done if a patient's TSH level remains elevated after 8 weeks of levothyroxine treatment?

A. Return in 4 weeks as more time is needed

B. Increase the dose to 100 mcg/day

C. Decrease the dose to 50 mcg/day

D. Switch to every other day dosing

When a patient's TSH (Thyroid Stimulating Hormone) level remains elevated after 8 weeks of levothyroxine treatment, an increase in the dose of levothyroxine is typically warranted. Levothyroxine is a synthetic form of the thyroid hormone thyroxine (T4) and is used to replace or supplement insufficient thyroid hormone levels in conditions such as hypothyroidism. The general guideline is to re-evaluate TSH levels approximately 6 to 8 weeks after initiating or adjusting levothyroxine therapy, as it takes about that time for the body to stabilize thyroid hormone levels in response to the medication. If TSH remains elevated, it indicates that the current dose is not sufficient to achieve the desired suppression of TSH. Increasing the dose allows for greater T4 availability, which can lead to a decrease in TSH levels and help bring the patient's thyroid function back to the normal range. It's important to note that the specific dose increase must be individualized based on the patient's response, weight, age, and any concomitant medical conditions. However, an increase in dosage is the most appropriate response when faced with persistently elevated TSH levels after a reasonable period of treatment.

SAMPLE

9. Which of the following indicates well-controlled asthma in a patient?

A. FEV1 of 75% predicted

B. Nighttime awakening occurring about once per week

C. Asthma symptoms occurring about once or twice per week

D. SABA use about every other day

Well-controlled asthma is characterized by infrequent symptoms and minimal limitations on daily activities. The correct answer indicates that asthma symptoms occur once or twice per week, which aligns with the criteria for good asthma management where patients aim to have symptoms less than twice a week. This frequency suggests the asthma is well-managed and that the patient is able to maintain normal activities without being significantly affected by the condition. In contrast, the other options suggest varying levels of poor asthma control. A FEV1 of 75% predicted indicates a moderate level of impairment, as a well-controlled patient would typically have a higher percentage of their predicted FEV1. Nighttime awakenings occurring about once per week signifies that the asthma is causing enough disruption to sleep, which is a key indicator of uncontrolled asthma. Finally, using short-acting beta-agonists (SABA) every other day suggests reliance on rescue medications for symptom relief, indicating that the asthma is not well-controlled since patients with adequately managed asthma should ideally use these medications infrequently or not at all.

10. For a child with increasing symptoms of acute bacterial rhinosinusitis, what is the recommended approach?

A. Sinus puncture and culture

B. Gram stain of sinus discharge

C. Head MRI

D. No additional diagnostic testing is needed as this is largely a clinical diagnosis

In cases of acute bacterial rhinosinusitis in children, the diagnosis is primarily clinical, based on the history and physical examination rather than extensive testing. Symptoms typically include nasal discharge, facial pain or pressure, and sometimes fever. These are usually sufficient to guide diagnosis and treatment without the need for additional diagnostic testing like imaging or invasive procedures. While sinus puncture and culture, Gram stain of sinus discharge, or head MRI could potentially provide more information, they are not standard practices for diagnosing acute bacterial rhinosinusitis in children due to concerns such as discomfort for the child, risk of complications, and added costs. Instead, the clinical diagnosis is often adequate, allowing healthcare providers to initiate the appropriate management, often with antibiotics if bacterial infection is suspected.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://fitzgeraldnppretest.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE