FOCUS ON ADULT HEALTH PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://focusonadulthealth.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Following two positive ELISA tests for HIV, which test is expected to confirm the diagnosis?**

   A. CD4+ T-cell count

   B. PCR test

   C. Western blot

   D. Bone marrow biopsy

2. **What are the typical symptoms of a myocardial infarction in women?**

   A. Chest pain and arm numbness

   B. Fatigue, shortness of breath, and nausea

   C. Severe headache and dizziness

   D. Palpitations and sweating

3. **What serum electrolyte level is a concern that may prompt the prescription of sodium polystyrene sulfonate in a client with acute kidney injury?**

   A. Sodium 135 mEq/L

   B. Calcium 9.0 mEq/L

   C. Potassium 5.9 mEq/L

   D. Magnesium 2.5 mEq/L

4. **What supplement should a nurse recommend daily for a client with hypoparathyroidism?**

   A. Vitamin C

   B. Calcium carbonate with vitamin D

   C. Iron supplements

   D. Magnesium

5. **Why is it important to monitor daily weights in patients with heart failure?**

   A. To detect fluid retention and worsening heart failure

   B. To assess physical activity levels

   C. To monitor dietary habits

   D. To calculate medication doses

SAMPLE

6. What does an elevated PT/INR indicate in patients on anticoagulants?

A. Lower blood pressure

B. Increased risk of bleeding

C. Reduced heart rate

D. Improved liver function

7. What is the primary characteristic of the TILTED position used in cardiac emergencies?

A. It increases cardiac output

B. It promotes venous return to the heart

C. It enhances respiratory function

D. It reduces blood pressure

8. What early sign of chronic kidney disease (CKD) should a nurse expect to assess upon admitting a client?

A. Increased urine output

B. Blood pressure of 168/94 mm Hg

C. Low grade fever

D. Decreased fatigue levels

9. Which recommendation should a nurse give a client on dietary modifications for improved nutritional status in COPD?

A. Encourage high-fat food choices

B. Incorporate gas-forming foods for digestion

C. Emphasize the importance of fluid intake during meals

D. Choose easy-to-chew, non-gas forming foods

10. Which assessment findings can indicate flail chest?

A. Ringing in the ears and nausea

B. Cyanosis and chest pain

C. Abdominal pain and vomiting

D. Low blood pressure and fever

SAMPLE

Answers

SAMPLE

1. C
2. B
3. C
4. B
5. A
6. B
7. B
8. B
9. D
10. B

SAMPLE

Explanations

SAMPLE

1. Following two positive ELISA tests for HIV, which test is expected to confirm the diagnosis?

A. CD4+ T-cell count

B. PCR test

C. Western blot

D. Bone marrow biopsy

The most reliable method for confirming a diagnosis of HIV after two positive ELISA tests is the Western blot test. This confirmatory test has a high specificity and is used to verify the presence of HIV antibodies in the bloodstream. The Western blot test detects specific proteins associated with the virus, providing a more definitive confirmation compared to initial screening tests like ELISA, which can sometimes yield false positives. While the CD4+ T-cell count and PCR test are important in the management and monitoring of HIV infection, they do not serve as definitive confirmatory tests after an initial positive diagnosis. The CD4+ T-cell count measures the immune system's health and is used to assess the impact of HIV on the body, while the PCR (Polymerase Chain Reaction) test detects the viral load but is often used in the context of tracking disease progression rather than confirming initial diagnosis. A bone marrow biopsy does not play a role in confirming HIV diagnoses. Thus, the Western blot test remains the gold standard for confirming the presence of HIV after initial positive ELISA results.

2. What are the typical symptoms of a myocardial infarction in women?

A. Chest pain and arm numbness

B. Fatigue, shortness of breath, and nausea

C. Severe headache and dizziness

D. Palpitations and sweating

The typical symptoms of a myocardial infarction (heart attack) in women often present differently than they do in men. While chest pain is a common symptom in both sexes, women frequently report symptoms that are less classic and can be more subtle. The symptoms such as fatigue, shortness of breath, and nausea are particularly significant because they can be misattributed to other conditions or stress, which adds to the challenge of timely recognition and treatment of a heart attack in women. Fatigue is a common and often overlooked symptom that may occur in the weeks leading up to a myocardial infarction, whereas shortness of breath can manifest with minimal exertion, and nausea can happen due to the visceral nature of cardiac pain—both of which are critical signals of a possible heart attack. In contrast, while other options contain symptoms that could be associated with cardiovascular issues, they are not as specific to myocardial infarctions in women as fatigue, shortness of breath, and nausea. Therefore, recognizing these symptoms is crucial for prompt medical intervention and improving outcomes for women experiencing a heart attack.

SAMPLE

3. What serum electrolyte level is a concern that may prompt the prescription of sodium polystyrene sulfonate in a client with acute kidney injury?

A. Sodium 135 mEq/L

B. Calcium 9.0 mEq/L

C. Potassium 5.9 mEq/L

D. Magnesium 2.5 mEq/L

The concern for prescribing sodium polystyrene sulfonate, commonly known as Kayexalate, centers around elevated serum potassium levels, especially in a patient with acute kidney injury (AKI). In this condition, the kidneys are unable to effectively filter and excrete potassium, leading to hyperkalemia, which is a potentially life-threatening electrolyte imbalance. A serum potassium level of 5.9 mEq/L is significantly elevated. Normal potassium levels range from about 3.5 to 5.0 mEq/L, and levels above 5.0 mEq/L indicate hyperkalemia. The risk factors associated with AKI, including the decreased ability to eliminate potassium, can cause levels to rise substantially. Immediate management is crucial, as high potassium levels can lead to serious complications such as cardiac arrhythmias or even cardiac arrest. Sodium polystyrene sulfonate works by exchanging sodium ions for potassium ions in the intestines, thus helping to reduce the body's potassium levels. In summary, the elevated potassium level at 5.9 mEq/L indicates a significant risk for serious health complications and justifies the use of sodium polystyrene sulfonate for treatment.

4. What supplement should a nurse recommend daily for a client with hypoparathyroidism?

A. Vitamin C

B. Calcium carbonate with vitamin D

C. Iron supplements

D. Magnesium

For a client with hypoparathyroidism, the daily recommendation of calcium carbonate with vitamin D is appropriate because this condition commonly leads to low levels of calcium in the blood due to insufficient parathyroid hormone. Parathyroid hormone is essential for regulating calcium levels, and its deficiency causes decreased absorption of calcium from the diet and increased excretion of calcium by the kidneys. Calcium carbonate serves as a direct source of calcium, which helps to correct the hypocalcemia associated with hypoparathyroidism. Additionally, vitamin D is necessary for the proper absorption of calcium in the intestines. Without sufficient vitamin D, the body would struggle to utilize dietary calcium effectively, regardless of intake levels. Therefore, this combination effectively addresses both the deficiency in blood calcium levels and promotes better absorption of calcium from the gastrointestinal tract. Other supplements, such as iron, magnesium, or vitamin C, do not directly address the core issue of calcium deficiency related to hypoparathyroidism and are not part of the standard management for this condition. Thus, recommending calcium carbonate with vitamin D directly targets the physiological needs of a client with hypoparathyroidism.

SAMPLE

5. Why is it important to monitor daily weights in patients with heart failure?

A. To detect fluid retention and worsening heart failure

B. To assess physical activity levels

C. To monitor dietary habits

D. To calculate medication doses

Monitoring daily weights in patients with heart failure is crucial because weight changes can be an early indicator of fluid retention, which is a common issue in this population. When heart failure worsens, the body may retain extra fluid due to the heart's reduced ability to pump effectively. A sudden increase in weight, often more than a couple of pounds in a day, can suggest that the patient is experiencing fluid overload. By tracking daily weights, healthcare providers and patients can identify these changes early and implement strategies such as adjusting medications, dietary modifications, or increasing diuretics to prevent further complications and hospitalizations. This proactive approach helps to manage symptoms and maintain the patient's overall well-being. The other choices do not focus specifically on the pivotal role that weight monitoring plays in managing heart failure. Assessing physical activity levels, monitoring dietary habits, and calculating medication doses, while relevant to overall patient care, do not directly reflect the immediate concern of fluid retention and its implications for heart failure management.

6. What does an elevated PT/INR indicate in patients on anticoagulants?

A. Lower blood pressure

B. Increased risk of bleeding

C. Reduced heart rate

D. Improved liver function

An elevated PT/INR indicates an increased risk of bleeding in patients who are on anticoagulants. PT (Prothrombin Time) and INR (International Normalized Ratio) are tests that measure how well and how quickly blood clots. Anticoagulants, such as warfarin, are medications that help reduce the clotting ability of blood to prevent conditions like thrombosis. When the PT/INR is elevated, it signifies that the blood is taking longer to clot, which can lead to an increased risk of bleeding complications. Maintaining the PT/INR within a therapeutic range is crucial for patients on anticoagulants to ensure that they benefit from the medication without experiencing adverse effects like excessive bleeding, which could occur if the levels are too high. This understanding emphasizes the importance of routine monitoring of PT/INR for patients on these medications to manage their treatment effectively. The other options do not relate directly to the implications of elevated PT/INR levels. For instance, lower blood pressure and reduced heart rate may occur due to various health conditions or treatments but do not correlate with clotting measures. Improved liver function is a different aspect altogether, and while liver function can affect clotting factors, it is not indicated

SAMPLE

7. What is the primary characteristic of the TILTED position used in cardiac emergencies?

A. It increases cardiac output

B. It promotes venous return to the heart

C. It enhances respiratory function

D. It reduces blood pressure

The primary characteristic of the TILTED position in cardiac emergencies is that it promotes venous return to the heart. This position is specifically designed to optimize blood flow back to the heart, which is essential in situations where cardiac output may be compromised. By encouraging venous return, the TILTED position can help maintain adequate preload and improve cardiac function, which is particularly crucial during emergencies when the body needs to maximize oxygen delivery to vital organs. Increasing venous return directly supports cardiac output because the heart works more effectively with a higher volume of blood returning to it. This can be beneficial in cases of hypotension or shock, where maintaining adequate circulation is a priority. While improving respiratory function and managing blood pressure are important considerations in cardiac care, they are not the primary focus of the TILTED position. Instead, enhancing venous return takes precedence, making this position a vital intervention in managing acute cardiac situations.

8. What early sign of chronic kidney disease (CKD) should a nurse expect to assess upon admitting a client?

A. Increased urine output

B. Blood pressure of 168/94 mm Hg

C. Low grade fever

D. Decreased fatigue levels

The presence of hypertension, particularly a blood pressure reading of 168/94 mm Hg, is a common early sign of chronic kidney disease (CKD). The kidneys play a crucial role in regulating blood pressure through the renin-angiotensin-aldosterone system. As kidney function declines, this regulation is disrupted, often leading to elevated blood pressure levels. Hypertension can be both a cause and a consequence of CKD, creating a cycle that may exacerbate kidney damage. Identifying elevated blood pressure upon admission is critical as it can prompt further assessments and interventions to manage kidney health and overall cardiovascular risk. In the context of CKD, increased urine output is typically not an early sign; instead, clients may experience altered urination patterns as the disease progresses. Similarly, a low-grade fever is not characteristic of CKD and may indicate a separate underlying infection or inflammatory process. Decreased fatigue levels would not be an expectation in CKD, as patients often report increased fatigue due to the accumulation of waste products in the body as kidney function declines. Recognizing hypertension as an early sign helps in monitoring and managing potential kidney-related complications effectively.

SAMPLE

9. Which recommendation should a nurse give a client on dietary modifications for improved nutritional status in COPD?

A. Encourage high-fat food choices

B. Incorporate gas-forming foods for digestion

C. Emphasize the importance of fluid intake during meals

D. Choose easy-to-chew, non-gas forming foods

For clients with Chronic Obstructive Pulmonary Disease (COPD), dietary modifications can play a significant role in managing symptoms and improving overall nutritional status. Recommending the selection of easy-to-chew, non-gas forming foods is particularly beneficial. Individuals with COPD often experience shortness of breath, which can be exacerbated by the physical effort required to eat. Foods that are easy to chew reduce the energy expenditure associated with eating, allowing the individual to focus their energy on breathing. Moreover, non-gas forming foods help minimize gastrointestinal discomfort and bloating, which can further impede breathing by putting pressure on the diaphragm and lungs. This dietary choice supports the dual needs of ensuring adequate nutrition while also considering the respiratory challenges that come with COPD. The other choices do not align with best practices for dietary management in this context. For example, high-fat foods can provide excess calories but may not be the most effective choice for overall nutrition, gas-forming foods can lead to increased discomfort and respiratory compromise, and emphasizing fluid intake during meals can increase the feeling of fullness and potentially decrease overall calorie intake. By focusing on easy-to-chew and non-gas forming foods, the nurse can guide the client towards a dietary strategy that supports both their nutritional requirements

10. Which assessment findings can indicate flail chest?

A. Ringing in the ears and nausea

B. Cyanosis and chest pain

C. Abdominal pain and vomiting

D. Low blood pressure and fever

Flail chest is characterized by a segment of the rib cage that becomes detached from the rest of the chest wall, typically due to multiple rib fractures. This condition can lead to impaired respiratory function and is often associated with significant trauma. The classic signs include chest pain, difficulty breathing, and paradoxical movement of the affected area during respiration. Cyanosis, which is a bluish discoloration of the skin and mucous membranes, indicates inadequate oxygenation and is common in patients with respiratory compromise, such as those with flail chest. Chest pain is also a typical symptom, stemming from rib fractures and the associated trauma. These assessment findings together point towards a compromised respiratory status due to the mechanical instability of the chest wall, making it a clear indicator of flail chest. The other options present symptoms that are more indicative of different conditions. Ringing in the ears and nausea often suggest neurological issues or the effects of certain medications but do not directly relate to respiratory trauma. Abdominal pain and vomiting are more representative of gastrointestinal disorders or internal injuries rather than flail chest. Low blood pressure and fever are signs that may indicate infection or systemic issues but do not specifically point to the mechanical disruption of the ribs typical of flail chest.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://focusonadulthealth.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE