FOUNDATION PHARMACIST RECRUITMENT PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://fdnpharmrecruitment.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is recommended for the treatment of chronic kidney disease in a primary care setting?

A. Use of calcium supplements

B. Atorvastatin 20mg OD and SGLT2 inhibitors

C. Increased dietary protein intake

D. Regular use of diuretics

2. Which medication interaction increases citalopram exposure?

A. Ranolazine

B. Proton pump inhibitors

C. Calcium channel blockers

D. Beta-blockers

3. Which drug class does bumetanide belong to?

A. Thiazide diuretic

B. Loop diuretic

C. Potassium-sparing diuretic

D. Carbonic anhydrase inhibitor

4. What is the first-line treatment for absence seizures?

A. Lamotrigine

B. Sodium valproate

C. Ethosuximide

D. Levetiracetam

5. What is the most appropriate pharmacological management for uncomplicated lower UTI in a pregnant woman?

A. Ciprofloxacin 500mg

B. Nitrofurantoin 100mg MR

C. Amoxicillin 500mg

D. Trimethoprim 200mg

SAMPLE

6. What would indicate hyperthyroidism in terms of TSH levels?

A. High TSH

B. Low TSH

C. Normal TSH

D. Variable TSH

7. Which of these drug classes can contribute to gastrointestinal bleeding if taken long-term without protective measures?

A. Analgesics

B. NSAIDs

C. Anticoagulants

D. Aspirin

8. What are the two main types of calcium channel blockers?

A. Dihydropyridine CCB and non-dihydropyridine CCB

B. Cardioselective and non-cardioselective CCB

C. Short-acting and long-acting CCB

D. Beta-adrenergic and calcium channel blockers

9. What is a primary characteristic of dihydropyridine calcium channel blockers (CCBs)?

A. Systemic vasodilation

B. Myocardium vasodilation

C. Reduction in heart rate

D. Increased contractility

10. Which impulse control issues can be a side effect of co-careldopa?

A. Increased motivation

B. Gambling, drugs, and sex

C. Social withdrawal

D. Enhanced focus

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. C
5. B
6. B
7. B
8. A
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. What is recommended for the treatment of chronic kidney disease in a primary care setting?

A. Use of calcium supplements

B. Atorvastatin 20mg OD and SGLT2 inhibitors

C. Increased dietary protein intake

D. Regular use of diuretics

The recommended treatment for chronic kidney disease (CKD) in a primary care setting includes the use of atorvastatin and SGLT2 inhibitors. Atorvastatin, a statin medication, is effective in managing cholesterol levels, which is crucial for patients with CKD as they have an increased risk of cardiovascular disease. SGLT2 inhibitors help in managing blood glucose levels, particularly in patients with diabetes, and have also been shown to provide renal protective effects by reducing progression to end-stage kidney disease and lowering the risk of hospitalization for heart failure. In contrast, while calcium supplements may be considered in specific scenarios related to mineral balance, particularly in patients with secondary hyperparathyroidism, they are not a standard treatment for CKD itself. Similarly, increased dietary protein intake is not typically recommended because it can exacerbate kidney function decline. Diuretics may be used to manage fluid overload in CKD patients but are not recommended as a primary treatment strategy for the condition itself. Hence, the combination of atorvastatin and SGLT2 inhibitors aligns with current guidelines for managing CKD, emphasizing cardiovascular health and renal protection.

2. Which medication interaction increases citalopram exposure?

A. Ranolazine

B. Proton pump inhibitors

C. Calcium channel blockers

D. Beta-blockers

Proton pump inhibitors (PPIs) are known to interact with various medications, including citalopram. The correct answer highlights that the use of PPIs can increase the exposure of citalopram in the body. This occurs due to the way these medications affect gastric pH levels and, consequently, the absorption and metabolism of citalopram. Citalopram is primarily metabolized by the liver enzymes of the cytochrome P450 family, particularly CYP2C19, and to some extent, CYP3A4. When proton pump inhibitors are taken, they can raise the pH in the stomach, which may modify the solubility and absorption characteristics of some drugs, including citalopram, resulting in higher plasma concentrations. Increased exposure to citalopram can elevate the risks of citalopram's side effects, such as QT interval prolongation, particularly in susceptible populations. Therefore, careful monitoring and adjustment of citalopram dosage may be necessary when a patient is concurrently taking a proton pump inhibitor to avoid potential adverse effects brought about by the interaction. Other medications, like ranolazine, calcium channel blockers, and beta-blockers, do not have a significant impact on the levels of c

SAMPLE

3. Which drug class does bumetanide belong to?

A. Thiazide diuretic

B. Loop diuretic

C. Potassium-sparing diuretic

D. Carbonic anhydrase inhibitor

Bumetanide is classified as a loop diuretic. This class of diuretics works primarily on the ascending loop of Henle in the kidneys, inhibiting the reabsorption of sodium and chloride ions. This leads to increased urine output, which is beneficial in managing conditions such as edema associated with heart failure, liver cirrhosis, or kidney disease. Loop diuretics are known for their potency and effectiveness, particularly in cases where rapid diuresis is required. They are different from thiazide diuretics, which act on the distal convoluted tubule and are typically used for hypertension and mild fluid retention. Potassium-sparing diuretics, on the other hand, are designed to conserve potassium in the body while promoting diuresis, and carbonic anhydrase inhibitors primarily work by reducing bicarbonate reabsorption in the kidneys. Thus, the mechanism of action and the anatomical sites where these drug classes exert their effects distinguish bumetanide as a loop diuretic, making it the correct answer to the question regarding its classification.

4. What is the first-line treatment for absence seizures?

A. Lamotrigine

B. Sodium valproate

C. Ethosuximide

D. Levetiracetam

The first-line treatment for absence seizures is ethosuximide. This medication is specifically designed to target and effectively manage absence seizures, which are characterized by brief episodes of impaired consciousness. Ethosuximide works by reducing the frequency of these seizures through its action on T-type calcium channels in the thalamic neurons, which play a crucial role in the generation of the rhythmic neuronal activity associated with absence seizures. While other medications, such as sodium valproate, lamotrigine, and levetiracetam, can be effective for various types of seizures including generalized seizures, ethosuximide is considered the most effective and preferred option specifically for the treatment of absence seizures due to its targeted mechanism of action and favorable side effect profile.

SAMPLE

5. What is the most appropriate pharmacological management for uncomplicated lower UTI in a pregnant woman?

A. Ciprofloxacin 500mg

B. Nitrofurantoin 100mg MR

C. Amoxicillin 500mg

D. Trimethoprim 200mg

The most appropriate pharmacological management for uncomplicated lower urinary tract infections (UTIs) in pregnant women is Nitrofurantoin 100mg MR. Nitrofurantoin is considered a first-line treatment for this condition during pregnancy due to its effectiveness against common uropathogens and its relatively safe profile for use in pregnant populations. During pregnancy, certain antibiotics are preferred or avoided due to potential effects on the developing fetus. Nitrofurantoin is generally utilized because it is thought to have minimal risks when taken during pregnancy, particularly during the second trimester. It acts by inhibiting bacterial cell wall synthesis, making it effective for treating uncomplicated UTIs. In contrast, other options may pose greater risks or are not recommended during pregnancy. For example, Ciprofloxacin is a fluoroquinolone, which is often avoided due to potential adverse effects on fetal development. Amoxicillin, while safe in pregnancy, may not be as effective against certain resistant strains commonly responsible for UTIs. Trimethoprim, especially when taken in the first trimester, raises concerns related to potential teratogenic effects. Thus, Nitrofurantoin stands out as the safest and most appropriate choice for managing uncomplicated lower UTIs in pregnant women, balancing efficacy and safety for both the mother and

6. What would indicate hyperthyroidism in terms of TSH levels?

A. High TSH

B. Low TSH

C. Normal TSH

D. Variable TSH

In the context of hyperthyroidism, low TSH (Thyroid-Stimulating Hormone) levels are indicative of the condition. Hyperthyroidism occurs when the thyroid gland is overactive, producing excess thyroid hormones (T3 and T4). When these hormones are elevated, they provide negative feedback to the pituitary gland, leading to a decreased secretion of TSH. As a result, when someone is experiencing hyperthyroidism, the expected laboratory finding would be low levels of TSH, often accompanied by high levels of T3 and T4. Understanding this feedback mechanism is crucial for interpreting thyroid function tests and diagnosing thyroid disorders accurately. Other options, like high, normal, or variable TSH levels, do not represent the characteristic findings of hyperthyroidism, as they do not align with the expected physiological response to elevated thyroid hormone levels.

SAMPLE

7. Which of these drug classes can contribute to gastrointestinal bleeding if taken long-term without protective measures?

A. Analgesics

B. NSAIDs

C. Anticoagulants

D. Aspirin

Long-term use of nonsteroidal anti-inflammatory drugs (NSAIDs) is well-documented to contribute to gastrointestinal bleeding due to their mechanism of action. NSAIDs work by inhibiting cyclooxygenase (COX) enzymes, which are important for the production of prostaglandins. Prostaglandins play a key role in maintaining the gastric mucosal lining, promoting mucus and bicarbonate secretion, and maintaining blood flow to the stomach lining. When these drugs inhibit prostaglandin synthesis, the protective mechanisms of the gastric mucosa are compromised, resulting in potential gastric irritation, ulceration, and bleeding. Although analgesics, anticoagulants, and aspirin can also affect the gastrointestinal tract, the specific association of NSAIDs with serious gastrointestinal complications is particularly strong. Aspirin, while related to the NSAID class, is often considered separately, as it has unique properties and gastrointestinal risks. However, it is important to note that when aspirin is used in a long-term setting, it can also cause bleeding; therefore, careful consideration and protective measures (like gastroprotective agents) are recommended for patients in need of long-term treatment with NSAIDs.

8. What are the two main types of calcium channel blockers?

A. Dihydropyridine CCB and non-dihydropyridine CCB

B. Cardioselective and non-cardioselective CCB

C. Short-acting and long-acting CCB

D. Beta-adrenergic and calcium channel blockers

Calcium channel blockers (CCBs) are categorized primarily into two categories: dihydropyridines and non-dihydropyridines. Dihydropyridine CCBs primarily target peripheral vascular smooth muscle, leading to vasodilation and are commonly used to treat hypertension and angina. Examples include amlodipine and nifedipine. On the other hand, non-dihydropyridine CCBs affect both the vascular smooth muscle and the conduction system of the heart, making them effective for conditions involving heart rhythm irregularities and supraventricular tachycardia. Examples include verapamil and diltiazem. Understanding the distinctions between these two types of CCBs is important for their appropriate clinical application, as they have different mechanisms of action and therapeutic uses. This classification aids pharmacists and healthcare providers in making informed decisions regarding treatment regimens for patients.

SAMPLE

9. What is a primary characteristic of dihydropyridine calcium channel blockers (CCBs)?

A. Systemic vasodilation

B. Myocardium vasodilation

C. Reduction in heart rate

D. Increased contractility

A primary characteristic of dihydropyridine calcium channel blockers (CCBs) is systemic vasodilation. These medications primarily target vascular smooth muscle, leading to the relaxation of these muscles and subsequent dilation of blood vessels. This mechanism results in decreased peripheral resistance, which can help reduce blood pressure in patients with hypertension. Dihydropyridine CCBs, such as amlodipine and nifedipine, do not significantly affect the heart's ability to contract or its rate of firing, which is why options related to myocardial vasodilation, reduction in heart rate, and increased contractility are not accurate primary characteristics. Their principal action is aimed at the vasculature rather than direct cardiac effects, differentiating them from non-dihydropyridine CCBs, which can affect the heart rate and contractility. Thus, the systemic vasodilation effect is the defining feature of dihydropyridine calcium channel blockers.

10. Which impulse control issues can be a side effect of co-careldopa?

A. Increased motivation

B. Gambling, drugs, and sex

C. Social withdrawal

D. Enhanced focus

Co-careldopa, a medication commonly used in the treatment of Parkinson's disease, has been associated with impulse control disorders as a potential side effect. Specifically, this can manifest in behaviors such as compulsive gambling, increased drug use, and hypersexuality. Understanding the mechanism, co-careldopa affects dopamine levels in the brain, which can enhance reward-seeking behaviors. As dopamine plays a crucial role in both motivation and the pleasure systems in the brain, elevating its activity can lead to exaggerated responses to stimuli that would typically be moderated. This can drive individuals to engage in impulsive behaviors that they might not have otherwise considered. Other choices do not accurately describe common side effects of co-careldopa. Increased motivation and enhanced focus may seem like potential outcomes of dopamine increase, but they do not reflect the negative side effects seen with impulse control disorders. Social withdrawal, on the other hand, generally reflects a lack of motivation or response to social stimuli, which is more likely a result of the underlying condition of Parkinson's rather than an impulse control issue linked to the medication.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://fdnpharmrecruitment.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE