HEALTHSTREAM PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://healthstream.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **An elevated central venous pressure (CVP) is most likely indicative of which condition?**

   A. Left heart failure

   B. Right heart failure

   C. Hypovolemia

   D. Fluid overload

2. **What is the priority assessment the nurse should make before administering digoxin (Lanoxin®)?**

   A. Blood pressure

   B. Respiratory rate

   C. Heart rate

   D. Fluid balance

3. **During an acute lateral myocardial infarction, which ECG leads will show changes?**

   A. I, aVL, V1, V2

   B. I, aVL, V5, V6

   C. II, III, aVF

   D. V3, V4, V5

4. **What symptom is most commonly reported by patients experiencing acute myocardial infarction?**

   A. Palpitations

   B. Crushing chest pain

   C. Fatigue

   D. Lightheadedness

5. **Why is it important to have an updated training curriculum in healthcare?**

   A. To maintain a peaceful work environment.

   B. To ensure alignment with changing regulations and medical standards.

   C. To enhance employee satisfaction.

   D. To reduce training costs.

SAMPLE

6. In pharmacology, how is the dose of a medication typically expressed?

A. As a percentage of body weight

B. As milligrams per body surface area

C. As units based on drug concentration

D. As milliliters of solution

7. What is the most effective way to evaluate a patient's understanding of instructions regarding digoxin administration?

A. Observation of adherence to the medication schedule

B. Teach-back method for verbalization of understanding

C. Written tests on medication side effects

D. Monthly follow-up appointments

8. A patient is newly intubated with normal breath sounds on one side and diminished on the other. What is the likely cause?

A. Intubation in the left mainstem bronchus

B. Intubation in the right mainstem bronchus

C. Pneumothorax on the left side

D. Congestive heart failure

9. What is the recommended rate at which IV saline should be administered to a patient with diabetic ketoacidosis during the first hour of treatment?

A. Drip continuously over 6 hours

B. Start a bolus infusion

C. Administer at a slow rate

D. Only when blood sugar levels are stable

10. What is a common treatment for severe sepsis with third-degree heart block?

A. Invasive monitoring

B. Transcutaneous pacing

C. Administration of beta-blockers

D. Fluid resuscitation

SAMPLE

Answers

SAMPLE

1. B
2. C
3. B
4. B
5. B
6. C
7. B
8. B
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. An elevated central venous pressure (CVP) is most likely indicative of which condition?

A. Left heart failure

B. Right heart failure

C. Hypovolemia

D. Fluid overload

An elevated central venous pressure (CVP) is primarily indicative of right heart failure. The CVP reflects the pressure in the thoracic vena cava, which gives an important insight into the filling pressures of the right atrium and overall right ventricular function. When the right side of the heart is unable to efficiently pump blood, it leads to an accumulation of blood in the venous system, which results in an increased CVP. In right heart failure, the heart's ability to manage the return flow of blood from the body can be compromised, causing elevated pressures in the venous system as blood backs up. This condition can present with associated symptoms like peripheral edema and jugular venous distention, further supporting the interpretation of elevated CVP. While elevated CVP can also occur in conditions like fluid overload, the primary condition that leads to this measurement being elevated is specific to right heart failure due to the direct impact it has on the circulation and venous return. Understanding this mechanism is key to diagnosing cardiovascular issues effectively.

2. What is the priority assessment the nurse should make before administering digoxin (Lanoxin®)?

A. Blood pressure

B. Respiratory rate

C. Heart rate

D. Fluid balance

The priority assessment before administering digoxin is the heart rate. Digoxin is a cardiac glycoside that affects heart function, and it can lead to bradycardia, or a significantly slowed heart rate, which can be dangerous. It is crucial to assess the patient's heart rate and ensure it is within a safe range before administration. A heart rate below the recommended threshold (usually less than 60 beats per minute for adults) may indicate that administration of digoxin could lead to further heart rate reduction, potentially resulting in adverse effects such as syncope, dizziness, or even cardiac arrest. Therefore, confirming that the heart rate is appropriate ensures patient safety and optimal therapeutic benefit from the medication. Other assessments like blood pressure, respiratory rate, and fluid balance are also important in overall patient evaluation but do not directly influence the immediate safety of administering digoxin as directly as heart rate does.

SAMPLE

3. During an acute lateral myocardial infarction, which ECG leads will show changes?

A. I, aVL, V1, V2

B. I, aVL, V5, V6

C. II, III, aVF

D. V3, V4, V5

In the context of an acute lateral myocardial infarction, the appropriate leads to monitor for changes are indeed I, aVL, V5, and V6. These leads correspond to the lateral wall of the left ventricle, which is the area of the heart that is typically affected during a lateral myocardial infarction. Lead I and aVL are positioned in such a way that they primarily observe the left lateral aspect of the heart. V5 and V6 are positioned on the left side of the chest and also visualize the lateral wall. Therefore, any changes such as ST-segment elevation or significant T-wave inversions observed in these leads would indicate ischemia or infarction in the lateral aspect of the heart. It is crucial for healthcare providers to identify these changes accurately, as they guide necessary interventions for patients experiencing myocardial infarctions. Recognition of these leads helps in assessing the extent and location of the myocardial damage, allowing for a more targeted treatment approach. The other choices represent other territories of the heart, which are not directly involved in a lateral MI, thus they would not show the characteristic changes associated with this specific type of infarction.

4. What symptom is most commonly reported by patients experiencing acute myocardial infarction?

A. Palpitations

B. Crushing chest pain

C. Fatigue

D. Lightheadedness

Patients experiencing acute myocardial infarction (AMI), commonly known as a heart attack, often report a symptom that is distinctly characterized as crushing chest pain. This type of pain is typically intense and may be described by patients as a feeling of pressure or tightness in the chest, sometimes radiating to the arms, back, neck, or jaw. The sensation is generally more severe than typical chest discomfort and can be accompanied by sweating, shortness of breath, or nausea. The crushing nature of the chest pain relates to the underlying pathophysiology of a heart attack, where the heart muscle is deprived of oxygen due to reduced blood flow, often resulting from a blocked coronary artery. Understanding this symptom is crucial for timely diagnosis and intervention, which can significantly affect patient outcomes in AMI cases. Other symptoms, while they may occur in conjunction with or instead of the crushing chest pain, are generally considered less characteristic of AMI when compared to this hallmark symptom. For instance, palpitations can be a sign of many different conditions, fatigue may be subtle and not as immediately alarming, and lightheadedness can result from various causes, including anxiety or low blood pressure. However, the classic presentation remains the intense chest pain that prompts immediate medical evaluation.

SAMPLE

5. Why is it important to have an updated training curriculum in healthcare?

A. To maintain a peaceful work environment.

B. To ensure alignment with changing regulations and medical standards.

C. To enhance employee satisfaction.

D. To reduce training costs.

Having an updated training curriculum in healthcare is essential to ensure alignment with changing regulations and medical standards. The healthcare sector is subject to frequent changes in laws, guidelines, and evidence-based practices. This dynamic environment requires healthcare professionals to stay informed about the latest standards to provide safe and effective patient care. By keeping training programs current, institutions can ensure that their staff is equipped with the necessary knowledge and skills to comply with new legal requirements, adhere to best practices, and implement updated medical technologies and techniques. Additionally, this alignment helps prevent compliance issues that could lead to legal repercussions or compromise patient safety. Regular updates to the training curriculum are important for addressing the evolving landscape of healthcare, including new treatment protocols, advances in medical technology, and shifts in patient care approaches. Thus, maintaining an up-to-date training curriculum directly impacts the quality of care provided and the organization's ability to function successfully in a regulated environment.

6. In pharmacology, how is the dose of a medication typically expressed?

A. As a percentage of body weight

B. As milligrams per body surface area

C. As units based on drug concentration

D. As milliliters of solution

In pharmacology, the dose of a medication is often expressed as units based on drug concentration. This method provides a standardized way to ensure that healthcare providers can accurately administer the correct amount of medication based on its potency and intended effect. By expressing the dose in terms of units, such as units of insulin or other medications, healthcare professionals can ensure that patients receive an appropriate and effective dose tailored to their specific needs. Understanding the concentration of the drug is essential, as different medications have different potencies. This approach allows for precise dosages that can be safely administered to patients, taking into account various factors such as age, weight, and health conditions. This method of expressing dosage is widely accepted in clinical practice and helps prevent dosing errors, ensuring patient safety and effective therapeutic outcomes.

SAMPLE

7. What is the most effective way to evaluate a patient's understanding of instructions regarding digoxin administration?

A. Observation of adherence to the medication schedule

B. Teach-back method for verbalization of understanding

C. Written tests on medication side effects

D. Monthly follow-up appointments

The teaching method known as "teach-back" is highly effective for evaluating a patient's understanding of instructions, especially in the context of medication administration like digoxin. This approach involves asking the patient to explain, in their own words, the instructions they were provided, such as how and when to take the medication. By having the patient verbalize the information, healthcare providers can determine not only if the patient has understood the instructions but also identify any misconceptions or gaps in knowledge. This method encourages active participation from the patient, fostering better retention and comprehension of the critical information related to their health. It is particularly relevant for medications like digoxin, where correct dosing and adherence are essential for effectiveness and safety. Observing adherence to the medication schedule, while useful, does not provide direct insight into the patient's understanding of the instructions. Similarly, written tests and monthly follow-up appointments may provide some information but do not directly assess comprehension at the time of teaching as effectively as the teach-back method does.

8. A patient is newly intubated with normal breath sounds on one side and diminished on the other. What is the likely cause?

A. Intubation in the left mainstem bronchus

B. Intubation in the right mainstem bronchus

C. Pneumothorax on the left side

D. Congestive heart failure

The scenario describes a patient who has been intubated and exhibits normal breath sounds on one side while having diminished breath sounds on the other. This clinical presentation is most consistent with intubation in the right mainstem bronchus. When an endotracheal tube is placed, it may inadvertently enter the right mainstem bronchus instead of the trachea. The right mainstem bronchus is anatomically straighter and shorter than the left, making it more likely for the tube to advance further into this bronchus. As a result, ventilation will primarily occur in the right lung, causing breath sounds to be normal on that side. Conversely, the left lung will receive inadequate ventilation, leading to diminished breath sounds on that side due to reduced airflow. While pneumothorax could also account for diminished breath sounds, it is less likely to be the immediate cause in a newly intubated patient where breath sounds were previously normal. Congestive heart failure typically presents with bilateral breath sounds on exam, not asymmetric findings as described. Therefore, the explanation for the clinical finding of diminished breath sounds on one side aligns most logically with intubation in the right mainstem bronchus.

SAMPLE

9. What is the recommended rate at which IV saline should be administered to a patient with diabetic ketoacidosis during the first hour of treatment?

A. Drip continuously over 6 hours

B. Start a bolus infusion

C. Administer at a slow rate

D. Only when blood sugar levels are stable

In the management of diabetic ketoacidosis (DKA), the initial treatment often includes the rapid administration of intravenous (IV) fluids. A bolus infusion of saline is recommended to quickly restore the patient's hydration status, improve tissue perfusion, and decrease glucose levels. This initial phase is crucial because DKA can lead to severe dehydration and electrolyte imbalances, which need to be corrected as a priority. Starting a bolus infusion ensures that the necessary fluid volume reaches the circulation quickly, effectively addressing the physiological needs of the patient right at the onset of treatment. It sets a foundation for subsequent management steps, such as insulin therapy and electrolyte replacement, which are critical in stabilizing the patient. The other options do not reflect the urgency and immediate action required in the first hour of DKA treatment. Continuous drip over several hours would delay the critical initial care. Administering at a slow rate would not sufficiently address the urgent fluid deficit seen in DKA. Waiting until blood sugar levels are stable before administering fluids is not advisable since rehydration needs to occur independently of glucose levels to support effective metabolic correction and patient stabilization.

10. What is a common treatment for severe sepsis with third-degree heart block?

A. Invasive monitoring

B. Transcutaneous pacing

C. Administration of beta-blockers

D. Fluid resuscitation

Transcutaneous pacing is often employed as a common treatment for severe sepsis complicated by third-degree heart block because it can temporarily stimulate the heart when there is a complete block in the electrical conduction system. In cases of third-degree heart block, the signals that regulate heartbeats can become ineffective, leading to bradycardia (a slower than normal heart rate) and potentially reducing cardiac output and tissue perfusion. Transcutaneous pacing uses external electrodes to deliver electrical impulses to the heart, effectively bypassing the block and increasing the heart rate. This intervention can be particularly critical in the setting of severe sepsis, where maintaining adequate circulation and organ perfusion is essential to prevent further complications. While other treatments may be necessary to address the underlying sepsis and stabilize the patient overall, transcutaneous pacing is a direct and immediate way to manage the heart's rhythm in the context of severe sepsis and heart block. The other options, while they may play important roles in managing sepsis or heart conditions separately, do not address the immediate need for pacing to manage the heart block effectively.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://healthstream.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE