INFECTIOUS DISEASE FIRST LINE TREATMENTS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://infectiousdiseasefirstline.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **What additional antibiotic is included in the empiric treatment for Bacterial Meningitis in patients over 50 years old?**

   A. Vancomycin

   B. Ampicillin

   C. Clofazimine

   D. Cefepime

2. **What is the first line treatment for Fascioliasis?**

   A. Praziquantel

   B. Triclabendazole

   C. Albendazole

   D. Ivermectin

3. **What is the recommended first line treatment for Pediculosis Corporis (Body Lice)?**

   A. Hygiene improvement and hot water washing

   B. Use of antibiotics

   C. Topical creams

   D. Oral medications

4. **What type of care is generally suggested for children with Chicken Pox?**

   A. Isolation and hydration

   B. Antibiotics and bed rest

   C. Corticosteroids and NSAIDs

   D. Analgesics and dietary restrictions

5. **What is the first line treatment for Leprosy (Paucibacillary)?**

   A. Dapsone + Rifampin

   B. Oral Griseofulvin

   C. IV Amphotericin B

   D. Topical Antifungals

SAMPLE

6. What is the recommended first line treatment for Zika Virus (ZIKV)?

A. Antibiotics and rest

B. Supportive care, fluids, and acetaminophen

C. Antiviral medication

D. Immediate vaccination

7. Which of the following is a primary method of controlling Pediculosis Corporis?

A. Frequent haircuts

B. Medication

C. Hygiene improvement

D. Quarantine

8. For the treatment of Chickungunya, what supportive measures are typically suggested?

A. NSAIDs, rest, and fluids

B. Cold compresses only

C. Analgesics and bed rest only

D. Acyclovir and fluids

9. What is the first line treatment for Cutaneous Larva Migrans?

A. Ivermectin

B. Albendazole

C. Chloroquine

D. Amoxicillin

10. Which treatment is typically used for Mycobacterium marinum infections?

A. Clarithromycin + Ethambutol

B. Dapsone + Rifampin

C. Amoxicillin + Clavulanate

D. Vancomycin

SAMPLE

Answers

SAMPLE

1. B
2. B
3. A
4. A
5. A
6. B
7. C
8. A
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. What additional antibiotic is included in the empiric treatment for Bacterial Meningitis in patients over 50 years old?

A. Vancomycin

B. Ampicillin

C. Clofazimine

D. Cefepime

In cases of bacterial meningitis, especially for patients over 50 years old, empiric treatment typically includes an antibiotic that covers Listeria monocytogenes due to the increased risk of this pathogen in older adults. Ampicillin is the antibiotic that provides effective coverage against Listeria, making it a critical part of the treatment regimen for this age group. In the context of empiric treatment, the combination of antibiotics usually includes a third-generation cephalosporin like ceftriaxone or cefotaxime and vancomycin for coverage against Streptococcus pneumoniae and Staphylococcus aureus, respectively. Including ampicillin in the regimen ensures that patients are protected against Listeria, addressing the specific risk factors associated with age. Other antibiotics listed do not provide the same level of coverage against Listeria. For example, vancomycin is primarily for gram-positive cocci coverage, and while it is an important drug in treating meningitis, it does not target Listeria. Clofazimine is not typically used for bacterial meningitis; it is primarily an anti-leprosy agent and has no relevance here. Cefepime is often utilized for broader-spectrum coverage but also does not specifically target Lister

2. What is the first line treatment for Fascioliasis?

A. Praziquantel

B. Triclabendazole

C. Albendazole

D. Ivermectin

The first line treatment for Fascioliasis is Triclabendazole. This anthelmintic medication specifically targets the liver flukes responsible for Fascioliasis, particularly Fasciola hepatica and Fasciola gigantica. Triclabendazole is highly effective against both the adult and immature stages of these parasites, which is crucial for treating the infection, especially since immature flukes can lead to more severe symptoms and complications. While other medications like Praziquantel, Albendazole, and Ivermectin are effective against various other parasitic infections, they do not demonstrate the same level of efficacy against Fasciola species. Praziquantel is primarily effective for schistosomiasis and some other trematodes, Albendazole is used for infections caused by nematodes and certain cestodes, and Ivermectin is typically used for roundworms and ectoparasites. Therefore, for treating Fascioliasis, Triclabendazole is the most appropriate choice due to its targeted action and proven success in clinical outcomes.

SAMPLE

3. What is the recommended first line treatment for Pediculosis Corporis (Body Lice)?

A. Hygiene improvement and hot water washing

B. Use of antibiotics

C. Topical creams

D. Oral medications

The recommended first-line treatment for Pediculosis Corporis (Body Lice) focuses on hygiene improvement and thorough washing of clothing and bedding in hot water. This approach is effective because body lice are primarily associated with poor hygiene and overcrowding, making improvements in cleanliness vital for controlling the infestation. By washing all clothing and affected bedding at high temperatures, typically above 130°F (54°C), you can kill adult lice as well as their eggs (nits). It is essential to dry the items on a hot setting as well to ensure that any remaining lice or nits are eradicated. While topical creams, oral medications, and antibiotics may be options in certain contexts for treating other types of lice or secondary infections, they are not the first line for body lice. Body lice do not require antibiotic treatment unless there is a secondary bacterial infection as a result of scratching and skin damage. Therefore, addressing hygiene and environmental factors is the most crucial and effective first step in managing Pediculosis Corporis.

4. What type of care is generally suggested for children with Chicken Pox?

A. Isolation and hydration

B. Antibiotics and bed rest

C. Corticosteroids and NSAIDs

D. Analgesics and dietary restrictions

For children with Chicken Pox, the recommended care primarily focuses on isolation and hydration. Isolation is crucial to prevent the spread of the varicella-zoster virus to others, especially vulnerable populations, such as newborns and immunocompromised individuals. Keeping the infected child at home until they are no longer contagious helps reduce transmission. Hydration is equally important, as it supports the child's overall well-being and helps manage symptoms. It's essential to ensure that the child drinks plenty of fluids, especially if they are not eating well or have fever, as dehydration can lead to further complications. The other options each feature treatments that do not align well with the standard management of Chicken Pox. Antibiotics are ineffective against viral infections like Chicken Pox; corticosteroids, while sometimes used in severe cases, may not be appropriate for routine cases due to potential adverse effects on the immune response; and dietary restrictions are not necessary as there are typically no specific dietary considerations for a child suffering from this illness. Analgesics may be used for symptom relief, but they are not the primary focus compared to the critical aspects of isolation and hydration.

SAMPLE

5. What is the first line treatment for Leprosy (Paucibacillary)?

A. Dapsone + Rifampin

B. Oral Griseofulvin

C. IV Amphotericin B

D. Topical Antifungals

The first-line treatment for Paucibacillary leprosy involves the combination of Dapsone and Rifampin. This regimen is specifically recommended because it effectively targets the Mycobacterium leprae bacterium, which causes leprosy. Dapsone works as a sulfone antibiotic that inhibits bacterial folate synthesis, while Rifampin is a potent rifamycin antibiotic that inhibits bacterial RNA synthesis. The combination of these two medications ensures a synergistic effect, which is essential for effectively managing the infection and preventing resistance. This treatment approach is aligned with the World Health Organization's guidelines for leprosy management, emphasizing the importance of multimodal therapy for effective patient outcomes. In cases of Paucibacillary leprosy, this dual therapy is particularly important as it reduces the risk of transmission and promotes healing of the skin lesions associated with the condition. Other treatments listed are not applicable for leprosy. For example, Griseofulvin is primarily used for dermatophyte infections of the skin, hair, and nails; Amphotericin B is an antifungal used for serious fungal infections; and topical antifungals address localized fungal infections but have no role in treating leprosy. Thus, the combination of D

6. What is the recommended first line treatment for Zika Virus (ZIKV)?

A. Antibiotics and rest

B. Supportive care, fluids, and acetaminophen

C. Antiviral medication

D. Immediate vaccination

The recommended first-line treatment for Zika Virus (ZIKV) focuses on supportive care, fluids, and the use of acetaminophen. This approach is appropriate because Zika virus infections are primarily self-limiting and often do not require specific antiviral therapies. Supportive care includes managing symptoms such as fever, rash, and body aches while ensuring that the patient remains hydrated, which is crucial for recovery. Acetaminophen is recommended to help reduce fever and alleviate pain, as it is generally considered safe and effective for symptomatic relief in these cases. While antibiotics are effective against bacterial infections, they do not treat viral infections like Zika. Antiviral medication is not specifically indicated for Zika, as no antiviral treatment has been proven effective against it to date. Vaccination is also not available for Zika virus prevention at this time, making supportive care the most suitable and recommended approach to manage the infection.

SAMPLE

7. Which of the following is a primary method of controlling Pediculosis Corporis?

A. Frequent haircuts

B. Medication

C. Hygiene improvement

D. Quarantine

The primary method of controlling Pediculosis Corporis, which is an infestation of body lice, is hygiene improvement. This approach focuses on the importance of maintaining cleanliness to reduce areas where lice thrive, such as in clothing and bedding. Regular washing of clothes, bedding, and personal items, as well as maintaining overall personal hygiene, can significantly diminish the presence of body lice and help prevent future infestations. While medication can be effective in treating Pediculosis Corporis by killing existing lice, it does not address the underlying issue of infestation cycles or re-infestation. Hygiene improvement gets to the root of the problem, reducing the contact with lice-infested materials. Frequent haircuts are not relevant in the case of body lice, as these lice infest clothing and not hair, which makes this choice ineffective. Quarantine might seem like a plausible option at first, but it does not directly address how to control or eliminate the lice, since the lice can easily spread through shared clothing or bedding rather than person-to-person contact. Thus, focusing on hygiene improvement is essential for effectively managing and controlling Pediculosis Corporis, making it the most appropriate choice among the options provided.

8. For the treatment of Chickungunya, what supportive measures are typically suggested?

A. NSAIDs, rest, and fluids

B. Cold compresses only

C. Analgesics and bed rest only

D. Acyclovir and fluids

The suggested supportive measures for the treatment of Chikungunya include nonsteroidal anti-inflammatory drugs (NSAIDs), rest, and increased fluid intake. NSAIDs are effective in managing the pain and inflammation associated with Chikungunya, which is crucial as this viral infection can lead to debilitating joint pain. Rest is essential to allow the body to recover, and staying hydrated supports overall health and aids in the recovery process. Cold compresses can be helpful for localized pain relief but are not a comprehensive treatment strategy for Chikungunya. Analgesics alone do not address the need for inflammation control or hydration, while antivirals such as acyclovir are not indicated for Chikungunya, as this condition does not respond to antiviral medications. Therefore, the combination of NSAIDs, rest, and fluids represents a holistic approach to managing the symptoms of Chikungunya effectively.

SAMPLE

9. What is the first line treatment for Cutaneous Larva Migrans?

A. Ivermectin

B. Albendazole

C. Chloroquine

D. Amoxicillin

The first-line treatment for cutaneous larva migrans is ivermectin. This is due to ivermectin's effectiveness against the larvae that cause this skin condition. Cutaneous larva migrans is typically caused by hookworm larvae, particularly Ancylostoma braziliense, which penetrate the skin and create a characteristic serpiginous rash. Ivermectin works by paralyzing and killing the larvae, leading to the resolution of symptoms. While other medications listed might treat different parasitic infections, they are not effective against the specific larvae causing cutaneous larva migrans. Albendazole is used for intestinal helminth infections but is less effective for skin larvae. Chloroquine is primarily used to treat malaria and some autoimmune diseases and doesn't have any effect on cutaneous larva migrans. Amoxicillin is an antibiotic that is not indicated for parasitic infections, making it irrelevant for the treatment of this condition. Thus, ivermectin is the preferred choice for its targeted action against the causative agents of cutaneous larva migrans.

10. Which treatment is typically used for Mycobacterium marinum infections?

A. Clarithromycin + Ethambutol

B. Dapsone + Rifampin

C. Amoxicillin + Clavulanate

D. Vancomycin

Mycobacterium marinum is a type of nontuberculous mycobacteria that can cause skin and soft tissue infections, often associated with exposure to aquatic environments. The first-line treatment for infections caused by Mycobacterium marinum typically involves a combination of clarithromycin and ethambutol. Clarithromycin is a macrolide antibiotic that inhibits bacterial protein synthesis, making it effective against the slow-growing Mycobacterium species. Ethambutol works by inhibiting the synthesis of the mycobacterial cell wall, complementing the action of clarithromycin and enhancing the overall effectiveness of the treatment regimen. The choice of this combination is based on their ability to penetrate the cells and target the mycobacterial infection effectively, as Mycobacterium marinum can be more resilient to treatment compared to other types of bacteria. This tailored approach helps in managing the infection while minimizing the risk of resistance development. In addition, other options provided, such as dapsone with rifampin, amoxicillin with clavulanate, and vancomycin, are not considered standard treatments for Mycobacterium marinum infections due to their differing mechanisms of action and effectiveness against this specific pathogen.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://infectiousdiseasefirstline.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE