INTERVENTIONAL RADIOLOGY REGISTRY PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://intervradiologyregistry.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which sterilization method has a short cycle time of 28-38 minutes?

A. Steam (Autoclaves)

B. Gas Plasma

C. Ethylene Oxide

D. Microwave

2. What is the purpose of using a stopcock during the flushing of a transducer?

A. To control the pressure in the device

B. To isolate the transducer

C. To prevent air bubbles

D. To increase flow rate

3. Which of the following is NOT a component of interventional supplies?

A. Needles

B. Wires

C. Embolization materials

D. Diagnostic contrast agents

4. Which equipment is NOT typically included in a TIPSS kit?

A. 14- or 16-ga needle

B. Occlusion balloon

C. Stethoscope

D. Self-expanding stent

5. Which arterial branch serves the descending colon and sigmoid flexure?

A. Anterior mesenteric artery

B. Sigmoid branches

C. Superior rectal artery

D. Left colic artery

6. Which of the following is NOT a characteristic of the Cobra catheter?

A. Visceral system usage

B. End hole design

C. Angled configuration

D. Side holes

SAMPLE

7. How long should a patient rest after undergoing kyphoplasty?

A. 1-2 hours

B. 12 hours

C. 24 hours

D. 48 hours

8. Which statement about AV block 1st degree is accurate?

A. All impulses are blocked from reaching the ventricles

B. Markedly shortened PR interval

C. Prolonged PR interval without missed beats

D. Isolated occurrence with no symptoms

9. What is a possible outcome for patients with contrast induced nephropathy before recovery?

A. Long-term dialysis

B. Avoidance of contrast media

C. Temporary dialysis

D. Increased medication dosage

10. Which item of equipment is important for the disposal of blood on a procedural tray?

A. Closed system for disposal of blood

B. Biohazard bag

C. Suction device

D. Waste bin with lid

SAMPLE

Answers

SAMPLE

1. B
2. C
3. D
4. C
5. B
6. D
7. C
8. C
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. Which sterilization method has a short cycle time of 28-38 minutes?

A. Steam (Autoclaves)

B. Gas Plasma

C. Ethylene Oxide

D. Microwave

The method of sterilization known as gas plasma is recognized for its efficient cycle time, typically lasting between 28 to 38 minutes. This technique utilizes low-temperature hydrogen peroxide vapor, which is then energized to produce gas plasma. This sterilization method is particularly advantageous for heat-sensitive medical instruments and devices that cannot withstand traditional sterilization methods such as steam. The relatively short cycle time allows for rapid turnover of instruments, making it a practical choice in busy healthcare settings where quick availability of sterile instruments is critical. Additionally, gas plasma sterilization does not leave toxic residues, as the byproducts of the process are benign, further enhancing its safety for both users and patients. Overall, gas plasma stands out in the sterilization landscape for its quick cycle duration, effectiveness against a wide range of organisms, and compatibility with sensitive materials, which is why it is the correct answer in this context.

2. What is the purpose of using a stopcock during the flushing of a transducer?

A. To control the pressure in the device

B. To isolate the transducer

C. To prevent air bubbles

D. To increase flow rate

Using a stopcock during the flushing of a transducer primarily serves to prevent air bubbles from entering the system. In the context of ultrasound or invasive monitoring, air bubbles can interfere with readings and create inaccuracies. By using the stopcock, the clinician can ensure a smooth flow of fluid, which helps to eliminate the risk of air being introduced into the transducer's pathway. This contributes not only to the reliability of the readings but also to the overall performance of the monitoring system. The design and function of stopcocks enable controlled access to the fluid pathway, allowing for the flushing process to be conducted smoothly. This control is crucial, as the presence of air bubbles could lead to erroneous measurements or create obstacles for proper fluid movement within the system. Overall, the use of a stopcock is integral to maintaining the integrity of the transducer's performance during procedures involving fluid dynamics.

SAMPLE

3. Which of the following is NOT a component of interventional supplies?

A. Needles

B. Wires

C. Embolization materials

D. Diagnostic contrast agents

In the context of interventional supplies, diagnostic contrast agents are not typically considered a direct component of the supplies used for the procedural aspect of interventional radiology. Interventional supplies generally refer to items that are actively used during invasive procedures to treat conditions or facilitate access to specific anatomical areas. These supplies include needles for puncturing vessels or tissues, wires that assist in guiding catheters or devices during a procedure, and embolization materials used to occlude blood vessels for various therapeutic reasons. Diagnostic contrast agents, on the other hand, are primarily used for imaging purposes before or during procedures to enhance the visibility of structures within the body on radiographic images. While they play a crucial role in the diagnostic phase of patient assessment, they do not serve as interventional supplies used directly to carry out the procedures themselves. Therefore, distinguishing between supplies that are directly utilized in interventions versus those used for imaging helps clarify why diagnostic contrast agents do not fit into the category of interventional supplies.

4. Which equipment is NOT typically included in a TIPSS kit?

A. 14- or 16-ga needle

B. Occlusion balloon

C. Stethoscope

D. Self-expanding stent

The equipment typically included in a TIPSS (Transjugular Intrahepatic Portosystemic Shunt) kit is specifically tailored to the interventional procedure involved in creating a shunt between the hepatic and portal veins to help manage complications arising from portal hypertension, such as variceal bleeding. The inclusion of a 14- or 16-gauge needle is critical for the initial access and the creation of the shunt. The occlusion balloon is essential for temporarily blocking the blood flow during the procedure, which aids in the proper placement of the stent. The self-expanding stent is necessary because it maintains the patency of the newly created shunt after the procedure. In contrast, a stethoscope is not a required tool for the TIPSS procedure itself. While stethoscopes are commonly used in clinical practice for auscultation or monitoring patients' heart and lung sounds, they do not play a direct role in the technical aspects of the TIPSS procedure. Thus, it does not belong in the specialized kit that contains the equipment necessary for performing this particular interventional radiology procedure.

SAMPLE

5. Which arterial branch serves the descending colon and sigmoid flexure?

A. Anterior mesenteric artery

B. Sigmoid branches

C. Superior rectal artery

D. Left colic artery

The choice related to the arterial branch that serves the descending colon and sigmoid flexure is indeed the sigmoid branches. The sigmoid branches arise from the inferior mesenteric artery and specifically supply blood to the sigmoid colon, which is part of the large intestine that connects the descending colon to the rectum. These branches ensure adequate blood flow to the sigmoid flexure, a crucial area involved in the storage of fecal matter before defecation. The descending colon receives blood supply primarily from the left colic artery, which is a branch of the inferior mesenteric artery. However, the specific supply to the sigmoid colon is accomplished by the sigmoid branches. This targeted vascular supply is essential for the functioning and health of the colon, facilitating nutrient absorption and waste management. Understanding the vascular anatomy of the colon is important in interventional radiology, particularly when considering procedures that might involve the inferior mesenteric artery or its branches. It is essential to know which arteries vascularize specific segments of the colon for effective treatment planning and intervention strategies.

6. Which of the following is NOT a characteristic of the Cobra catheter?

A. Visceral system usage

B. End hole design

C. Angled configuration

D. Side holes

The Cobra catheter is primarily recognized for its unique characteristics that enhance its performance in vascular access and procedures. Its design allows for easy navigation through the vascular system, particularly when accessing the visceral arteries. Considering the characteristics outlined, the end hole design is significant because it facilitates efficient contrast flow during angiographic procedures. The angled configuration is also essential, as it aids in steering the catheter into the optimal position within the vascular anatomy. These features collectively contribute to its effectiveness in various interventional radiology procedures. However, the Cobra catheter is not characterized by side holes. Unlike other catheter designs that incorporate multiple side holes to allow for lateral drainage or distribution of fluids, the Cobra catheter operates predominantly through its end hole, which is designed to minimize turbulence and provide a streamlined passage for contrast media. Therefore, identifying the absence of side holes as a characteristic sets this option apart from the others that are inherent to the Cobra catheter's design and function.

7. How long should a patient rest after undergoing kyphoplasty?

A. 1-2 hours

B. 12 hours

C. 24 hours

D. 48 hours

After undergoing kyphoplasty, a patient is typically advised to rest for approximately 24 hours. This period allows the body to stabilize and reduces the potential for complications such as bleeding or a post-procedure fracture. During this time, patients are encouraged to minimize physical activity and avoid lifting heavy objects to promote healing. The 24-hour rest period is significant as it helps in the initial recovery after an invasive procedure where cement is injected into a vertebral body to stabilize fractures and relieve pain. Following this initial recovery phase, patients can usually begin to resume normal activities with caution as advised by their healthcare provider. It's important to provide patients with guidance on their activity levels following the procedure to ensure optimal healing and avoid unnecessary strain on the affected area.

SAMPLE

8. Which statement about AV block 1st degree is accurate?

A. All impulses are blocked from reaching the ventricles

B. Markedly shortened PR interval

C. Prolonged PR interval without missed beats

D. Isolated occurrence with no symptoms

The statement regarding AV block first degree that is accurate describes prolonged PR interval without missed beats. In first-degree AV block, there is a delay in the conduction of electrical impulses through the atrioventricular node, which manifests as a prolonged PR interval on an electrocardiogram (ECG). This prolonged interval indicates that the time taken for impulses to travel from the atria to the ventricles is longer than normal. In first-degree AV block, every impulse generated in the atria successfully reaches the ventricles, which is why there are no missed beats. Patients typically do not experience symptoms due to the typically benign nature of this block, although it could be an indication of an underlying condition in some patients. This clearly distinguishes first-degree AV block from more severe types of AV block, such as second-degree or third-degree blocks, where some impulses are indeed blocked from reaching the ventricles. The other options do not accurately represent the characteristics of first-degree AV block, specifically the PR interval's prolonged nature.

9. What is a possible outcome for patients with contrast induced nephropathy before recovery?

A. Long-term dialysis

B. Avoidance of contrast media

C. Temporary dialysis

D. Increased medication dosage

The situation described involves contrast-induced nephropathy (CIN), which is an impairment of kidney function that can occur after the administration of contrast media during diagnostic procedures or interventional radiology. A possible outcome for patients who experience CIN before they begin to recover is the need for temporary dialysis. Temporary dialysis may be necessary to manage the acute kidney injury that occurs as a result of CIN. This type of dialysis serves to support the kidneys while they heal and return to normal function. In many cases, the kidneys can recover fully over time, and once they regain adequate function, the patient can be weaned off dialysis. This outcome is pertinent to patients who may have experienced a significant decline in their renal function due to the contrast agent, especially if there are underlying conditions, such as diabetes or hypertension, that could exacerbate the situation. On the other hand, long-term dialysis is typically associated with chronic kidney failure rather than with acute events like CIN, which often resolves. Avoidance of contrast media, while a sensible precaution for high-risk patients, doesn't directly address the immediate consequences of CIN for those who have already received it. Lastly, increasing medication dosage is not a standard response to managing CIN; rather, careful management of the patient's existing medications may

SAMPLE

10. Which item of equipment is important for the disposal of blood on a procedural tray?

A. Closed system for disposal of blood

B. Biohazard bag

C. Suction device

D. Waste bin with lid

The correct response highlights the importance of a closed system for the disposal of blood during medical procedures. A closed system minimizes exposure to blood and helps prevent spills or contamination risks that can occur if blood is not contained properly. In interventional radiology, where the use of instruments and drainage of blood are routine, utilizing a closed system ensures that any blood generated during a procedure is directed safely and efficiently into a secure receptacle, reducing risks of infection and environmental hazards. While other options, like biohazard bags or waste bins with lids, also play roles in handling biohazardous material, they do not specifically emphasize the containment and elimination of blood products in a procedural context. A suction device, on the other hand, is used to remove blood during a procedure but does not address the final disposal of the blood once it is collected. Therefore, a closed system specifically designed for blood disposal aligns best with safety protocols and best practices in interventional radiology.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://intervradiologyregistry.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE