ITE CARDIOLOGY PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://itecardiology.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. In cardiogenic shock with low cardiac index and MAP around 75 mmHg, which vasoactive agent is preferred initially?

A. Norepinephrine

B. Dobutamine

C. Vasopressin

D. Epinephrine

2. Which statement best describes vasopressin's relation to dobutamine in cardiogenic shock?

A. It provides inotropy and reduces afterload.

B. It has no effect on vascular tone.

C. It increases heart rate exclusively.

D. It increases SVR and lacks inotropy; thus less favorable with dobutamine.

3. In heart transplant recipients, which autonomic innervation is absent at baseline?

A. Parasympathetic

B. Sympathetic

C. Both

D. None

4. According to ACC/AHA guidelines, patients with no active cardiac conditions and functional capacity greater than 4 METs generally require what regarding preoperative cardiac testing?

A. Routine noninvasive testing

B. No further cardiac testing

C. Preoperative echocardiography

D. Stress testing always in all cases

5. Which method is quickest and most reliable to confirm proper placement of a central venous line prior to dilation?

A. Ultrasound

B. Echocardiography

C. Observation of blood color

D. Pressure monitoring (manometry)

SAMPLE

6. For a patient with aortic stenosis undergoing anesthesia, which pre-induction strategy helps reduce the risk of hypotension?

A. Phenylephrine infusion prior to induction

B. High-dose volatile induction without vasopressors

C. Rapid sequence induction with no fluids

D. Balanced anesthesia with high end-diastolic volume augmentation

7. In an ECG, a large R wave in lead V1 should prompt consideration of which diagnosis?

A. Posterior wall MI

B. Right ventricular hypertrophy

C. Muscular dystrophy

D. Right atrial enlargement

8. Which volatile anesthetic is most associated with ischemic preconditioning?

A. Sevoflurane

B. Isoflurane

C. Desflurane

D. Nitrous oxide

9. What is a key perioperative goal for LQTS patients?

A. Provide adequate anesthesia and analgesia to prevent sympathetic stimulation

B. Induce hypotension to reduce heart workload

C. Avoid anxiolysis to maintain alertness

D. Advance rapid extubation without analgesia

10. Why do centrifugal cardiopulmonary bypass circuits require flowmeters on the arterial limb?

A. Flow is constant during nonpulsatile perfusion

B. Flow varies with preload and afterload

C. To measure venous return

D. To regulate circuit temperature

SAMPLE

Answers

SAMPLE

1. B
2. D
3. A
4. B
5. D
6. A
7. B
8. A
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. In cardiogenic shock with low cardiac index and MAP around 75 mmHg, which vasoactive agent is preferred initially?

A. Norepinephrine

B. Dobutamine

C. Vasopressin

D. Epinephrine

When perfusion is mainly limited by low cardiac output rather than by extreme hypotension, the first priority is to increase the heart's ability to pump. Dobutamine is a beta-1 agonist that boosts myocardial contractility and stroke volume, which raises cardiac index. Because it also has mild beta-2–mediated vasodilation, it can reduce afterload a bit, helping the weakened ventricle eject blood more effectively. With a mean arterial pressure around 75 mmHg, this inotropic boost can improve forward flow without needing strong vasoconstriction. If blood pressure were to fall or remain insufficient, a vasopressor such as norepinephrine can be added to maintain perfusion pressure. In contrast, vasopressin is typically used as an adjunct to raise MAP, and epinephrine raises contractility but carries higher risks of tachyarrhythmias and increased myocardial oxygen demand, making them less ideal as the initial choice in this scenario.

2. Which statement best describes vasopressin's relation to dobutamine in cardiogenic shock?

A. It provides inotropy and reduces afterload.

B. It has no effect on vascular tone.

C. It increases heart rate exclusively.

D. It increases SVR and lacks inotropy; thus less favorable with dobutamine.

The key idea is how these drugs affect afterload and contractility. Dobutamine mainly boosts heart contractility (inotropy) and can lower systemic vascular resistance a bit, helping forward flow. Vasopressin, on the other hand, raises systemic vascular resistance through V1 receptor–mediated vasoconstriction and has no direct inotropic effect. So adding vasopressin to dobutamine increases afterload without increasing contractility, which can blunt the improvement in cardiac output that dobutamine provides. That's why the statement describing vasopressin as increasing SVR and lacking inotropy, making it less favorable with dobutamine, is best.

3. In heart transplant recipients, which autonomic innervation is absent at baseline?

A. Parasympathetic

B. Sympathetic

C. Both

D. None

The key idea is that a transplanted heart is denervated from the moment of surgery. The parasympathetic (vagal) input to the heart is lost at baseline because the vagus connections are severed during the transplant. Without vagal tone, the heart lacks the usual braking effect on heart rate, so resting heart rate tends to be higher and beat-to-beat variability is reduced. In contrast, any influence on the heart that remains comes from circulating catecholamines acting on the heart's receptors, and some reinnervation—especially sympathetic—may occur slowly over time in some patients, but parasympathetic innervation is not restored reliably in the early period.

SAMPLE

4. According to ACC/AHA guidelines, patients with no active cardiac conditions and functional capacity greater than 4 METs generally require what regarding preoperative cardiac testing?

A. Routine noninvasive testing

B. No further cardiac testing

C. Preoperative echocardiography

D. Stress testing always in all cases

Having functional capacity above 4 METs shows you have enough cardiac reserve to handle most noncardiac surgeries without routine testing. When there are no active cardiac conditions and the patient can perform activities at this level, ACC/AHA guidelines recommend no further preoperative cardiac testing, because such testing doesn't improve outcomes in this scenario. Testing is kept for those with symptoms, poor functional capacity (<4 METs), active cardiac disease, or situations where test results would change management.

5. Which method is quickest and most reliable to confirm proper placement of a central venous line prior to dilation?

A. Ultrasound

B. Echocardiography

C. Observation of blood color

D. Pressure monitoring (manometry)

Confirming proper placement of a central venous line quickly relies on real-time pressure feedback to verify you're in the vein before proceeding to dilation. By connecting the catheter site to a pressure transducer (manometer), you can observe the waveform and the absolute pressure. Venous access shows a low, non-arterial pressure waveform, consistent with central venous pressures, whereas arterial placement would reveal a much higher, pulsatile arterial waveform. This immediate pressure verification lets you proceed confidently with dilation only when venous placement is confirmed, and halt or reposition if an arterial pattern is detected. Ultrasound, while excellent for guiding vessel entry, doesn't by itself confirm that the catheter remains in the venous system beyond the entry point. Echocardiography can visualize catheter position but requires more time and equipment. Observing blood color is unreliable.

SAMPLE

6. For a patient with aortic stenosis undergoing anesthesia, which pre-induction strategy helps reduce the risk of hypotension?

A. Phenylephrine infusion prior to induction

B. High-dose volatile induction without vasopressors

C. Rapid sequence induction with no fluids

D. Balanced anesthesia with high end-diastolic volume augmentation

The main idea is that in aortic stenosis, the heart relies on a stable afterload to maintain blood pressure because the stroke volume is fixed. Induction of anesthesia brings vasodilation and venous pooling, which can cause a big drop in blood pressure and compromise coronary perfusion. Starting a phenylephrine infusion before induction provides its pure alpha-1 vasoconstrictive effect, maintaining systemic vascular resistance and arterial pressure as anesthesia takes effect. This helps preserve preload and afterload, keeping the left ventricle as stable as possible and reducing the risk of hypotension during induction. Other approaches that avoid pre-induction vasopressor support tend to allow larger drops in SVR or fail to optimize afterload, increasing the chances of hypotension. For instance, inducing with high-dose volatile agents without vasopressors can cause significant vasodilation; rapid sequence induction with no fluids risks volume depletion; while optimizing preload with fluids alone may not adequately counteract the vasodilatory phase. Phenylephrine before induction directly counters the main problem in AS during anesthesia: loss of vascular tone leading to hypotension.

7. In an ECG, a large R wave in lead V1 should prompt consideration of which diagnosis?

A. Posterior wall MI

B. Right ventricular hypertrophy

C. Muscular dystrophy

D. Right atrial enlargement

A large R wave in V1 indicates the heart's depolarization vector is directed toward the right ventricle. In the normal pattern, the R wave in V1 is small and gradually grows as you move laterally to the chest leads. When the right ventricle enlarges or experiences strain, the electrical vector shifts to the right, making the R wave in V1 and often V2 larger (R/S ratio in V1 >1). This is a classic sign of right ventricular hypertrophy and is often accompanied by right axis deviation and secondary repolarization changes. While a tall R wave in V1 can appear with posterior myocardial infarction due to mirror-image effects, that scenario would be accompanied by other posterior-involved changes on the ECG, and the pattern aligns most directly with RVH. Right atrial enlargement would manifest as tall, peaked P waves rather than a prominent R in V1, and muscular dystrophy lacks a specific ECG hallmark like this.

SAMPLE

8. Which volatile anesthetic is most associated with ischemic preconditioning?

A. Sevoflurane

B. Isoflurane

C. Desflurane

D. Nitrous oxide

Ischemic preconditioning can be mimicked by volatile anesthetics, which trigger protective signaling in the heart that helps it withstand a later period of ischemia. The protective effect comes from pathways that preserve mitochondrial function, such as opening mitochondrial ATP-sensitive potassium channels and activating downstream kinases, which together reduce calcium overload and limit cell injury during reperfusion. Sevoflurane is most associated with this phenomenon because the data consistently show a robust and reliable cardioprotective response with sevoflurane exposure before ischemic events. This translates into smaller infarct sizes and better outcomes in models and clinical settings of ischemia-reperfusion. While other volatile anesthetics can also induce preconditioning, sevoflurane has the strongest and most consistent association in the literature, whereas nitrous oxide does not produce this effect.

9. What is a key perioperative goal for LQTS patients?

A. Provide adequate anesthesia and analgesia to prevent sympathetic stimulation

B. Induce hypotension to reduce heart workload

C. Avoid anxiolysis to maintain alertness

D. Advance rapid extubation without analgesia

Blunting sympathetic stimulation during the perioperative period is crucial in LQTS patients because pain, anxiety, and airway/line stimulation raise catecholamines, which can prolong the QT interval and precipitate dangerous arrhythmias like torsades de pointes. Providing adequate anesthesia and effective analgesia minimizes these autonomic surges, helping keep heart rate, blood pressure, and the QT interval more stable. Inducing hypotension or skipping analgesia would not address the underlying trigger and could worsen perfusion or trigger stress responses, while avoiding anxiolysis would do the opposite by increasing sympathetic drive; rapid emergence without analgesia similarly risks a surge in sympathetic tone.

10. Why do centrifugal cardiopulmonary bypass circuits require flowmeters on the arterial limb?

A. Flow is constant during nonpulsatile perfusion

B. Flow varies with preload and afterload

C. To measure venous return

D. To regulate circuit temperature

On centrifugal CPB circuits, flow is not fixed by pump speed alone; the delivered flow depends on the pressure the circuit presents (afterload) and how much venous return comes in (preload). If afterload increases or preload decreases, the same pump speed yields less flow. Because of this, an arterial limb flowmeter is essential to know the actual systemic flow in real time and to adjust pump speed to achieve the target perfusion and oxygen delivery. This is especially important in nonpulsatile perfusion, where you don't have natural pulsatile cues to gauge flow. The arterial flowmeter specifically measures the flow delivered to the patient, not venous return, and it isn't used to regulate temperature.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://itecardiology.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE