
ITE NEPHROLOGY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://itenephrology.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. If SIADH does not respond to fluid restriction in the outpatient setting, which agent is commonly used?

A. Furosemide

B. Demeclocycline

C. Hypertonic saline

D. Conivaptan

2. Which clinical features strongly suggest ANCA-associated GN (Wegener's granulomatosis)?

A. Nasal congestion, ulceration, chronic sinusitis, and pulmonary nodules/infiltrates

B. Weight gain and edema only

C. Isolated proteinuria

D. Hyperthyroidism symptoms

3. Cryoglobulinemia typically presents with which cluster of features?

A. Hematuria with red cell casts

B. Isolated peripheral edema

C. Palpable purpura, arthritis, neuropathy, and digital ischemia

D. Hypertension with proteinuria only

4. Which mechanism describes how nifedipine or tamsulosin aid distal ureteral stone passage?

A. Increasing urine calcium excretion

B. Narrowing ureter

C. Inducing ureteral dilatation and relaxation

D. Decreasing peristalsis

5. For a kidney stone that is less than 5 mm, what is the typical outcome?

A. Pass spontaneously

B. Invasive procedure

C. Intravenous fluids

D. Diet changes

SAMPLE

6. How is acute kidney injury defined?

A. Abrupt increase in serum creatinine or a decrease in urine output

B. Gradual decline in eGFR over months

C. Stable creatinine with a normal urine output

D. Chronic proteinuria with preserved creatinine

7. AKI with hydronephrosis on ultrasound is most likely due to obstruction and may be relieved by which intervention?

A. Bladder catheterization

B. High-volume IV fluids

C. Antibiotics

D. Dialysis

8. Isopropanol toxicity presents with which combination?

A. Somnolence/coma with metabolic acidosis

B. Somnolence/coma with normal acid-base

C. Visual disturbances

D. Severe pancreatitis

9. Which test is used to differentiate central from nephrogenic diabetes insipidus?

A. Desmopressin challenge

B. Water deprivation

C. Serum cortisol

D. ACTH stimulation

10. Which medication is commonly associated with hyperkalemia?

A. Cyclosporine

B. ACE inhibitors

C. Furosemide

D. Thiazide

SAMPLE

Answers

SAMPLE

1. B
2. D
3. C
4. C
5. A
6. A
7. A
8. B
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. If SIADH does not respond to fluid restriction in the outpatient setting, which agent is commonly used?

A. Furosemide

B. Demeclocycline

C. Hypertonic saline

D. Conivaptan

In SIADH, too much ADH causes the kidneys to retain water, leading to hyponatremia. After fluid restriction, an outpatient option that directly limits water reabsorption is useful. Demeclocycline works by inducing nephrogenic diabetes insipidus in the collecting ducts, which reduces the kidney's response to ADH and promotes free-water excretion. This helps raise the serum sodium over several days and is suitable for gradual correction in the outpatient setting. By contrast, hypertonic saline is reserved for acute, severe symptoms and usually given in hospital; furosemide can aid water excretion but doesn't address the ADH effect and risks volume depletion; vasopressin receptor antagonists like conivaptan block ADH signaling but are typically IV and require inpatient monitoring. Demeclocycline's oral administration and its mechanism make it a common outpatient choice when fluid restriction alone is not enough.

2. Which clinical features strongly suggest ANCA-associated GN (Wegener's granulomatosis)?

A. Nasal congestion, ulceration, chronic sinusitis, and pulmonary nodules/infiltrates

B. Weight gain and edema only

C. Isolated proteinuria

D. Hyperthyroidism symptoms

ANCA-associated GN, such as granulomatosis with polyangiitis, classically presents with both upper airway disease and lung involvement along with kidney inflammation. The nasal congestion, ulceration, and chronic sinusitis reflect granulomatous inflammation of the upper airway, while nodules or infiltrates in the lungs indicate granulomatous and necrotizing lung disease. Together, these features form a characteristic pattern that strongly points to GPA as the cause of glomerulonephritis. Options describing weight gain with edema, isolated proteinuria, or hyperthyroid symptoms don't capture this distinctive combination of upper airway and pulmonary involvement, so they're not as suggestive.

3. Cryoglobulinemia typically presents with which cluster of features?

A. Hematuria with red cell casts

B. Isolated peripheral edema

C. Palpable purpura, arthritis, neuropathy, and digital ischemia

D. Hypertension with proteinuria only

Cryoglobulinemia is an immune complex–mediated vasculitis that classically affects small vessels, leading to a multi-system pattern. The most characteristic cluster includes skin involvement with palpable purpura from leukocytoclastic vasculitis, joint involvement causing arthralgia or arthritis, nerve involvement causing neuropathy, and vascular compromise in the digits leading to digital ischemia. This combination reflects the typical way cryoglobulinemic vasculitis manifests: immune complex deposition in skin, nerves, and small vessels of the extremities, often in the setting of conditions like Hepatitis C. Other patterns such as isolated hematuria with red cell casts or isolated edema or hypertension with proteinuria alone don't capture this specific vasculitic syndrome, which is why the combination of purpura, arthritis, neuropathy, and digital ischemia is the best fit.

SAMPLE

4. Which mechanism describes how nifedipine or tamsulosin aid distal ureteral stone passage?

A. Increasing urine calcium excretion

B. Narrowing ureter

C. Inducing ureteral dilatation and relaxation

D. Decreasing peristalsis

Relaxing the smooth muscle of the ureter to widen the lumen makes it easier for a distal ureteral stone to pass. Tamsulosin blocks alpha-1 receptors in ureteral smooth muscle, reducing tone in the distal ureter, while nifedipine (a calcium channel blocker) lowers intracellular calcium in smooth muscle, causing further relaxation. The resulting ureteral dilation lowers resistance to flow and helps peristaltic movement push the stone toward the bladder. Increasing urine calcium excretion wouldn't directly aid passage, narrowing the ureter would hinder it, and decreasing peristalsis would slow transit—so the key mechanism is ureteral dilatation and relaxation.

5. For a kidney stone that is less than 5 mm, what is the typical outcome?

A. Pass spontaneously

B. Invasive procedure

C. Intravenous fluids

D. Diet changes

Small kidney stones under five millimeters are likely to pass spontaneously through the urinary tract. Their small size lets them travel down the ureter and be excreted in urine, with many passing within a few weeks. The typical plan is conservative: keep well hydrated, manage pain, and sometimes use medical expulsive therapy to help the stone move along. Invasive procedures are reserved for stones that are larger, cause persistent blockage, or complications. IV fluids are supportive but don't by themselves determine the outcome, and diet changes are mainly for prevention of future stones rather than the immediate passage. So, spontaneous passage is the expected outcome for a stone this small.

6. How is acute kidney injury defined?

A. Abrupt increase in serum creatinine or a decrease in urine output

B. Gradual decline in eGFR over months

C. Stable creatinine with a normal urine output

D. Chronic proteinuria with preserved creatinine

Acute kidney injury is a sudden loss of kidney function, defined by an abrupt rise in serum creatinine or a significant drop in urine output. This captures the acute nature of the problem, which can develop over hours to days and requires urgent assessment and management. The other options describe chronic or stable states: a gradual decline in eGFR over months points to chronic kidney disease, stable creatinine with normal urine output suggests no kidney injury, and chronic proteinuria with preserved creatinine indicates a chronic nephropathy rather than an acute event.

SAMPLE

7. AKI with hydronephrosis on ultrasound is most likely due to obstruction and may be relieved by which intervention?

A. Bladder catheterization

B. High-volume IV fluids

C. Antibiotics

D. Dialysis

The key idea is that hydronephrosis on ultrasound signals obstruction causing post-renal AKI. When urine flow is blocked, pressure backs up into the kidneys, impairing glomerular filtration. Relieving the blockage promptly can restore urine drainage and allow renal function to recover. Bladder catheterization directly decompresses the bladder, reducing the outlet pressure and back-pressure on the kidneys. This quick intervention addresses the obstruction, which is the root cause of the AKI in this scenario, making it the most effective initial treatment. High-volume IV fluids won't fix an obstruction and may worsen issues if the kidneys aren't able to drain urine. Antibiotics treat infection, not the obstruction. Dialysis is a supportive measure for severe, uncompensated kidney failure or specific metabolic derangements and doesn't resolve the underlying obstruction.

8. Isopropanol toxicity presents with which combination?

A. Somnolence/coma with metabolic acidosis

B. Somnolence/coma with normal acid-base

C. Visual disturbances

D. Severe pancreatitis

The main concept is how isopropanol affects the body and its acid-base profile. Isopropanol causes central nervous system depression, leading to somnolence or coma. Unlike many other alcohols, its major metabolite, acetone, is not acidic, so it does not produce a significant metabolic acidosis. Patients often have an elevated osmolal gap, but the acid-base status remains normal. Visual disturbances point more toward methanol or ethylene glycol toxicity, and pancreatitis is not a typical feature of isopropanol poisoning. Therefore, the presentation that best fits is somnolence or coma with a normal acid-base status.

9. Which test is used to differentiate central from nephrogenic diabetes insipidus?

A. Desmopressin challenge

B. Water deprivation

C. Serum cortisol

D. ACTH stimulation

Differentiating central from nephrogenic diabetes insipidus relies on assessing the kidney's response to ADH replacement. In central DI there is an ADH deficiency, so giving desmopressin, a synthetic vasopressin analogue, should markedly raise urine osmolality and reduce urine volume as the kidneys reabsorb more water. In nephrogenic DI the kidneys are unresponsive to ADH due to receptor or signaling defects, so there is little or no change in urine osmolality or volume after desmopressin. The water deprivation test helps confirm DI by showing an inability to concentrate urine, but it doesn't distinguish the cause as specifically as the desmopressin challenge. Serum cortisol and ACTH stimulation test adrenal function and are not used to differentiate DI types.

SAMPLE

10. Which medication is commonly associated with hyperkalemia?

A. Cyclosporine

B. ACE inhibitors

C. Furosemide

D. Thiazide

The main idea here is how certain drugs disturb kidney potassium handling to raise serum potassium. Cyclosporine, a calcineurin inhibitor often used in transplant patients, is well known for causing hyperkalemia. It narrows the afferent arteriole, which lowers renal blood flow and glomerular filtration rate. With a lower GFR, less potassium is filtered and excreted, and the distal nephron's ability to secrete potassium can also be impaired. Additionally, there may be a tubular dysfunction and blunted aldosterone response, further reducing potassium disposal. In contrast, diuretics like furosemide and thiazides promote potassium loss and typically cause hypokalemia, not hyperkalemia. ACE inhibitors can cause hyperkalemia by reducing aldosterone-driven potassium excretion, but in this context cyclosporine is the medication most classically linked to hyperkalemia due to its nephrotoxic effects on renal handling of potassium.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://itenephrology.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE