JBL OBSTETRICS AND PEDIATRICS PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://jblobstetricspeds.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the common position of the baby's head upon delivery?

A. With the face up

B. Posteriorly, face down

C. Posteriorly, to one side

D. Anteriorly, with the chin up

2. What growth chart percentile indicates underweight in children?

A. Below the 10th percentile

B. Below the 5th percentile

C. Between the 5th and 15th percentile

D. Above the 25th percentile

3. How often should well-child visits occur during the first two years of life?

A. Quarterly

B. At 1 month, 2 months, 4 months, 6 months, 9 months, and 12 months

C. Bi-annually

D. Only during the first year

4. In assessing an infant, what is an effective way to calm and comfort during an examination?

A. Allow the infant to play with medical equipment

B. Perform the examination quickly while holding the child

C. Involve the parent in holding the infant

D. Keep the infant swaddled and restrict movement

5. At what age do most children achieve bladder control during the day?

A. By age 1

B. By age 2

C. By 3-5 years of age

D. By age 6

SAMPLE

6. What should you do for a 4-month-old male who suddenly became limp, cyanotic, and apneic but is now breathing adequately?

A. Administer oxygen via nonrebreathing mask

B. Allow a parent to transport him to the hospital

C. Contact the infant's pediatrician to seek advice

D. Transport to the hospital and monitor him en route

7. What are the signs of dehydration in infants?

A. Excessive crying and fever

B. Dry mouth, lack of tears, and decreased urine output

C. Increased appetite and weight gain

D. Cool skin and low energy

8. What dietary recommendation is advised for toddlers to prevent choking?

A. Introduce whole nuts at age 3

B. Avoid whole nuts and hard candies

C. Encourage soft fruits and vegetables

D. Limit protein intake

9. What is the most common congenital heart defect in children?

A. Atrial septal defect (ASD)

B. Ventricular septal defect (VSD)

C. Patent ductus arteriosus (PDA)

D. Coarctation of the aorta

10. At what stage do children typically begin to develop stranger anxiety?

A. Infant.

B. Neonate.

C. Toddler.

D. Preschooler.

SAMPLE

Answers

SAMPLE

1. C
2. B
3. B
4. C
5. C
6. D
7. B
8. B
9. B
10. C

SAMPLE

Explanations

SAMPLE

1. What is the common position of the baby's head upon delivery?

A. With the face up

B. Posteriorly, face down

C. Posteriorly, to one side

D. Anteriorly, with the chin up

The common position of the baby's head upon delivery is typically with the occiput (the back part of the head) oriented towards the mother's pubic bone. This position often leads to what is described as the "vertex presentation," which occurs when the baby is positioned to exit the birth canal in a way that allows the smallest part of the head to present first. In the most favorable delivery scenario, the baby's head is often turned slightly to the side, which helps navigate the shoulders and body through the birth canal. This position, where the back of the head is facing one of the mother's sides, often results in a more manageable delivery process by enabling the head to rotate as it descends. Understanding the dynamics of these positions is crucial for practitioners as they can influence labor and delivery outcomes significantly. The other options may represent less favorable positions that could complicate delivery or require interventions like manual rotation or the use of forceps.

2. What growth chart percentile indicates underweight in children?

A. Below the 10th percentile

B. Below the 5th percentile

C. Between the 5th and 15th percentile

D. Above the 25th percentile

To determine whether a child is underweight, healthcare providers typically reference growth charts that categorize a child's weight relative to their peers. A child whose weight falls below the 5th percentile is considered to be underweight. This means that less than 5% of children of the same age and sex weigh less than this specific child. Being in this percentile indicates a significant deviation from what is considered healthy growth patterns and raises concerns about possible underlying health issues, nutritional deficiencies, or growth disorders. In clinical practice, identifying underweight status is crucial because it prompts further evaluation and intervention to ensure the child's health and development are supported appropriately. The other options do not accurately represent the clinical threshold for underweight. For example, the 10th percentile suggests a child is on the lower end of weight but not necessarily classified as underweight. The range between the 5th and 15th percentile indicates that while some children may be lighter, they do not meet the criteria for underweight. Finally, being above the 25th percentile signifies that a child is within or above the average weight range compared to peers, thus not falling into the underweight category.

SAMPLE

3. How often should well-child visits occur during the first two years of life?

A. Quarterly

B. At 1 month, 2 months, 4 months, 6 months, 9 months, and 12 months

C. Bi-annually

D. Only during the first year

The recommendation for well-child visits during the first two years of life emphasizes the importance of regular monitoring and assessment of a child's growth and development. The specific schedule of visits at 1 month, 2 months, 4 months, 6 months, 9 months, and 12 months allows for timely vaccinations, developmental screenings, and parental guidance on nutrition, safety, and other critical aspects of childcare. This frequency is crucial because early childhood is a period of rapid growth and development, and these visits provide an opportunity to monitor milestones, address any medical concerns, and provide necessary immunizations. Each visit is tailored to track developmental progress and identify potential issues early, thus ensuring that children are healthy and thriving. In contrast, the other options suggest less frequent visits or omit important ages. Quarterly visits might seem reasonable, but they don't align with the established guidelines for regular checks during these formative months. Bi-annual visits do not provide the necessary frequency to monitor growth and development during critical periods. Visits only during the first year would completely overlook key developmental stages in the second year of life when additional assessments and vaccinations are necessary.

4. In assessing an infant, what is an effective way to calm and comfort during an examination?

A. Allow the infant to play with medical equipment

B. Perform the examination quickly while holding the child

C. Involve the parent in holding the infant

D. Keep the infant swaddled and restrict movement

Involving the parent in holding the infant during an examination is particularly effective for calming and comforting the child. Infants often feel safer and more secure when they are close to their primary caregiver. The presence of the parent can provide emotional support, reduce anxiety, and help the infant associate the medical environment with a sense of safety, which can facilitate a smoother examination process. Additionally, parents can engage with the infant through soothing sounds or gentle motions while the healthcare provider conducts the examination, further aiding in comfort and reducing distress. Infants may feel overwhelmed in an unfamiliar environment, so having their parent close by can help bridge that gap, making the process less intimidating. It fosters trust not just in the parent but also in the healthcare provider as they gently and effectively perform their duties. This approach aligns with pediatric practices that emphasize the importance of family-centered care.

SAMPLE

5. At what age do most children achieve bladder control during the day?

A. By age 1

B. By age 2

C. By 3-5 years of age

D. By age 6

Most children achieve bladder control during the day between the ages of 3 to 5 years. By this age range, many children have developed the necessary physical and neurological maturity to recognize the urge to urinate and to control their bladder effectively. This developmental milestone is part of toilet training, which generally begins as children approach their third birthday. During this period, they typically gain confidence in their ability to communicate their needs and have increased bladder capacity, allowing them to stay dry for longer periods. Younger children, such as those around the age of 1 or 2, are still developing their bodily awareness and coordination, making it challenging for them to consistently control their bladder. As for children aged 6, while they might be solidly trained, the majority have already achieved daytime bladder control by the earlier age range of 3 to 5 years. Therefore, recognizing that bladder control is a part of broader developmental milestones helps frame the timeline accurately.

6. What should you do for a 4-month-old male who suddenly became limp, cyanotic, and apneic but is now breathing adequately?

A. Administer oxygen via nonrebreathing mask

B. Allow a parent to transport him to the hospital

C. Contact the infant's pediatrician to seek advice

D. Transport to the hospital and monitor him en route

Transporting the infant to the hospital and monitoring him en route is the appropriate action in this scenario. The infant's initial presentation of becoming limp, cyanotic, and apneic suggests a potentially serious condition that could require immediate medical intervention. Although he is breathing adequately now, this episode indicates that he could be at risk for further complications. In situations with young children, prompt medical evaluation is critical, especially when they exhibit signs of distress or changes in their condition. The fact that the infant previously experienced respiratory distress underscores the importance of continuous monitoring during transport. This allows for immediate response if his condition deteriorates again. Other options may not provide the necessary level of care and intervention. Administering oxygen via a nonrebreathing mask might seem appropriate, but the focus should remain on ensuring the infant is under professional medical care as soon as possible. Allowing a parent to transport the child could result in delays in receiving urgent care. While contacting the pediatrician could provide valuable advice, it does not replace the need for immediate assessment and potential emergency care at a facility. Therefore, transporting to the hospital ensures that the child receives the necessary medical attention and monitoring without delay.

SAMPLE

7. What are the signs of dehydration in infants?

A. Excessive crying and fever

B. Dry mouth, lack of tears, and decreased urine output

C. Increased appetite and weight gain

D. Cool skin and low energy

The signs of dehydration in infants are critical to recognize, as infants can become dehydrated quickly, which poses serious health risks. Among the options presented, one of the key indicators is the presence of a dry mouth, a lack of tears, and decreased urine output. A dry mouth suggests that the infant is not receiving enough fluids, and without the production of tears when crying, it is an additional sign that the body is conserving water. Decreased urine output is particularly telling; healthy infants typically have a certain amount of wet diapers throughout the day, and a noticeable reduction can indicate a deficiency in hydration levels. While excessive crying and fever can be associated with illness or discomfort, they are not definitive signs of dehydration. Similarly, increased appetite and weight gain would typically suggest that the infant is receiving adequate nourishment and hydration rather than being dehydrated. Cool skin and low energy can also signal various health issues, but they are not as specific as the signs listed in the correct choice for indicating dehydration. Understanding these signs helps caregivers and healthcare providers respond promptly to infants at risk of dehydration, ensuring their safety and well-being.

8. What dietary recommendation is advised for toddlers to prevent choking?

A. Introduce whole nuts at age 3

B. Avoid whole nuts and hard candies

C. Encourage soft fruits and vegetables

D. Limit protein intake

The recommendation to avoid whole nuts and hard candies is crucial for preventing choking in toddlers. Choking hazards are a significant concern for this age group due to their developing chewing and swallowing skills. Whole nuts are particularly problematic, as their small, hard, and unyielding nature can easily block a toddler's airway, especially if the child does not have the ability to adequately chew them. Similarly, hard candies can dissolve in the mouth but pose a risk if they are bitten into, leading to potential choking. The avoidance of these foods aligns with guidelines provided by pediatric health organizations, which emphasize ensuring that foods offered to young children are age-appropriate in size and texture. The focus should be on foods that are soft, easily chewed, and can be safely swallowed. This is why the correct recommendation highlights the need to steer clear of both whole nuts and hard candies to mitigate the risk of choking accidents in toddlers.

9. What is the most common congenital heart defect in children?

A. Atrial septal defect (ASD)

B. Ventricular septal defect (VSD)

C. Patent ductus arteriosus (PDA)

D. Coarctation of the aorta

Ventricular septal defect (VSD) is recognized as the most common congenital heart defect in children. It involves a defect in the ventricular septum, which is the wall that separates the left and right ventricles of the heart. This condition allows for the mixing of oxygenated and deoxygenated blood, leading to increased blood flow to the lungs and potential complications if left untreated. VSDs can vary in size and severity, with smaller defects sometimes closing spontaneously over time. However, larger defects can lead to heart failure, pulmonary hypertension, and other serious issues, necessitating medical surveillance or surgical intervention. The high prevalence of VSDs in the pediatric population contributes to their categorization as the most common congenital heart defect, which is critical for healthcare professionals to understand when diagnosing and managing these conditions.

SAMPLE

10. At what stage do children typically begin to develop stranger anxiety?

A. Infant.

B. Neonate.

C. Toddler.

D. Preschooler.

Stranger anxiety is a psychological phenomenon commonly observed in infants and young children as they develop their social and emotional skills. This anxiety typically emerges in response to the presence of unfamiliar individuals, indicating that the child is becoming more aware of social boundaries and attachments. The correct answer reflects that stranger anxiety primarily begins in the infant stage, particularly around 6 to 8 months of age. During this time, infants start forming strong attachments to their primary caregivers and may react with distress or fear when confronted with strangers. This behavior is a normal part of development as children learn to differentiate between familiar caregivers and unfamiliar individuals, marking a critical stage in their emotional growth. This understanding is essential because it highlights the normal developmental milestones children experience. While toddlers may continue to exhibit signs of apprehension towards strangers, the initial onset typically occurs during infancy, thus aligning with the correct answer regarding the development of stranger anxiety.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://jblobstetricspeds.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE