JRCALC RESUSCITATION PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://jrcalcresuscitation.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What does JRCALC stand for?

A. Joint Royal Colleges Ambulance Liaison Committee

B. Joint Rescue Committee for Ambulance Liaison and Care

C. Joint Response Council for Ambulance and Life Support

D. Joint Regulation for Cardiac Ambulance Liaison Consulting

2. What is a potential risk of excessive compression depth during CPR?

A. Increased cardiac output

B. Increased risk of organ damage

C. Decrease in airway patency

D. Faster heart rhythm stabilization

3. How many initial rescue breaths are recommended for a drowned pediatric patient?

A. 3

B. 5

C. 7

D. 10

4. What is the recommended action for a child with a suspected airway obstruction?

A. Perform abdominal thrusts only

B. Administer back blows and chest thrusts

C. Encourage the child to cough forcefully

D. Provide oxygen immediately

5. When should high-flow oxygen be administered?

A. For patients with suspected hypoxemia or respiratory distress

B. Only for patients who are unconscious

C. When a patient's oxygen saturation levels are normal

D. For patients with chest pain regardless of symptoms

SAMPLE

6. What immediate action should be taken when a patient is unresponsive and not breathing?

A. Call for emergency help and start CPR

B. Wait for the patient to regain consciousness

C. Administer oxygen if available

D. Position the patient in the recovery position

7. What are the phases of ventricular fibrillation?

A. Electrical phase, Metabolic phase, Mechanical phase

B. Electrical phase, Circulatory phase, Metabolic phase

C. Electrical phase, Neurogenic phase, Metabolic phase

D. Electrical phase, Circulatory phase, Respiratory phase

8. What should be done first in the event of a suspected cardiac arrest?

A. Start CPR

B. Call for emergency help

C. Check the patient's pulse

D. Use an AED

9. What should be done if an AED prompts a "no shock advised" message?

A. Continue CPR until advanced help arrives

B. Disconnect the AED and wait

C. Re-analyze the rhythm again after one minute

D. Administer medication immediately

10. What should be the first step when encountering an unresponsive adult?

A. Call for help

B. Check for breathing

C. Perform CPR

D. Check for a pulse

SAMPLE

Answers

SAMPLE

1. A
2. B
3. B
4. B
5. A
6. A
7. B
8. B
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. What does JRCALC stand for?

A. Joint Royal Colleges Ambulance Liaison Committee

B. Joint Rescue Committee for Ambulance Liaison and Care

C. Joint Response Council for Ambulance and Life Support

D. Joint Regulation for Cardiac Ambulance Liaison Consulting

JRCALC stands for Joint Royal Colleges Ambulance Liaison Committee. This organization is essential in developing clinical guidelines and protocols for ambulance services in the UK, focusing on providing high-quality patient care. The involvement of the "Royal Colleges" signifies collaboration among various professional bodies within the medical field, ensuring that the guidelines are evidence-based and reflect best practices in emergency medical care. This committee emphasizes the importance of standardizing care across different ambulance services, which ultimately improves patient outcomes in emergency situations. Other options do not accurately reflect the established name or mission of the organization, as they contain variations and terms that do not correspond to the official terminology recognized in clinical practice within the ambulance services.

2. What is a potential risk of excessive compression depth during CPR?

A. Increased cardiac output

B. Increased risk of organ damage

C. Decrease in airway patency

D. Faster heart rhythm stabilization

The chosen answer, which highlights the increased risk of organ damage, is foundational to understanding the mechanics of CPR and the physiological effects exerted on the body during resuscitation efforts. During cardiopulmonary resuscitation (CPR), compressing the chest too deeply or with excessive force can lead to significant internal trauma. Organs such as the lungs, liver, and spleen are particularly susceptible to injury from high-pressure compressions. For instance, deep compressions might cause rib fractures, which can puncture the lungs or lead to internal bleeding, especially in patients who might already have weakened bone structures or pre-existing health issues. This understanding emphasizes the importance of maintaining the recommended compression depth during CPR, which is 5 to 6 centimeters in adults, rather than exceeding it in an attempt to enhance circulation. While effective compressions are critical for maintaining blood flow to vital organs, it is essential to balance the force applied to prevent additional harm.

3. How many initial rescue breaths are recommended for a drowned pediatric patient?

A. 3

B. 5

C. 7

D. 10

For a drowned pediatric patient, the recommended practice is to administer 5 initial rescue breaths. This approach is fundamental because it aims to provide oxygen to the lungs before starting chest compressions, particularly in drowning cases where the primary issue is often the lack of oxygen due to water inhalation. Administering these initial breaths helps to ventilate the lungs and establish a minimum level of oxygenation in the body, which is critical in restoring circulation and addressing the hypoxic state of the patient. These breaths should be delivered effectively and at a rate that ensures that the chest rises visibly, indicating that the air is entering the lungs. This practice is reflective of current guidelines aimed at optimizing the chances of survival and recovery in pediatric patients who have experienced drowning or near-drowning situations. It balances the need for immediate intervention with the specifics of how respiratory distress from drowning should be addressed.

SAMPLE

4. What is the recommended action for a child with a suspected airway obstruction?

A. Perform abdominal thrusts only

B. Administer back blows and chest thrusts

C. Encourage the child to cough forcefully

D. Provide oxygen immediately

For a child with a suspected airway obstruction, administering back blows and chest thrusts is the recommended action. This approach is effective because back blows help to create a forceful impact on the back, which can aid in dislodging the obstruction in the airway, while chest thrusts can further help to expel the object if back blows alone are insufficient. This technique is particularly suitable for children who are over the age of one because it leverages their anatomy and physiology to assist in clearing the airway. Encouraging the child to cough forcefully is typically a good initial response for conscious individuals with mild obstruction, as it can help to expel the object. However, if the obstruction is severe and the child is unable to breathe, cough, or speak, then the combination of back blows and chest thrusts is the recommended response rather than simply encouraging coughing. Administering oxygen immediately is not appropriate in this scenario because oxygen will not resolve the obstruction; the priority must be to clear the airway. Abdominal thrusts are more suited for adults or larger children, and are less effective for infants and younger children, making them not the best option in this case.

5. When should high-flow oxygen be administered?

A. For patients with suspected hypoxemia or respiratory distress

B. Only for patients who are unconscious

C. When a patient's oxygen saturation levels are normal

D. For patients with chest pain regardless of symptoms

High-flow oxygen should be administered to patients with suspected hypoxemia or respiratory distress because these conditions indicate that a patient may not be receiving adequate oxygenation. Hypoxemia refers to low levels of oxygen in the blood, which can lead to serious complications if not treated promptly. Patients in respiratory distress may also struggle to breathe effectively, further exacerbating the risk of low oxygen levels. Providing high-flow oxygen can help restore adequate oxygen saturation levels and improve overall respiratory function, facilitating better outcomes. The other choices do not meet the criteria for when high-flow oxygen is appropriately indicated. Administering high-flow oxygen solely to unconscious patients neglects the broader scope of individuals who may need intervention, such as those conscious but experiencing hypoxemia. Administering oxygen to patients with normal oxygen saturation levels is unnecessary and could even lead to complications such as oxygen toxicity in certain situations. Providing high-flow oxygen just for patients with chest pain does not take into account the need for clinical assessment of oxygen levels or respiratory distress, as chest pain can arise from various underlying conditions that may not require supplemental oxygen.

SAMPLE

6. What immediate action should be taken when a patient is unresponsive and not breathing?

A. Call for emergency help and start CPR

B. Wait for the patient to regain consciousness

C. Administer oxygen if available

D. Position the patient in the recovery position

When a patient is unresponsive and not breathing, the immediate action of calling for emergency help and starting CPR is critical. This response is vital because time is of the essence in cardiorespiratory emergencies. The lack of responsiveness and absence of breathing indicate that the patient is likely in cardiac arrest or experiencing a life-threatening situation where circulation and respiration have ceased. Initiating CPR immediately helps to maintain blood flow to vital organs, especially the brain and heart, which can sustain life long enough for emergency services to arrive. CPR compressions help to artificially circulate blood and can significantly increase the chances of survival when performed promptly. This immediate action addresses both the unresponsiveness and failure to breathe, while waiting or administering oxygen do not provide the necessary life-saving intervention that CPR offers in this critical moment.

7. What are the phases of ventricular fibrillation?

A. Electrical phase, Metabolic phase, Mechanical phase

B. Electrical phase, Circulatory phase, Metabolic phase

C. Electrical phase, Neurogenic phase, Metabolic phase

D. Electrical phase, Circulatory phase, Respiratory phase

The correct answer highlights the phases of ventricular fibrillation as the electrical phase, circulatory phase, and metabolic phase. In a cardiac arrest scenario involving ventricular fibrillation, the electrical phase occurs immediately, characterized by the disorganized electrical activity in the heart that prevents effective pumping. This phase lasts for a few minutes, during which defibrillation is most likely to be successful. Following this, the circulatory phase begins after a few minutes. In this phase, the absence of effective blood flow leads to increased cellular death and ischemia. It's during this time that the chances of successful resuscitation decline if definitive measures such as defibrillation are not taken. The metabolic phase follows, which can begin around 10 minutes after the onset of cardiac arrest. During this phase, the body's tissues are in a state of acidosis and require extensive time and intervention to restore function, given the severe metabolic derangements that occur. Together, these phases underscore the sequence of physiological changes that occur during ventricular fibrillation, emphasizing the urgent need for prompt and effective resuscitative measures at each stage. This knowledge is essential for providing timely and appropriate care during cardiac emergencies.

SAMPLE

8. What should be done first in the event of a suspected cardiac arrest?

A. Start CPR

B. Call for emergency help

C. Check the patient's pulse

D. Use an AED

The first action that should be taken in the event of a suspected cardiac arrest is to call for emergency help. This step is crucial because it ensures that advanced medical support is on the way while other life-saving measures, such as CPR, can be initiated. Rapid response from emergency services can significantly increase the chances of survival for the patient experiencing a cardiac arrest. Calling for help activates the emergency response system, allowing for more skilled intervention, including advanced airway management and medication administration, that may be necessary once the first responders arrive. Immediate recognition of cardiac arrest and alerting emergency services provides the foundation for a comprehensive and effective response, benefiting both the patient and the individuals performing CPR. While initiating CPR is paramount, it is recommended to first ensure that help is called in order to coordinate the best chances of survival. Checking the patient's pulse or using an AED can come after these critical initial steps, but without calling for emergency help, the patient might not receive the necessary advanced care in a timely manner.

9. What should be done if an AED prompts a "no shock advised" message?

A. Continue CPR until advanced help arrives

B. Disconnect the AED and wait

C. Re-analyze the rhythm again after one minute

D. Administer medication immediately

When an AED indicates "no shock advised," it means that the device has analyzed the heart rhythm and determined that it is not in a shockable state, such as ventricular fibrillation or pulseless VT. Continuing CPR is essential because it helps maintain blood circulation and oxygenation to vital organs, particularly the brain. High-quality CPR enhances the chances of successful defibrillation if the rhythm changes back to a shockable state later on. During CPR, the rescuer should ensure that chest compressions are uninterrupted and adequate in depth and rate, providing ventilations if trained to do so. Waiting for advanced help without continuing CPR would not provide any meaningful support for the cardiac arrest victim. Therefore, continuing CPR until advanced help arrives is the appropriate action to take following an AED's "no shock advised" message.

10. What should be the first step when encountering an unresponsive adult?

A. Call for help

B. Check for breathing

C. Perform CPR

D. Check for a pulse

When encountering an unresponsive adult, the initial step is to call for help. This is crucial because it ensures that emergency services are notified while you're assessing the situation and providing care. Calling for help not only brings additional resources to the scene but also allows the responder to focus on potentially life-saving interventions without delay. In many cases, once assistance is summoned, other responders can take over or provide support, allowing for a more effective and safe response. Following this initial step, checking for breathing, performing CPR, or checking for a pulse are important actions that would typically follow. However, without first ensuring that trained assistance is on the way, your ability to provide effective resuscitation measures could be compromised due to a lack of support or necessary equipment.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://jrcalcresuscitation.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE