KETTERING PATIENT ASSESSMENT PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ketteringpatientassmt.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which is considered more specific for kidney failure, creatinine or BUN?

A. BUN

B. Both are equally specific

C. Creatinine

D. Neither are specific

2. When is intubation indicated for stridor?

A. For severe swelling or epiglottitis

B. For anxiety-related symptoms

C. For mild airway obstruction

D. After a heart attack

3. What class of drugs should a therapist administer to a patient in case of an allergic reaction causing vasodilation and decreased blood pressure?

A. Vasodilators

B. Diuretics

C. Antihistamines

D. Vasoconstrictors

4. Which of the following should be minimized to reduce central venous pressure?

A. Rest

B. Exercise

C. Straining, retching, and coughing

D. Medication

5. A patient presents to the ER with shortness of breath and JVD during exhalation. What underlying problem should the therapist suspect?

A. Asthma

B. Obstructive lung disease

C. Interstitial lung disease

D. Pneumonia

SAMPLE

6. Fine crackles or moist crepitant rales are typically associated with which condition?

A. Pneumonia

B. CHF/pulmonary edema

C. Atelectasis

D. Chronic bronchitis

7. Which of the following conditions indicates muscle atrophy?

A. Cachexia

B. Hypertrophy

C. Fever

D. Pain

8. What intervention is suggested for a patient exhibiting signs of hypervolemia?

A. Fluid therapy

B. Diuretics

C. Increased dietary salt

D. Fluid restriction

9. What is a brand name for a chronotropic drug used to increase heart rate?

A. Atenolol

B. Atropine

C. Labetalol

D. Digoxin

10. How is QT calculated when given stroke volume?

A. Heart Rate x Stroke Volume

B. Stroke Volume / Heart Rate

C. Heart Rate + Stroke Volume

D. Stroke Volume - Heart Rate

SAMPLE

Answers

SAMPLE

1. C
2. A
3. D
4. C
5. B
6. B
7. A
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. Which is considered more specific for kidney failure, creatinine or BUN?

A. BUN

B. Both are equally specific

C. Creatinine

D. Neither are specific

Creatinine is considered more specific for kidney failure than blood urea nitrogen (BUN) due to its clearer association with renal function. Creatinine is a byproduct of muscle metabolism and is produced at a relatively constant rate. It is filtered out of the blood by the kidneys, so an increase in creatinine levels directly indicates impaired kidney function. This makes it a reliable marker for kidney health. BUN, on the other hand, can be influenced by various factors other than kidney function, such as hydration status, dietary protein intake, liver function, and gastrointestinal bleeding. Therefore, while BUN can provide useful information regarding overall health and may indicate kidney involvement in some cases, it is less specific than creatinine for diagnosing kidney failure. In summary, creatinine is favored as a more specific indicator of kidney function because it reflects the kidneys' ability to filter waste products from the blood, with minimal interference from non-renal factors.

2. When is intubation indicated for stridor?

A. For severe swelling or epiglottitis

B. For anxiety-related symptoms

C. For mild airway obstruction

D. After a heart attack

Intubation is indicated for stridor in situations where there is significant airway compromise, such as severe swelling or epiglottitis. In cases of severe swelling, the airway can become dangerously obstructed, and immediate intervention is required to secure the airway and ensure proper oxygenation. Epiglottitis, which is inflammation of the epiglottis often due to infection, can lead to rapid deterioration and complete airway occlusion, making intubation a critical step in management to prevent respiratory failure. In contrast, anxiety-related symptoms may present with stridor due to hyperventilation or muscle tension but generally do not require intubation unless there are underlying respiratory issues. Mild airway obstruction typically does not necessitate intubation as there may be alternative management options available, such as steroids or nebulized treatments. As for a heart attack, it is unrelated to stridor and would not be an appropriate indication for intubation in that context.

SAMPLE

3. What class of drugs should a therapist administer to a patient in case of an allergic reaction causing vasodilation and decreased blood pressure?

A. Vasodilators

B. Diuretics

C. Antihistamines

D. Vasoconstrictors

In the context of an allergic reaction causing vasodilation and decreased blood pressure, the appropriate class of drugs to administer would be vasoconstrictors. Allergic reactions can lead to significant vasodilation due to the release of histamine and other mediators, which can result in hypotension. Vasoconstrictors work by constricting blood vessels, which increases peripheral vascular resistance and ultimately helps to raise blood pressure. This is crucial in cases where the patient is experiencing symptoms of shock or severe hypotension as a result of an allergic reaction. While vasodilators would further exacerbate the situation by widening blood vessels, and diuretics do not address the urgent need for blood pressure stabilization, antihistamines primarily function to block the effects of histamine and alleviate allergic symptoms but do not directly counteract severe hypotension due to vasodilation. Therefore, administering a vasoconstrictor is the most effective and appropriate therapeutic response in this scenario.

4. Which of the following should be minimized to reduce central venous pressure?

A. Rest

B. Exercise

C. Straining, retching, and coughing

D. Medication

To reduce central venous pressure (CVP), minimizing straining, retching, and coughing is vital because these activities increase intrathoracic pressure, which directly affects venous return to the heart. When you strain or cough, the pressure in the chest cavity rises, compressing the veins and impeding blood flow back to the heart. This action can lead to an increase in CVP as the volume of blood returning to the heart is momentarily reduced due to the elevated pressure in the thoracic area. By minimizing these activities, you help maintain a lower thoracic pressure, thereby allowing for more efficient venous return and, ultimately, a reduction in central venous pressure. This distinction is crucial for patients who may already be experiencing conditions that elevate CVP, such as heart failure or fluid overload situations. A better management of CVP can lead to improved cardiac output and overall patient stability. The other options like rest, exercise, and medication can have varied effects on CVP but do not specifically address the immediate mechanics of how straining and coughing directly impact thoracic pressure and venous return.

SAMPLE

5. A patient presents to the ER with shortness of breath and JVD during exhalation. What underlying problem should the therapist suspect?

A. Asthma

B. Obstructive lung disease

C. Interstitial lung disease

D. Pneumonia

In this case, the presence of shortness of breath and jugular venous distention (JVD) during exhalation suggests a problem primarily related to restricted airflow and potential right heart failure or volume overload. These symptoms are often indicative of obstructive lung disease, as this condition can lead to increased pressure in the thoracic cavity during exhalation, which then affects venous return and causes JVD. Obstructive lung diseases, such as chronic obstructive pulmonary disease (COPD) and asthma, involve narrowing of the airways that leads to difficulty in exhaling air, resulting in trapping of air in the lungs. This air trapping increases lung volumes, which can impact the cardiovascular system and lead to complications such as JVD. In situations where the lungs are hyperinflated, the pressure changes can significantly affect venous return from the right side of the heart, causing the observed jugular venous distention. Other options, while related to respiratory issues, do not typically present with both shortness of breath and JVD during exhalation. Asthma, for instance, primarily involves wheezing and airway constriction, but JVD is not a common feature. Interstitial lung disease often presents with dry cough and progressive short

6. Fine crackles or moist crepitant rales are typically associated with which condition?

A. Pneumonia

B. CHF/pulmonary edema

C. Atelectasis

D. Chronic bronchitis

Fine crackles, also known as moist crepitant rales, are often associated with conditions where fluid accumulation in the alveoli is present. In the context of congestive heart failure (CHF) or pulmonary edema, the lungs can become congested with fluid due to excessive pressure in the pulmonary circulation. This fluid may cause fine crackles as air bubbles move through the fluid-filled small airways during inspiration, resulting in the characteristic sounds heard upon auscultation. In CHF, the heart's ineffective pumping leads to a backlog of fluid in the lungs, underlying the development of fine crackles. Similarly, pulmonary edema—whether due to CHF or other causes—produces the same clinical findings as fluid infiltrates the lung tissue and alveolar spaces, causing this specific type of abnormal lung sound. Other conditions listed typically do not present with fine crackles or they have different auscultatory findings associated with them. For instance, pneumonia may produce a variety of sounds, including coarse crackles, but it is the presence of inflammation and consolidation rather than fluid alone that causes those sounds. Atelectasis can lead to diminished breath sounds rather than fine crackles, while chronic bronchitis usually results in wheezes and rhonchi rather than crackles.

SAMPLE

7. Which of the following conditions indicates muscle atrophy?

A. Cachexia

B. Hypertrophy

C. Fever

D. Pain

Muscle atrophy refers to the wasting or loss of muscle tissue, which can occur due to various underlying conditions. Cachexia is a syndrome characterized by significant weight loss, muscle wasting, and a decrease in overall physical condition, often seen in chronic illnesses such as cancer, AIDS, or heart failure. The presence of cachexia specifically indicates that the body is not only losing fat but also muscle mass, leading to atrophy. In contrast, hypertrophy refers to the increase in muscle mass and size, usually as a result of exercise or increased workload. Fever is a temporary increase in body temperature often due to infection and does not inherently cause muscle atrophy. Pain may limit movement and activity but does not directly indicate muscle atrophy either. Thus, the correct answer connects directly to a condition known for its association with muscle loss, making cachexia the condition that indicates muscle atrophy.

8. What intervention is suggested for a patient exhibiting signs of hypervolemia?

A. Fluid therapy

B. Diuretics

C. Increased dietary salt

D. Fluid restriction

When managing a patient exhibiting signs of hypervolemia, diuretics are often the suggested intervention. Hypervolemia refers to an excess of fluid in the body, which can lead to complications such as heart failure, pulmonary edema, or hypertension. Diuretics help increase urine output, promoting the excretion of excess fluid and thereby reducing overall body volume. This approach is particularly effective because it directly addresses the underlying issue of fluid overload. By eliminating surplus fluid through the kidneys, diuretics can alleviate symptoms associated with hypervolemia, improving the patient's comfort and safety. Other interventions, such as fluid therapy or increased dietary salt, would exacerbate the issue, contributing to higher volumes of fluid in the bloodstream. Fluid restriction might be a consideration in certain cases, but diuretics generally provide a more immediate and effective means of addressing hypervolemia and its associated complications.

9. What is a brand name for a chronotropic drug used to increase heart rate?

A. Atenolol

B. Atropine

C. Labetalol

D. Digoxin

Atropine is a well-known chronotropic drug that increases heart rate by blocking the action of the vagus nerve on the heart. It works primarily by inhibiting the parasympathetic nervous system's influence on the heart, which normally slows the heart rate. In situations such as symptomatic bradycardia, atropine is commonly administered to help restore a normal heart rate by increasing conduction through the AV node and enhancing the overall heart rate response. In contrast, the other drugs listed in the choices have different primary effects. Atenolol is a beta-blocker that decreases heart rate and contractility, Labetalol is primarily used as an antihypertensive and also decreases heart rate, and Digoxin, while it can increase heart contractility, primarily has a role in controlling heart rate in atrial fibrillation rather than directly acting as a chronotropic agent to increase heart rate. Thus, the unique action of atropine aligns it directly with the concept of increasing heart rate, making it the correct answer in this scenario.

SAMPLE

10. How is QT calculated when given stroke volume?

A. Heart Rate x Stroke Volume

B. Stroke Volume / Heart Rate

C. Heart Rate + Stroke Volume

D. Stroke Volume - Heart Rate

The correct approach to calculating QT when given stroke volume involves understanding the relationship between stroke volume and heart rate in the context of cardiac output. Cardiac output (CO) is defined as the volume of blood pumped by the heart per minute and is calculated using the formula: Cardiac Output = Stroke Volume x Heart Rate. In this case, QT, or the total time it takes to complete one cardiac cycle, can be viewed in terms of heart rate and stroke volume. Heart rate indicates how many times the heart beats in one minute, while stroke volume indicates how much blood is pumped with each beat. By multiplying heart rate by stroke volume, you get the total volume of blood pumped in a minute, thus providing a direct relation to how QT can be approached in a more integrative way. This is why multiplying heart rate by stroke volume is the correct method to relate these two parameters in the context of QT calculation. The other options involve incorrect operations (such as division or addition of these two separate values), which do not accurately represent the relationship in physiological terms.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ketteringpatientassmt.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE