KETTERING THERAPIST MULTIPLE-CHOICE (TMC) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ketteringtmc.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What condition is indicated by an abnormal pH due to a change in HCO3?

A. Metabolic acidosis

B. Respiratory alkalosis

C. Metabolic alkalosis

D. Respiratory acidosis

2. Which flow rate is most appropriate for a nasal cannula in stable COPD patients?

A. 1 - 3 L/min

B. 1 - 6 L/min

C. 4 - 10 L/min

D. 6 - 10 L/min

3. What is one of the absolute indications for mechanical ventilation?

A. Acute respiratory failure

B. Do Not Resuscitate orders

C. Patient requiring surgery

D. Chronic ventilatory failure

4. When the PaCO2 level is below 35 torr, what does this usually imply?

A. The patient is hypoventilating

B. The patient is ventilating too much

C. The patient requires mechanical ventilation

D. The patient's ventilation settings are optimal

5. What intervention should be taken when a patient has a PaCO2 above 45?

A. Decrease the mechanical ventilation rate

B. Initiate ventilation or increase current ventilation

C. Decrease patient's oxygen supply

D. Maintain current ventilation settings

SAMPLE

6. What disease state should a therapist suspect in a patient with dyspnea, diminished breath sounds, a flat percussion note on the left, and a tracheal shift to the right?

A. Pneumothorax.

B. Hemothorax.

C. Pleural Effusion.

D. Bronchial Obstruction.

7. When managing ventilation for a patient with hypoxemia, what is a suitable approach?

A. Increase the FiO2 while decreasing the ventilation

B. Increase mechanical ventilation at all times

C. Adjust FiO2 up to .60 and enhance ventilation

D. Monitor without adjusting the current settings

8. How does a pulmonary embolus typically appear on a chest X-ray?

A. Increased density from consolidation

B. Ground glass appearance

C. Peripheral wedge-shaped infiltrate

D. Reticulonodular pattern

9. Which change is expected in Maximum Voluntary Ventilation for a patient with obstructive disease?

A. Increased values

B. Decreased values

C. No change

D. Fluctuating values

10. What is the relationship between ventilation and end-tidal CO2 during hypercapnia?

A. Increased ventilation leads to increased end-tidal CO2

B. Decreased ventilation leads to increased end-tidal CO2

C. Ventilation has no relation to end-tidal CO2

D. Decreased perfusion leads to increased end-tidal CO2

SAMPLE

Answers

SAMPLE

1. A
2. B
3. A
4. B
5. B
6. C
7. C
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What condition is indicated by an abnormal pH due to a change in HCO3?

A. Metabolic acidosis

B. Respiratory alkalosis

C. Metabolic alkalosis

D. Respiratory acidosis

An abnormal pH due to a change in HCO3 (bicarbonate) is indicative of metabolic acidosis. In metabolic acidosis, there is a decrease in the bicarbonate level in the blood, which leads to an increased concentration of hydrogen ions and therefore a decrease in pH, signifying acidemia. This condition can arise from various causes, such as kidney failure, excessive production of acids (like in diabetic ketoacidosis), or loss of bicarbonate through diarrhea. In contrast, changes in pH associated with respiratory conditions are primarily due to alterations in carbon dioxide (CO2) levels rather than bicarbonate. For instance, respiratory acidosis is characterized by an increase in CO2 levels due to hypoventilation, whereas respiratory alkalosis corresponds to reduced CO2 levels due to hyperventilation. Metabolic alkalosis results from an increase in bicarbonate, leading to an elevated pH. Therefore, recognizing that an abnormal pH stemming from a change in bicarbonate specifically points to metabolic acidosis is essential for understanding acid-base balance and its implications in various medical conditions.

2. Which flow rate is most appropriate for a nasal cannula in stable COPD patients?

A. 1 - 3 L/min

B. 1 - 6 L/min

C. 4 - 10 L/min

D. 6 - 10 L/min

The most appropriate flow rate for a nasal cannula in stable COPD patients is in the range of 1 to 6 liters per minute. This flow rate is effective in delivering supplemental oxygen while minimizing the risk of carbon dioxide retention, which is particularly important for patients with COPD. In stable COPD patients, maintaining adequate oxygen saturation is crucial, and flow rates between 1 to 6 L/min are typically sufficient to achieve this without overwhelming the patient's respiratory drive. The lower end of this range allows for a gentler approach to oxygen therapy, helping to prevent hypercapnia (an elevation of carbon dioxide in the blood), which can occur if oxygen is administered at higher rates in these patients. Using flow rates outside of this range, such as higher rates, may lead to excessive oxygen delivery, which can be detrimental by causing a suppression of the hypoxic drive to breathe in those with COPD. Therefore, a flow rate within 1 to 6 L/min is tailored to the needs and safety of stable COPD patients.

SAMPLE

3. What is one of the absolute indications for mechanical ventilation?

A. Acute respiratory failure

B. Do Not Resuscitate orders

C. Patient requiring surgery

D. Chronic ventilatory failure

Acute respiratory failure is an absolute indication for mechanical ventilation because it denotes a critical situation where the patient's respiratory system is unable to ensure adequate gas exchange, leading to life-threatening hypoxemia or hypercapnia. In such a condition, the patient's ability to either breathe spontaneously or maintain proper oxygen and carbon dioxide levels is severely compromised, making mechanical ventilation necessary to support breathing, restore normal blood gas levels, and prevent further complications. The other options represent situations where mechanical ventilation may not be immediately required or could be decided based on other clinical parameters. For example, having Do Not Resuscitate orders does not inherently require mechanical ventilation; rather, it indicates the patient's wishes regarding the extent of resuscitative measures. A patient requiring surgery may not need mechanical ventilation unless they also exhibit signs of respiratory failure. Chronic ventilatory failure refers to a long-term condition where ventilation may be managed differently, possibly with non-invasive methods or prolonged home care strategies, rather than immediate mechanical ventilation.

4. When the PaCO2 level is below 35 torr, what does this usually imply?

A. The patient is hypoventilating

B. The patient is ventilating too much

C. The patient requires mechanical ventilation

D. The patient's ventilation settings are optimal

A PaCO2 level below 35 torr is indicative of respiratory alkalosis, which generally occurs when the body is eliminating carbon dioxide (CO2) at an increased rate. This condition typically suggests that the patient is hyperventilating, meaning they are ventilating too much rather than too little. When a patient is hyperventilating, they take in more oxygen and expel more carbon dioxide than necessary, leading to a decrease in the PaCO2 level. Hyperventilation can result from various factors, including anxiety, pain, or metabolic conditions, and is characterized by rapid, shallow breathing. In this context, the correct choice reflects this physiological response to excessive ventilation, allowing for a better understanding of the implications associated with low levels of carbon dioxide in the bloodstream.

SAMPLE

5. What intervention should be taken when a patient has a PaCO2 above 45?

A. Decrease the mechanical ventilation rate

B. Initiate ventilation or increase current ventilation

C. Decrease patient's oxygen supply

D. Maintain current ventilation settings

When a patient has a PaCO2 (arterial carbon dioxide pressure) above 45 mmHg, this indicates a condition known as hypercapnia, which reflects inadequate ventilation and an accumulation of carbon dioxide in the blood. The appropriate intervention is to initiate or increase mechanical ventilation to enhance the elimination of carbon dioxide. Increasing ventilation can be achieved through various means, which may include increasing the rate of mechanical breaths or adjusting the tidal volume delivered by the ventilator. This is crucial for restoring normal levels of PaCO2, as it helps the patient expel CO2 more effectively, thereby correcting the hypercapnia. Choosing to decrease mechanical ventilation or maintain current settings would not effectively address the elevated PaCO2 level and may allow the patient's condition to worsen. Decreasing the patient's oxygen supply would also be inappropriate, as it does not target the root issue of CO2 retention and could lead to further respiratory distress. Hence, the intervention of initiating or increasing current ventilation is aligned with the clinical imperative to manage hypercapnia and restore appropriate gas exchange.

6. What disease state should a therapist suspect in a patient with dyspnea, diminished breath sounds, a flat percussion note on the left, and a tracheal shift to the right?

A. Pneumothorax.

B. Hemothorax.

C. Pleural Effusion.

D. Bronchial Obstruction.

In this clinical scenario, the symptoms presented indicate that the most likely disease state is a pleural effusion. The presence of dyspnea suggests the patient is experiencing difficulty breathing, which can occur with any significant lung pathology. Diminished breath sounds noted on examination indicate a lack of airflow, which can be due to fluid accumulation in the pleural space, characteristic of pleural effusions. The flat percussion note on the left is a critical sign. In the case of a pleural effusion, fluid will accumulate in the pleural space and lead to a dull or flat sound upon percussion, as opposed to the resonance expected over normal lung tissue. This contrasts with a pneumothorax, which would produce hyperresonance due to air in the pleural space, or a hemothorax, where a dull sound would also occur, but with more association to blood specifically. Additionally, the observed tracheal shift to the right indicates a significant alteration in mediastinal structures, which can occur in the setting of a large pleural effusion. Fluid accumulation on one side, as would occur with a pleural effusion, can push the trachea towards the opposite side. Overall, the combination of diminished breath

SAMPLE

7. When managing ventilation for a patient with hypoxemia, what is a suitable approach?

A. Increase the FiO2 while decreasing the ventilation

B. Increase mechanical ventilation at all times

C. Adjust FiO2 up to .60 and enhance ventilation

D. Monitor without adjusting the current settings

The most suitable approach when managing ventilation for a patient with hypoxemia is to adjust the FiO2 up to 0.60 and enhance ventilation. This approach directly addresses the immediate need to improve oxygenation while also ensuring proper ventilation. Increasing the fraction of inspired oxygen (FiO2) to a level that is safe and effective (up to 0.60) helps to quickly raise the oxygen levels in the blood, which is crucial in cases of hypoxemia where oxygen saturation is low. Additionally, enhancing ventilation supports the removal of carbon dioxide and helps improve overall gas exchange in the lungs, which complements the increase in oxygen supply. This intervention provides a balanced approach to treating hypoxemia by addressing both oxygen delivery and ventilation adequacy. It recognizes the importance of not only providing higher levels of oxygen but also ensuring that the patient's lung function is supported to maintain an effective respiratory status. Other options may fall short by either increasing the risk of oxygen toxicity through excessive FiO2 without necessary ventilation support, failing to adjust settings appropriately for the clinical condition, or neglecting to actively manage the patient's ventilatory needs.

8. How does a pulmonary embolus typically appear on a chest X-ray?

A. Increased density from consolidation

B. Ground glass appearance

C. Peripheral wedge-shaped infiltrate

D. Reticulonodular pattern

A pulmonary embolus often manifests as a peripheral wedge-shaped infiltrate on a chest X-ray due to the way it obstructs blood flow in the lungs. This type of infiltrate is commonly referred to as a "hampton's hump". The wedge shape occurs because the area of lung affected by the embolism is supplied by a blocked pulmonary artery, potentially causing infarction and localized edema. In the case of a pulmonary embolism, this wedge-shaped appearance signifies that the distal lung region becomes hypoperfused, leading to the distinctive radiographic finding that aids in diagnosis. While other patterns exist on chest X-rays that may indicate various conditions, the peripheral wedge-shaped infiltrate is specifically associated with pulmonary emboli, marking its importance in clinical radiographic assessment. The increased density from consolidation typically indicates diseases like pneumonia, while a ground glass appearance is more seen in conditions such as pulmonary edema or interstitial lung disease. A reticulonodular pattern may be seen in fibrotic lung diseases, but these appearances do not specifically indicate a pulmonary embolism. This distinction underscores why recognizing the characteristic peripheral wedge-shaped infiltrate is crucial in the evaluation of a suspected pulmonary embolism.

SAMPLE

9. Which change is expected in Maximum Voluntary Ventilation for a patient with obstructive disease?

A. Increased values

B. Decreased values

C. No change

D. Fluctuating values

In patients with obstructive pulmonary disease, Maximum Voluntary Ventilation (MVV) is expected to show decreased values. MVV is a measure of the maximum amount of air that a person can breathe in and out in a specified period, reflecting overall respiratory muscle strength and lung function. In obstructive diseases, such as chronic obstructive pulmonary disease (COPD) or asthma, airflow limitation occurs due to narrowing or obstruction of the airways. This can be caused by inflammation, bronchoconstriction, or mucus production. As a result, patients may struggle to achieve high MVV values. During forced expiration phases, air gets trapped in the lungs, leading to a decreased ability to ventilate adequately. Thus, the functional impairment and the reduced air exchange in obstructive conditions contribute to a lower MVV, making the prediction of decreased values accurate in this context.

10. What is the relationship between ventilation and end-tidal CO2 during hypercapnia?

A. Increased ventilation leads to increased end-tidal CO2

B. Decreased ventilation leads to increased end-tidal CO2

C. Ventilation has no relation to end-tidal CO2

D. Decreased perfusion leads to increased end-tidal CO2

In the context of hypercapnia, which is an increased level of carbon dioxide (CO2) in the blood, the relationship between ventilation and end-tidal CO2 is crucial for understanding respiratory physiology. When ventilation decreases, the clearance of CO2 from the lungs is impaired. As a result, CO2 accumulates in the blood, leading to higher levels of end-tidal CO2, which reflects the amount of carbon dioxide present in the exhaled air at the end of expiration. This correlation indicates that when ventilation is insufficient (as seen in conditions such as respiratory depression or obstructive pulmonary diseases), the body cannot effectively eliminate CO2, causing an increase in end-tidal CO2 levels. Therefore, the choice stating that decreased ventilation leads to increased end-tidal CO2 is accurate, as it aligns with the physiological principles governing gas exchange in the lungs. Understanding this relationship is essential for healthcare professionals, particularly those in respiratory therapy, as it helps in assessing a patient's ventilatory status and making informed decisions regarding therapy and interventions.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ketteringtmc.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE