LEARNING SYSTEMS MED SURG PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://learningsystemsmedsurg.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is an early sign of infection in a surgical patient?

A. Increased urination

B. Fever and chills

C. Low blood sugar

D. Chest pain

2. Which assessment should be included for patients receiving anticoagulants?

A. Assessment of blood pressure

B. Monitoring for signs of internal bleeding

C. Checking blood glucose levels

D. Evaluating kidney function

3. What action should a nurse take when preparing to remove a client's chest tube?

A. Have the client breathe normally during removal

B. Instruct the client to perform the Valsalva maneuver

C. Immediately pull the tube out swiftly

D. Apply pressure to the incision site afterward

4. What statement by a client with systemic lupus erythematosus indicates understanding of discharge teaching?

A. "I will need to take methotrexate even if I'm in remission"

B. "I can stop my medication once I feel better"

C. "I do not need to follow up with my doctor regularly"

D. "I should avoid sun exposure at all times"

5. What is the primary goal of treatment for right-sided heart failure?

A. Complete recovery of heart function

B. Control symptoms and improve quality of life

C. Reduce risk of surgery

D. Eliminate need for medication

SAMPLE

6. What is a common symptom of acute pancreatitis?

A. Mild abdominal cramps

B. Severe abdominal pain radiating to the back

C. Nausea without vomiting

D. Loss of appetite

7. Which laboratory finding is significant in diagnosing renal failure?

A. Low potassium levels

B. High creatinine levels

C. High albumin levels

D. Low urine specific gravity

8. What symptom might suggest a patient is experiencing a myocardial infarction?

A. Severe chest pain radiating to the left arm

B. Dizziness and shortness of breath

C. Pain in the lower abdomen

D. Fatigue and nausea

9. What is the primary concern in the management of a patient with a suspected stroke?

A. Administering antibiotics

B. Time-sensitive administration of thrombolytics

C. Monitoring blood pressure

D. Providing anticoagulants

10. When monitoring a patient in anaphylactic shock, what should be prioritized?

A. Fluid resuscitation

B. Airway management

C. Monitoring heart rate

D. Administering antihistamines

SAMPLE

Answers

SAMPLE

1. B
2. B
3. B
4. A
5. B
6. B
7. B
8. A
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is an early sign of infection in a surgical patient?

A. Increased urination

B. Fever and chills

C. Low blood sugar

D. Chest pain

Fever and chills are considered early signs of infection or an inflammatory response in a surgical patient. After surgery, the body's immune system may react to surgical trauma, and any increase in temperature, even if mild, can indicate that the body is fighting off an infection. The presence of chills often accompanies a fever, further reinforcing the body's response to an infectious process. Monitoring vital signs, including temperature, is a crucial aspect of post-operative care, as early identification of fever allows for prompt intervention and management, which is essential to prevent further complications. Increased urination may relate to various factors post-surgery, such as fluid management or diuretics, but it does not specifically indicate an early infection. Low blood sugar is also not directly linked to infection; rather, it may occur due to fasting or changes in metabolic state after anesthesia. Chest pain could result from multiple causes post-surgery, such as respiratory complications or cardiovascular issues, but it is not a typical early indicator of infection. Therefore, recognizing fever and chills is essential in the context of surgical recovery and complications associated with infections.

2. Which assessment should be included for patients receiving anticoagulants?

A. Assessment of blood pressure

B. Monitoring for signs of internal bleeding

C. Checking blood glucose levels

D. Evaluating kidney function

Monitoring for signs of internal bleeding is crucial for patients receiving anticoagulants, as these medications are designed to prevent blood clots but can also increase the risk of bleeding. Patients on anticoagulant therapy are particularly vulnerable to both minor and severe internal bleeding, which can present as symptoms such as unusual bruising, blood in urine or stool, or severe headache and dizziness, indicative of potential intracranial bleeding. Early detection of these signs can significantly impact patient safety and outcomes, allowing for timely intervention. While assessing blood pressure is important in a comprehensive evaluation, it does not specifically address the risks associated with anticoagulant therapy. Checking blood glucose levels is necessary for patients with diabetes or those at risk but is not a direct concern of anticoagulation management. Evaluating kidney function is essential, especially since renal impairment can influence how anticoagulants are metabolized and excreted, but it does not directly relate to the immediate risks posed by the medications in terms of bleeding complications. Thus, monitoring for signs of internal bleeding is the most pertinent assessment for patients on anticoagulants.

SAMPLE

3. What action should a nurse take when preparing to remove a client's chest tube?

A. Have the client breathe normally during removal

B. Instruct the client to perform the Valsalva maneuver

C. Immediately pull the tube out swiftly

D. Apply pressure to the incision site afterward

Instructing the client to perform the Valsalva maneuver during chest tube removal is important as it helps to create a transient increase in intrathoracic pressure. This action aids in preventing air from entering the pleural space as the tube is pulled out, thereby minimizing the risk of pneumothorax. The Valsalva maneuver involves the client taking a deep breath and holding it while bearing down, which helps to stabilize the pleural cavity during the procedure. To ensure safe and effective removal of the chest tube, the nurse should educate the client about this maneuver beforehand. Proper technique is essential, as it contributes to better patient outcomes following the removal. Additionally, this approach is supported by clinical practice guidelines that advocate for protective measures against complications associated with chest tube extraction. In contrast, having the client breathe normally may leave the pleural space vulnerable to air entry during removal. Swiftly pulling the tube out can disrupt the intrathoracic pressure protective mechanisms and cause discomfort or injury. Furthermore, while applying pressure to the incision site is a critical step after removal to minimize bleeding and seal the site, it is not performed until after the tube has been safely extracted. Therefore, performing the Valsalva maneuver is the most appropriate action to

4. What statement by a client with systemic lupus erythematosus indicates understanding of discharge teaching?

A. "I will need to take methotrexate even if I'm in remission"

B. "I can stop my medication once I feel better"

C. "I do not need to follow up with my doctor regularly"

D. "I should avoid sun exposure at all times"

The statement indicating understanding of discharge teaching for a client with systemic lupus erythematosus is that the client acknowledges the need to continue taking methotrexate even when in remission. This reflects a comprehensive understanding of the chronic nature of lupus, which often involves periods of flare and remission. In many cases, treatment protocols may require ongoing medication to manage the disease effectively and prevent potential flare-ups, even when symptoms are well-controlled. Continuing such medications is essential because stopping them prematurely could lead to a resurgence of symptoms or complications associated with the disease. The other statements demonstrate a lack of understanding regarding the management of lupus. For example, stopping medication upon feeling better suggests a misunderstanding of the need for consistent treatment. Failing to follow up with a healthcare provider can lead to unmanaged disease progression, and ignoring sun exposure precautions can trigger skin rashes or flares, which are common in systemic lupus erythematosus. Therefore, recognizing the importance of ongoing medication and careful management is critical for maintaining long-term health in patients with this condition.

SAMPLE

5. What is the primary goal of treatment for right-sided heart failure?

A. Complete recovery of heart function

B. Control symptoms and improve quality of life

C. Reduce risk of surgery

D. Eliminate need for medication

The primary goal of treatment for right-sided heart failure is to control symptoms and improve quality of life. In patients with this condition, the right side of the heart struggles to pump blood effectively to the lungs, leading to symptoms such as swelling, fatigue, and shortness of breath. Managing these symptoms is crucial for helping patients maintain their daily activities and overall well-being. While complete recovery of heart function may be an ideal outcome, many patients with right-sided heart failure may not achieve this, and treatments typically focus on symptom management. The emphasis is on medications, lifestyle adjustments, and sometimes device interventions that can help manage the condition rather than aiming for a complete reversal of heart function. Moreover, reducing the risk of surgery or eliminating the need for medication are not realistic primary goals for patients with this form of heart failure. Instead, the focus is on making sure that the patient can lead a more comfortable and functional life despite the chronic nature of the disease. Therefore, symptom control and enhancing the quality of life are fundamental in the management strategy for right-sided heart failure.

6. What is a common symptom of acute pancreatitis?

A. Mild abdominal cramps

B. Severe abdominal pain radiating to the back

C. Nausea without vomiting

D. Loss of appetite

Severe abdominal pain radiating to the back is a hallmark symptom of acute pancreatitis. This pain often presents in the upper abdomen and intensifies after eating or drinking, particularly fatty foods. The reason for the pain's radiating quality is linked to the pancreas's anatomical position; when inflamed, irritation can extend to surrounding structures, including the nerves that can transmit pain signals to the back. This symptom is significant because it helps clinicians differentiate acute pancreatitis from other abdominal conditions. While other symptoms may accompany acute pancreatitis, such as nausea, vomiting, and loss of appetite, the presence of severe abdominal pain that radiates to the back is the most defining and critical for diagnosis. Understanding the presentation of symptoms, particularly the nature and location of pain, is essential for effective management and treatment of this condition.

SAMPLE

7. Which laboratory finding is significant in diagnosing renal failure?

A. Low potassium levels

B. High creatinine levels

C. High albumin levels

D. Low urine specific gravity

High creatinine levels are a significant indicator in diagnosing renal failure because creatinine is a waste product produced from normal muscle metabolism that is usually filtered out of the blood by the kidneys. When the kidneys are not functioning properly, they are unable to adequately filter creatinine from the bloodstream, leading to elevated levels. This increase in creatinine levels is often used as a key biomarker in assessing kidney function and is a critical part of diagnosing acute and chronic renal failure. Elevated creatinine levels provide valuable information about the glomerular filtration rate (GFR), which is a measure of how well the kidneys filter blood. When GFR decreases due to kidney damage or disease, serum creatinine levels rise, indicating renal impairment. Therefore, high levels of creatinine are directly related to renal dysfunction and are a primary focus in renal failure diagnosis. Other laboratory findings mentioned, such as low potassium levels, high albumin levels, and low urine specific gravity, may indicate different conditions and are not as directly linked to the diagnosis of renal failure. For example, while high potassium levels (hyperkalemia) can occur in renal failure, low potassium does not indicate kidney dysfunction. High levels of albumin can suggest other conditions, like liver disease or nephrotic syndrome

8. What symptom might suggest a patient is experiencing a myocardial infarction?

A. Severe chest pain radiating to the left arm

B. Dizziness and shortness of breath

C. Pain in the lower abdomen

D. Fatigue and nausea

Severe chest pain radiating to the left arm is a classic symptom of myocardial infarction, commonly referred to as a heart attack. This symptom is often described as a pressure, squeezing, or intense discomfort in the chest that can extend to other areas such as the arms, neck, jaw, or back. The radiating pain to the left arm is particularly significant because it aligns with the pattern of referred pain often experienced during cardiac events due to the way nerves are interconnected in the body. Recognizing this symptom is crucial for timely intervention, as a myocardial infarction can quickly lead to serious complications or death if not treated promptly. While dizziness and shortness of breath, pain in the lower abdomen, and fatigue accompanied by nausea can sometimes occur during a myocardial infarction, they are not as specific or characteristic as chest pain radiating to the left arm. These symptoms can also indicate a variety of other conditions, which makes them less definitive in diagnosing a heart attack. Therefore, the combination of severe chest pain and its radiation to the left arm is a more reliable indicator of myocardial infarction, highlighting the importance of chest pain in cardiac diagnoses.

SAMPLE

9. What is the primary concern in the management of a patient with a suspected stroke?

A. Administering antibiotics

B. Time-sensitive administration of thrombolytics

C. Monitoring blood pressure

D. Providing anticoagulants

The primary concern in the management of a patient with a suspected stroke is the time-sensitive administration of thrombolytics. This approach is crucial because certain types of strokes, specifically ischemic strokes, occur due to a blockage in the blood vessels supplying the brain. Thrombolytics, or clot-dissolving medications, can significantly improve outcomes if given promptly, ideally within a narrow therapeutic window, typically within 3 to 4.5 hours of symptom onset. Timely intervention can reduce brain damage and disability by restoring blood flow to affected areas in the brain. Therefore, recognizing stroke symptoms, assessing the patient efficiently, and rapidly initiating treatment with thrombolytics are vital steps in the emergency management of a stroke. Other choices, while they may play a role in stroke management, do not address the urgent need to restore blood flow as effectively as thrombolytics do. Administering antibiotics is pertinent in managing infections, which may be a concern in post-stroke care but is not a priority during the acute phase. Monitoring blood pressure is essential in managing a stroke patient but does not directly treat the underlying issue of the blocked blood vessel. Providing anticoagulants may be necessary later in stroke management, particularly in the event of an embolic stroke,

10. When monitoring a patient in anaphylactic shock, what should be prioritized?

A. Fluid resuscitation

B. Airway management

C. Monitoring heart rate

D. Administering antihistamines

In the context of anaphylactic shock, airway management is of utmost priority because this condition can lead to severe and rapid obstruction of the airway due to swelling and bronchoconstriction. During an anaphylactic reaction, the release of histamines and other inflammatory mediators can cause the throat and airways to swell, making it difficult for the patient to breathe. When a patient is experiencing anaphylaxis, they can develop stridor, wheezing, or even complete loss of airway patency, which necessitates immediate intervention to secure the airway. If the airway is not established promptly, the patient could suffer from hypoxia, leading to respiratory failure and potential death. While fluid resuscitation, monitoring heart rate, and administering antihistamines are all important components of treating anaphylactic shock, they are secondary to ensuring that the patient has a clear and open airway. Administering epinephrine is also critical, as it helps counteract the allergic reaction quickly, but airway management should always be prioritized to prevent life-threatening complications.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://learningsystemsmedsurg.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE