LEIK FAMILY NURSE PRACTITIONER (FNP) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://leikfamilynursepracticioner.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which factor is not associated with an increased risk of osteopenia in teenage girls?

A. Drinking one glass of low-fat milk daily

B. Anorexia nervosa

C. Participation in sports

D. A normal BMI (basal metabolic index)

2. Which microorganism is the most common cause of otitis externa?

A. Pseudomonas aeruginosa

B. Streptococcus pyogenes

C. Haemophilus influenza

D. Moraxella catarrhalis

3. If a patient experiences a severe allergic reaction to Bactrim, what is the correct advice regarding sulfonylurea medications?

A. She cannot take any pills in the sulfonylurea class

B. She can take some of the pills in the sulfonylurea class

C. She can take any of the pills in the sulfonylurea class

D. None of the above

4. Koplik's spots are associated with which of the following conditions?

A. Poxvirus infections

B. Measles

C. Kawasaki's disease

D. Reye's syndrome

5. Cullen's sign is most commonly associated with which medical condition?

A. Acute pancreatitis

B. Myocardial infarction

C. Acute pyelonephritis

D. Preeclampsia

SAMPLE

6. Which complication is associated with Chronic lung infections due to Pseudomonas aeruginosa?

A. Increased immune response

B. Chronic fibrotic lung disease

C. Rapid recovery from infections

D. Decreased inflammation response

7. What is a common clinical presentation of diabetes mellitus?

A. Increased blood pressure

B. Excessive sweating

C. Frequent urination

D. Visual disturbances

8. Which drug is safe to use when taking erythromycin, considering potential drug interactions?

A. Theophylline (Theo-Dur)

B. Warfarin (Coumadin)

C. Diazepam (Valium)

D. Furosemide (Lasix)

9. All of the following statements about oral contraceptives are correct except?

A. The actual failure rate of oral contraceptives is 3%

B. Desogestrel belongs to the progesterone family of drugs

C. The newer low-dose birth control pills require backup during the first 2 weeks of use

D. Oral contraceptives are contraindicated for women 35 or older who smoke

10. Which sign can mimic testicular torsion but is not an emergent condition?

A. The "blue dot" sign

B. Swollen testicle with penile discharge

C. Acute onset of dysuria and frequency

D. A varicocele

SAMPLE

Answers

SAMPLE

1. A
2. A
3. A
4. B
5. A
6. B
7. C
8. D
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. Which factor is not associated with an increased risk of osteopenia in teenage girls?

A. Drinking one glass of low-fat milk daily

B. Anorexia nervosa

C. Participation in sports

D. A normal BMI (basal metabolic index)

Osteopenia is a condition characterized by lower than average bone density, which can increase the risk of fractures and osteoporosis later in life. Various factors can contribute to increased risk, especially in developing individuals like teenage girls, who are in a crucial bone-building phase. Drinking one glass of low-fat milk daily provides an important source of calcium and vitamin D, both essential nutrients for bone health. Adequate calcium intake is vital during adolescence for achieving peak bone mass, and a consistent intake may help to reduce the risk of osteopenia. Therefore, the inclusion of low-fat milk in the diet would not be associated with an increased risk of developing osteopenia; rather, it could be protective against it. In contrast, other factors such as anorexia nervosa commonly lead to malnutrition and insufficient calorie and nutrient intake, directly impacting bone density and increasing the risk of osteopenia. Participation in sports, particularly those that emphasize leanness or have weight restrictions, can also contribute to disordered eating patterns and, subsequently, low bone density. Lastly, having a low BMI can reflect inadequate nutritional intake or high levels of physical activity without sufficient caloric compensation, further putting girls at risk for lower bone density. Thus, considering these factors,

2. Which microorganism is the most common cause of otitis externa?

A. Pseudomonas aeruginosa

B. Streptococcus pyogenes

C. Haemophilus influenza

D. Moraxella catarrhalis

Pseudomonas aeruginosa is recognized as the most common pathogen responsible for otitis externa, often referred to as swimmer's ear. This condition typically arises when water becomes trapped in the ear canal, creating a moist environment conducive to bacterial growth. Pseudomonas aeruginosa thrives in such environments; it's a gram-negative bacterium commonly found in water and soil, which explains its prevalence in cases of otitis externa, especially in individuals who swim frequently or have increased exposure to water. The virulence of Pseudomonas aeruginosa is attributed to its ability to form biofilms, resist antibiotics, and cause inflammatory responses within the ear canal. This makes it particularly effective at establishing infections in the moist environment of the ear, where it can flourish and lead to the classic symptoms of otitis externa, such as pain, itching, and discharge. Other organisms like Streptococcus pyogenes, Haemophilus influenza, and Moraxella catarrhalis are associated with various ear infections, but they more commonly cause otitis media rather than otitis externa. Therefore, recognizing Pseudomonas aeruginosa's role in otitis externa is crucial for appropriate evaluation and management in patients presenting with symptoms indicative of this condition.

SAMPLE

3. If a patient experiences a severe allergic reaction to Bactrim, what is the correct advice regarding sulfonylurea medications?

A. She cannot take any pills in the sulfonylurea class

B. She can take some of the pills in the sulfonylurea class

C. She can take any of the pills in the sulfonylurea class

D. None of the above

A patient who has experienced a severe allergic reaction to Bactrim, which contains sulfamethoxazole, is at a heightened risk for cross-reactivity with sulfonylureas, a class of medications used to treat diabetes. Sulfonylureas, such as glyburide, glipizide, and glimepiride, are chemically related to sulfonamide antibiotics and can provoke similar allergic responses in patients with sulfonamide allergies. Due to the potential for serious allergic reactions, it is advisable to avoid all medications within the sulfonylurea class for someone with a known severe allergy to Bactrim. The recommendation stems from the concern that susceptibility to allergic reactions may persist across similar chemical structures, making it prudent to steer clear of these medications altogether. On the other hand, while some may consider that not every sulfonylurea medication may cause an allergic response, the risk is significant enough to warrant avoidance of all drugs in this category for individuals with a severe allergy history.

4. Koplik's spots are associated with which of the following conditions?

A. Poxvirus infections

B. Measles

C. Kawasaki's disease

D. Reye's syndrome

Koplik's spots are characteristic lesions that appear in the oral cavity and are specifically associated with measles (rubeola). These spots are small, bluish-white lesions with a red halo that appear on the buccal mucosa, typically two to three days before the rash associated with measles develops. They serve as an important clinical sign that helps healthcare providers identify and diagnose measles early in the course of the illness. In contrast, the other conditions listed do not present with Koplik's spots. Poxvirus infections, such as those caused by variola or vaccinia, are associated with different types of skin lesions, typically vesicular or pustular lesions on the skin. Kawasaki's disease is a vasculitis affecting children and is characterized by high fever, rash, and conjunctivitis but does not include Koplik's spots as a symptom. Reye's syndrome is linked to a viral infection and aspirin use in children, presenting primarily with encephalopathy and liver dysfunction rather than oral lesions like Koplik's spots. Thus, the association of these spots specifically with measles makes it clear that option B is the correct choice.

SAMPLE

5. Cullen's sign is most commonly associated with which medical condition?

A. Acute pancreatitis

B. Myocardial infarction

C. Acute pyelonephritis

D. Preeclampsia

Cullen's sign is a clinical manifestation characterized by a discoloration or bruising around the umbilical area, usually indicating internal bleeding. It is most commonly associated with acute pancreatitis, particularly when there is necrotizing pancreatitis, which can lead to hemorrhage in the retroperitoneal space that ultimately tracks to the umbilical region. This sign may occur due to the presence of blood in the peritoneal cavity. In the context of acute pancreatitis, Cullen's sign can help in the initial assessment of the severity of the disease and potential complications, guiding further diagnostic and therapeutic interventions. While other conditions, such as ectopic pregnancy or hemorrhagic conditions, can also present with Cullen's sign, it is most closely tied to acute pancreatitis in clinical practice scenarios. The other conditions listed—myocardial infarction, acute pyelonephritis, and preeclampsia—do not typically present with Cullen's sign, which helps to reiterate that this sign is distinctive in the context of diagnosing acute pancreatitis.

6. Which complication is associated with Chronic lung infections due to Pseudomonas aeruginosa?

A. Increased immune response

B. Chronic fibrotic lung disease

C. Rapid recovery from infections

D. Decreased inflammation response

Chronic lung infections caused by Pseudomonas aeruginosa are associated with chronic fibrotic lung disease because this bacterium is notorious for causing long-lasting infections that can lead to significant lung tissue damage. Pseudomonas aeruginosa is often implicated in cystic fibrosis and other underlying conditions that weaken lung function. The persistent inflammation and immune response triggered by these chronic infections can lead to the release of inflammatory mediators and fibroblast activation, causing collagen deposition and resulting in fibrosis of the lung tissue. This fibrosis impairs lung function as it replaces healthy lung tissue with scar tissue, limiting the ability of the lungs to expand and contract effectively. Over time, this can lead to progressive respiratory failure and a decline in overall health in affected individuals. Therefore, understanding the relationship between Pseudomonas aeruginosa and chronic fibrotic lung disease is crucial for managing patients with recurrent lung infections effectively.

SAMPLE

7. What is a common clinical presentation of diabetes mellitus?

A. Increased blood pressure

B. Excessive sweating

C. Frequent urination

D. Visual disturbances

Frequent urination, also known as polyuria, is a hallmark symptom of diabetes mellitus. This occurs as a result of elevated blood glucose levels. When blood sugar is high, the kidneys work to filter out the excess glucose. As glucose is excreted in urine, it pulls water along with it, leading to increased urine production and frequency of urination. This can be particularly noticeable in instances where blood sugar levels remain uncontrolled. The osmotic diuresis caused by high glucose levels is a crucial aspect of the pathophysiology of diabetes. In contrast, while increased blood pressure, excessive sweating, and visual disturbances can be associated with diabetes, they are not as direct or common a presenting symptom as polyuria. Increased blood pressure can be a complication of long-standing diabetes but is not primarily a presenting symptom. Excessive sweating may occur in some individuals, particularly if affected by autonomic neuropathy, but it's not a classic symptom of diabetes. Visual disturbances can arise from diabetes-related complications, such as diabetic retinopathy or cataracts, but they typically develop over time, rather than serving as an early warning sign of the disease and are not often seen in initial presentations.

8. Which drug is safe to use when taking erythromycin, considering potential drug interactions?

A. Theophylline (Theo-Dur)

B. Warfarin (Coumadin)

C. Diazepam (Valium)

D. Furosemide (Lasix)

Erythromycin is an antibiotic that can significantly affect the metabolism of various drugs due to its inhibition of the cytochrome P450 system, particularly CYP3A4. This interaction can lead to increased plasma levels of other medications that are metabolized by the same pathway, potentially causing toxicity. Furosemide, a loop diuretic, is not primarily metabolized by CYP3A4 and therefore is considered safe to use concomitantly with erythromycin. It works by a different mechanism, primarily by inhibiting sodium and chloride reabsorption in the ascending loop of Henle, leading to diuresis without the risk of serious drug interactions with erythromycin. In contrast, the other listed medications pose significant risks when taken with erythromycin. Theophylline, for example, can have its clearance decreased, leading to increased plasma concentrations and potential toxicity. Warfarin has an increased risk of bleeding due to altered metabolism, and diazepam could result in elevated blood levels, increasing the risk of sedation and respiratory depression. Therefore, furosemide stands out as the safest option among these drugs for co-administration with erythromycin, minimizing the risk of adverse effects.

SAMPLE

9. All of the following statements about oral contraceptives are correct except?

A. The actual failure rate of oral contraceptives is 3%

B. Desogestrel belongs to the progesterone family of drugs

C. The newer low-dose birth control pills require backup during the first 2 weeks of use

D. Oral contraceptives are contraindicated for women 35 or older who smoke

The assertion regarding newer low-dose birth control pills requiring backup during the first two weeks of use is often misunderstood. In general, it is advised that women begin using a backup method of contraception, such as condoms, during the initial week of using hormonal contraceptives, specifically if the pills are not initiated during the first five days of their menstrual cycle. However, current guidelines suggest that in many cases, this might not be necessary if the pill is taken consistently and correctly, as the efficacy can be established more quickly than previous formulations indicated. This contrasts with the knowledge of how oral contraceptives work and the traditional recommendations that may no longer apply consistently, especially with modern advancements in contraceptive formulations. As such, while women new to oral contraceptives are often advised to use backup, newer low-dose pills might not have the same stringent requirement for backup for as long as two weeks, depending on the specific guidelines of the product and the healthcare provider's recommendations. Thus, the statement lacks a general applicability regarding current practices and recommendations surrounding oral contraceptive use.

10. Which sign can mimic testicular torsion but is not an emergent condition?

A. The "blue dot" sign

B. Swollen testicle with penile discharge

C. Acute onset of dysuria and frequency

D. A varicocele

The "blue dot" sign is indicative of a torsion of the appendage of the testis, specifically the tunica vaginalis or accessory testicular tissue. This sign occurs when the blood supply to the torted appendage is compromised, leading to localized ischemia and subsequent necrosis, giving rise to the characteristic blue appearance. While this sign can mimic testicular torsion, it is not considered a true emergency as the underlying issue does not typically jeopardize the viability of the testis itself. In contrast, the other options describe conditions that may not necessarily present with the same acute vascular emergency as testicular torsion. A swollen testicle with penile discharge typically indicates an infectious process, which, while requiring urgent care, does not present the same acute risk as torsion. Acute dysuria and frequency suggest a urinary tract infection rather than torsion, and a varicocele represents a venous dilation, which is generally a chronic and non-emergent condition. Understanding the distinct clinical presentations and implications behind each symptom allows for a more accurate assessment in differentiating conditions that require immediate medical intervention from those that are less acute.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://leikfamilynursepracticioner.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE