MEDICAL HISTORY COMPETENCY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://medhistorycompetency.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. How should language barriers be managed and interpreters used during history-taking?

A. Use automated translation apps only.

B. Use professional interpreters, avoid family members, brief interpreter, verify accuracy, consider patient comfort.

C. Skip questions.

D. Rely on family members.

2. What are the three types of diabetes and their defining characteristics?

A. Type 1: juvenile, insulin-dependent, little or no insulin produced; Type 2: insulin resistant, body produces insulin but doesn't know how to use it; Type 3: gestational diabetes during pregnancy.

B. Type 1: adult-onset, insulin-dependent; Type 2: insulin dependent; Type 3: gestational diabetes.

C. Type 1: gestational diabetes; Type 2: obesity-related diabetes; Type 3: juvenile diabetes.

D. Type 1: juvenile, insulin-dependent; Type 2: gestational diabetes; Type 3: obesity-related diabetes.

3. What is the main mechanism behind angina pectoris according to the source?

A. Thrombosis in an atherosclerotic artery

B. Bacterial infection of coronary arteries

C. Low blood pressure

D. Dehydration

4. Reactivation of which virus can cause shingles?

A. Herpes simplex virus

B. Epstein-Barr virus

C. Varicella-Zoster virus

D. Cytomegalovirus

5. PT stands for what in coagulation testing?

A. Prothrombin Time

B. Partial Thromboplastin Time

C. Platelet Time

D. Prothrombin Test

SAMPLE

6. What is the standard dose of Cephalexin used as an alternative for penicillin-allergic patients?

A. 1 g

B. 500 mg

C. 2 g

D. 750 mg

7. What characterizes a nonjudgmental approach to discussing social history topics like tobacco, alcohol, and substance use?

A. Normalize disclosures about tobacco, alcohol, illicit drugs; discuss lifestyle factors nonjudgmentally.

B. Guilt and admonition.

C. Only ask about smoking.

D. Avoid discussing lifestyle.

8. What approach is recommended when evaluating a patient with multiple health issues to improve care?

A. Focus only on the most recent problem

B. Use a structured approach, including chief concerns, review of systems, medications, and risk factors

C. Skip review of systems

D. Limit to one paragraph

9. What does pack-years quantify in tobacco exposure history?

A. The current nicotine level in blood.

B. The number of times the patient has smoked in life.

C. The average cigarette nicotine content.

D. The product of packs smoked per day and years of smoking.

10. Which food is associated with latex allergy?

A. Carrot

B. Rice

C. Avocado

D. Chicken

SAMPLE

Answers

SAMPLE

1. B
2. B
3. C
4. C
5. A
6. C
7. A
8. B
9. D
10. C

SAMPLE

Explanations

SAMPLE

1. How should language barriers be managed and interpreters used during history-taking?

A. Use automated translation apps only.

B. Use professional interpreters, avoid family members, brief interpreter, verify accuracy, consider patient comfort.

C. Skip questions.

D. Rely on family members.

When taking history from a patient who speaks a different language, the focus is on accurate, respectful communication through a trained interpreter. Using professional interpreters ensures medical terminology is conveyed correctly and that confidentiality and neutrality are maintained, which is essential for a truthful and complete history. Avoid involving family members. They may introduce bias, omit or alter details, or unintentionally pressure the patient, and confidentiality can be breached. A brief, pre-session with the interpreter helps set expectations: explain the clinical goals, highlight any sensitive topics that may come up, and outline preferred terminology or patient preferences. This background helps the interpreter render messages as clearly and accurately as possible. Verifying accuracy is important. After the history is gathered, or during it, check that the patient understands key points and that the interpreter has conveyed responses faithfully. Using simple language, pausing for questions, and, when appropriate, using teach-back techniques helps confirm understanding. Also consider patient comfort and dignity: respect cultural norms, offer a choice of interpreter gender if possible, ensure privacy, and allocate enough time so the patient doesn't feel rushed. Automated translation apps are unreliable for medical history, and skipping questions or relying on family members can compromise accuracy, confidentiality, and clinician-patient rapport.

2. What are the three types of diabetes and their defining characteristics?

A. Type 1: juvenile, insulin-dependent, little or no insulin produced; Type 2: insulin resistant, body produces insulin but doesn't know how to use it; Type 3: gestational diabetes during pregnancy.

B. Type 1: adult-onset, insulin-dependent; Type 2: insulin dependent; Type 3: gestational diabetes.

C. Type 1: gestational diabetes; Type 2: obesity-related diabetes; Type 3: juvenile diabetes.

D. Type 1: juvenile, insulin-dependent; Type 2: gestational diabetes; Type 3: obesity-related diabetes.

Understanding diabetes types comes from how insulin is produced and used in the body. Type 1 is an autoimmune process that destroys insulin-producing cells, so the body makes little or no insulin and people usually need insulin therapy. Type 2 starts with insulin resistance—cells don't respond well to insulin—and often with insufficient insulin production to meet demand; it's commonly seen in adults and is frequently linked to obesity, with management that may begin with lifestyle changes and oral medications and may progress to insulin. Gestational diabetes is a distinct form that appears during pregnancy due to hormones that raise insulin resistance, and it typically resolves after delivery although it raises future diabetes risk. The description that fits these concepts is the one that says Type 1 is juvenile and insulin-dependent with little or no insulin produced, Type 2 is insulin resistant with continued insulin production, and Type 3 is gestational diabetes during pregnancy. The other options mix up onset age, insulin dependence, or the pregnancy-associated nature of gestational diabetes.

SAMPLE

3. What is the main mechanism behind angina pectoris according to the source?

A. Thrombosis in an atherosclerotic artery

B. Bacterial infection of coronary arteries

C. Low blood pressure

D. Dehydration

Angina pectoris comes from the heart not getting enough oxygen to meet its needs, a mismatch between supply and demand. The source highlights the driving factor as reduced coronary perfusion pressure. When systemic blood pressure drops, the pressure gradient pushing blood into the coronary vessels falls, especially during diastole when the coronaries are normally perfused. In someone with already narrowed (atherosclerotic) arteries, this reduction in driving pressure can markedly limit oxygen-rich blood reaching the heart muscle. As the heart's workload increases—during exercise or stress—oxygen demand rises while supply remains constrained, leading to chest pain. So, low blood pressure is described as the main mechanism because it directly lowers the coronary blood flow available to the myocardium, triggering ischemia and anginal symptoms in susceptible individuals.

4. Reactivation of which virus can cause shingles?

A. Herpes simplex virus

B. Epstein-Barr virus

C. Varicella-Zoster virus

D. Cytomegalovirus

Shingles arises when the varicella-zoster virus, which lies dormant after the initial chickenpox infection, reactivates. This virus hides in the dorsal root or cranial nerve ganglia and, upon reactivation, travels along a sensory nerve to the skin, producing a painful, unilateral vesicular rash confined to a dermatome. Other herpesviruses don't cause the classic shingles presentation: herpes simplex reactivates mainly as cold sores or genital lesions, cytomegalovirus and Epstein-Barr virus have different reactivation patterns and illnesses, particularly in people with weakened immune systems. The specific reactivation that leads to the shingles rash is characteristic of the varicella-zoster virus.

5. PT stands for what in coagulation testing?

A. Prothrombin Time

B. Partial Thromboplastin Time

C. Platelet Time

D. Prothrombin Test

PT stands for Prothrombin Time. It measures how long it takes for plasma to clot after adding tissue factor (thromboplastin) and calcium, reflecting the extrinsic pathway of coagulation and the common pathway. It is especially influenced by vitamin K–dependent factors II, VII, IX, and X, making it sensitive to vitamin K deficiency and to anticoagulants like warfarin. Clinically, PT is used to monitor warfarin therapy and to assess liver function and clotting factor deficiencies; results are often reported in seconds and standardized as an INR for comparison across labs. The other terms refer to different concepts: Partial Thromboplastin Time assesses the intrinsic pathway, Platelet Time isn't a standard coagulation test, and Prothrombin Test isn't the conventional name.

SAMPLE

6. What is the standard dose of Cephalexin used as an alternative for penicillin-allergic patients?

A. 1 g

B. 500 mg

C. 2 g

D. 750 mg

Cephalexin is chosen as an alternative when penicillin can't be used because it provides similar coverage against the common oral flora and skin pathogens, and it is reliably absorbed when taken by mouth. For penicillin-allergic adults undergoing procedures where bacterial exposure is a concern, the standard approach is to give a single, relatively large oral dose to raise drug levels quickly so protection is in place right when it's needed. After that initial dose, smaller doses are used to maintain levels if therapy continues. This strategy works better than starting with smaller, more frequent doses, which may delay achieving effective concentrations. In practice, cephalexin is used cautiously in penicillin-allergic patients, recognizing there is some cross-reactivity risk, but it remains a common alternative when penicillin cannot be used.

7. What characterizes a nonjudgmental approach to discussing social history topics like tobacco, alcohol, and substance use?

A. Normalize disclosures about tobacco, alcohol, illicit drugs; discuss lifestyle factors nonjudgmentally.

B. Guilt and admonition.

C. Only ask about smoking.

D. Avoid discussing lifestyle.

The main idea is creating a nonjudgmental, nonstigmatizing way to take a social history about tobacco, alcohol, and substance use. When you approach these topics with neutral language and an accepting tone, patients feel safer sharing information about their use. This openness is crucial because accurate details about how substances affect health, risk behaviors, and readiness to change depend on trusting communication. Normalizing disclosures—framing tobacco, alcohol, and drugs as common topics in health care and not something to be shamed for—helps reduce fear of judgment and underreporting, enabling better assessment and management. Guilt and admonition, by contrast, can trigger defensiveness and concealment, undermining honesty. Only asking about smoking narrows the conversation and misses important context about other substances and lifestyle factors that matter for health. Avoiding discussion of lifestyle altogether leaves critical risk information undiscovered, hindering appropriate care.

SAMPLE

8. What approach is recommended when evaluating a patient with multiple health issues to improve care?

A. Focus only on the most recent problem

B. Use a structured approach, including chief concerns, review of systems, medications, and risk factors

C. Skip review of systems

D. Limit to one paragraph

When a patient has multiple health issues, gathering information in a structured, holistic way is essential. Start by identifying the chief concerns to anchor what matters most to the patient and guide priorities for the visit. Then perform a thorough review of systems to uncover symptoms the patient may not volunteer spontaneously, ensuring you don't miss problems in areas outside the most obvious complaint. Next, review every medication the patient takes—prescribed drugs, over-the-counter items, and supplements—to spot polypharmacy risks, potential drug interactions, duplications, and allergies. Finally, assess risk factors and broader context—smoking, alcohol use, nutrition, activity, social determinants, safety, and preventive needs—to shape management across all conditions. This structured approach helps integrate information from all problems, reduces the chance of missing important issues, and supports safer, more coordinated care. For example, in someone with diabetes, hypertension, and depression, this method reveals not only current symptoms but also how medications interact, what barriers to adherence may exist, and what preventive steps are needed, guiding a more effective overall plan. Focusing only on the most recent problem can miss chronic or overlapping issues; skipping the review of systems risks missing relevant symptoms; restricting the note to a single paragraph can prevent capturing the full scope needed for comprehensive care.

9. What does pack-years quantify in tobacco exposure history?

A. The current nicotine level in blood.

B. The number of times the patient has smoked in life.

C. The average cigarette nicotine content.

D. The product of packs smoked per day and years of smoking.

Pack-years quantify cumulative tobacco exposure by combining how much and how long a person has smoked. It multiplies the average number of cigarette packs smoked per day by the number of years the person has smoked, yielding a score that reflects lifetime exposure. This helps estimate risk for smoking-related diseases like COPD and lung cancer. For example, smoking one pack per day for 20 years equals 20 pack-years; smoking two packs per day for 15 years equals 30 pack-years. It is not a measure of nicotine level in the blood, nor simply a count of smoking events, nor an average nicotine content of cigarettes, but a composite that reflects both intensity and duration of smoking.

SAMPLE

10. Which food is associated with latex allergy?

A. Carrot

B. Rice

C. Avocado

D. Chicken

Latex allergy can cross-react with certain plant proteins, a phenomenon known as latex-fruit syndrome. In people with latex allergy, the immune system may recognize proteins in some fruits as if they were latex, triggering an allergic reaction after ingestion. Avocado is a classic example of a food that can cross-react with latex because it contains proteins (such as certain class I chitinases with hevein-like domains) that resemble latex allergens closely enough to provoke a reaction in susceptible individuals. The other items listed—carrot, rice, and chicken—do not commonly share these cross-reactive latex-like proteins, so they aren't typically associated with latex allergy.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://medhistorycompetency.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE