
MEDICAL SURGICAL NURSING PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://medsurgnursing.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Which imaging technique should be performed in cases of suspected orbital fractures?**

   A. X-ray of the face

   B. Ultrasound of the eye

   C. CT scan or MRI of the head

   D. Standard eye examination

2. **Which eye condition can lead to transient proptosis?**

   A. Orbital cellulitis

   B. Glaucoma

   C. Dry eye syndrome

   D. Cataracts

3. **Which symptom is associated with visual field defect over time in retinal detachment?**

   A. Complete blindness

   B. Gradual reduction in visual acuity

   C. Improved night vision

   D. Brightness perception increases

4. **What management step is essential for orbital cellulitis diagnosis?**

   A. CT scan

   B. Systemic antibiotics

   C. Blood tests

   D. Drainage of the abscess

5. **Which of the following findings is NOT associated with diabetic retinopathy?**

   A. Retinal neovascularization

   B. Blot haemorrhages in one quadrant

   C. Postpartum retinal detachment

   D. Cotton wool spots

SAMPLE

6. Which medication is classified as a second-line treatment for allergic conjunctivitis?

A. Oral antihistamines

B. Sodium cromoglycate

C. Opatanol

D. Topical corticosteroids

7. When is chemotherapy indicated for patients with retinoblastoma?

A. After surgical removal of the tumor

B. For intraocular groups C and D

C. When the patient is asymptomatic

D. As a first-line treatment for all cases

8. What test helps detect abnormalities associated with sarcoidosis in suspected uveitis patients?

A. Liver function tests

B. Chest radiography

C. Thyroid function tests

D. Electrocardiogram

9. What is the most common management step for pre-septal cellulitis?

A. Confirmation of diagnosis

B. CT scan

C. Admission for observation

D. Systemic antibiotics

10. What is one effect of vascular endothelial growth factor (VEGF) in diabetic retinopathy?

A. It increases blood flow to the retina

B. It promotes the formation of new blood vessels

C. It reduces retinal swelling

D. It stabilizes existing retinal vessels

SAMPLE

Answers

SAMPLE

1. C
2. A
3. B
4. A
5. C
6. C
7. B
8. B
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. Which imaging technique should be performed in cases of suspected orbital fractures?

A. X-ray of the face

B. Ultrasound of the eye

C. CT scan or MRI of the head

D. Standard eye examination

When evaluating suspected orbital fractures, a CT scan or MRI of the head is the preferred imaging technique due to its ability to provide detailed cross-sectional images of the bony structures, soft tissues, and any potential intracranial involvement. CT scanning is especially valuable in emergency settings for its rapid acquisition of images, allowing for immediate assessment of complex fractures and any associated injuries to the orbit and surrounding areas. It can reveal displaced bones, a presence of hematomas, and other complications that require prompt medical intervention. MRI, while not typically the first line for fractures, can provide excellent soft tissue detail and is useful in cases where there is concern for damage to the ocular muscles or nerves. However, the speed and accessibility of CT make it the go-to modality in trauma settings involving the orbit. In summary, the use of a CT scan or MRI is crucial in effectively diagnosing orbital fractures and guiding subsequent management, highlighting its importance in clinical practice.

2. Which eye condition can lead to transient proptosis?

A. Orbital cellulitis

B. Glaucoma

C. Dry eye syndrome

D. Cataracts

Transient proptosis, or the temporary protrusion of the eyeball, is often associated with conditions that involve inflammation or increased pressure within the orbit. Orbital cellulitis, an infection of the orbital soft tissues, can lead to swelling and inflammation, resulting in the displacement of the eyeball. This condition is characterized by symptoms such as pain, swelling, redness, and sometimes fever, which are indicative of an acute inflammatory process that can cause the eye to protrude. In contrast, conditions like glaucoma, dry eye syndrome, and cataracts do not typically present with transient proptosis. Glaucoma involves increased intraocular pressure but does not cause the eye to protrude; instead, it may lead to vision loss if left untreated. Dry eye syndrome primarily affects tear production and eye lubrication but also does not result in proptosis. Cataracts are related to lens opacification and do not influence the position of the eyeball within the orbit. Therefore, orbital cellulitis is correctly identified as a condition that can result in transient proptosis due to the inflammatory processes involved.

SAMPLE

3. Which symptom is associated with visual field defect over time in retinal detachment?

A. Complete blindness

B. Gradual reduction in visual acuity

C. Improved night vision

D. Brightness perception increases

A gradual reduction in visual acuity is indeed a symptom associated with visual field defects over time in retinal detachment. As the condition progresses, the retina becomes more compromised, leading to a decline in the clarity of the image that reaches the brain. This results in difficulty seeing details and may worsen as the detachment spreads. Unlike complete blindness, which can occur in advanced cases, the gradual reduction in visual acuity manifests earlier and is a key indicator that patients should seek prompt medical evaluation. Other options, such as improved night vision or increased brightness perception, do not align with the typical progression of retinal detachment, as these conditions usually become worse rather than better in the presence of this issue. Understanding these symptoms can help both healthcare providers and patients recognize the need for intervention to mitigate the risks associated with retinal detachment.

4. What management step is essential for orbital cellulitis diagnosis?

A. CT scan

B. Systemic antibiotics

C. Blood tests

D. Drainage of the abscess

For diagnosing orbital cellulitis, performing a CT scan is essential as it allows for an accurate assessment of the extent of the infection and any possible complications, such as the presence of an abscess or involvement of the surrounding structures, including the eye socket and sinuses. The CT scan provides a detailed image that can help differentiate orbital cellulitis from other conditions that present with similar symptoms, such as preseptal cellulitis or abscess formation. While systemic antibiotics are crucial in the management of orbital cellulitis, they are typically initiated after the diagnosis is confirmed. Blood tests may offer additional information regarding infection but are not definitive for diagnosis. Drainage of an abscess may be necessary in cases where a localized collection of pus is identified, but obtaining imaging through a CT scan is the critical first step in confirming the diagnosis and determining the appropriate management plan.

SAMPLE

5. Which of the following findings is NOT associated with diabetic retinopathy?

A. Retinal neovascularization

B. Blot haemorrhages in one quadrant

C. Postpartum retinal detachment

D. Cotton wool spots

Diabetic retinopathy is a complication of diabetes that primarily affects the blood vessels in the retina, leading to various characteristic findings. Among the choices provided, postpartum retinal detachment is not a finding typically associated with diabetic retinopathy. Retinal neovascularization, which occurs in response to ischemia due to damaged retinal blood vessels, is a hallmark of proliferative diabetic retinopathy. It involves the growth of new, fragile blood vessels that can lead to further complications such as hemorrhage or retinal detachment. Blot hemorrhages are another common finding in diabetic retinopathy. They result from broken blood vessels leaking blood into the deeper layers of the retina and can be observed in multiple quadrants of the retina, particularly in cases of non-proliferative diabetic retinopathy. Cotton wool spots are also associated with diabetic retinopathy. They represent localized areas of ischemia and are formed from accumulated axoplasmic material within the nerve fiber layer of the retina, typically seen in the context of poor glucose control and retinal vascular changes. In contrast, postpartum retinal detachment is not a classic manifestation of diabetic retinopathy and relates more to other risk factors, such as changes associated with pregnancy, rather than direct consequences of diabetes and

6. Which medication is classified as a second-line treatment for allergic conjunctivitis?

A. Oral antihistamines

B. Sodium cromoglycate

C. Opatanol

D. Topical corticosteroids

Opatanol is correctly identified as a second-line treatment for allergic conjunctivitis. It contains the active ingredient olopatadine, which is an antihistamine that helps relieve symptoms such as itching and redness in the eyes. While it can be effective for the management of allergic conjunctivitis, it is usually considered after first-line treatments—primarily oral antihistamines or topical mast cell stabilizers like sodium cromoglycate—do not provide adequate relief or when the condition is more persistent or severe. Topical corticosteroids, while effective in managing inflammation, are generally reserved for more severe cases of allergic conjunctivitis due to their potential side effects with prolonged use, such as elevated intraocular pressure or cataract formation. Therefore, Opatanol serves as a useful option when first-line therapies do not suffice, striking a balance between effectiveness and safety for mild to moderate cases of allergic conjunctivitis.

SAMPLE

7. When is chemotherapy indicated for patients with retinoblastoma?

A. After surgical removal of the tumor

B. For intraocular groups C and D

C. When the patient is asymptomatic

D. As a first-line treatment for all cases

Chemotherapy is indicated for patients with retinoblastoma primarily for intraocular groups C and D because these classifications indicate a more advanced disease state where the tumors are large or have spread within the eye. In these cases, chemotherapy can help reduce the size of the tumors or facilitate other treatment options, such as laser therapy or cryotherapy, ultimately aiming to preserve vision and control the disease. Intraocular groups C and D indicate that treatment is necessary to effectively manage the disease, as lump sizes and potential invasion of surrounding tissues increase the risk of complications. Chemotherapy works by targeting rapidly dividing cancer cells, making it a useful modality to shrink tumors before other interventions or to treat aggressive disease. Other potential treatment options, such as surgical removal of the tumor, might not be directly applicable in the earlier stages represented by intraocular groups, or surgeries might not achieve the same outcomes when lesions are still present or when there is a significant risk of metastasis. The other options either suggest interventions inappropriate for the disease stage or imply a lack of treatment when chemotherapy could be beneficial. Thus, for groups C and D, utilizing chemotherapy is essential to optimize patient care and outcomes in retinoblastoma.

8. What test helps detect abnormalities associated with sarcoidosis in suspected uveitis patients?

A. Liver function tests

B. Chest radiography

C. Thyroid function tests

D. Electrocardiogram

Chest radiography is particularly useful for detecting abnormalities associated with sarcoidosis in patients suspected of having uveitis. Sarcoidosis is a systemic granulomatous disease that can affect multiple organs, including the eyes. When uveitis is suspected, imaging of the chest is valuable because pulmonary involvement is common in sarcoidosis. In a chest X-ray, one might see hilar lymphadenopathy or pulmonary infiltrates, which are classic signs of sarcoidosis. If these findings are present, they strengthen the suspicion of sarcoidosis as an underlying cause of uveitis. Therefore, chest radiography serves as an important diagnostic tool to evaluate for this condition, guiding subsequent management and treatment. Other tests, while useful in other contexts, do not specifically target the abnormalities associated with sarcoidosis in the same way that chest radiography does.

SAMPLE

9. What is the most common management step for pre-septal cellulitis?

A. Confirmation of diagnosis

B. CT scan

C. Admission for observation

D. Systemic antibiotics

In the context of managing pre-septal cellulitis, the most common initial management step is the confirmation of diagnosis. Establishing a precise diagnosis is crucial because pre-septal cellulitis can often have similar presentations to other conditions such as orbital cellulitis, which is more severe and requires different management strategies. Confirming the diagnosis may involve a thorough clinical evaluation, including a detailed history and physical examination to assess symptoms like redness, swelling, and pain around the eye. By accurately diagnosing pre-septal cellulitis, healthcare providers can ensure that patients receive appropriate treatment and avoid unnecessary interventions that might arise from misdiagnosis. While imaging studies like a CT scan may be utilized in certain cases, particularly if there is suspicion for orbital involvement or complications, it is not routinely performed for straightforward cases of pre-septal cellulitis. Admission for observation is generally not the standard management unless there are complications or severe cases requiring careful monitoring. The use of systemic antibiotics is an important step in treatment, but it is typically initiated once the diagnosis is confirmed rather than as the first step.

10. What is one effect of vascular endothelial growth factor (VEGF) in diabetic retinopathy?

A. It increases blood flow to the retina

B. It promotes the formation of new blood vessels

C. It reduces retinal swelling

D. It stabilizes existing retinal vessels

Vascular endothelial growth factor (VEGF) plays a significant role in diabetic retinopathy, particularly through its promotion of angiogenesis, which is the formation of new blood vessels. In the context of diabetic retinopathy, elevated levels of VEGF are often associated with hypoxia or inadequate blood supply to the retina due to damaged vasculature resulting from diabetes. When VEGF is secreted, it stimulates endothelial cells to proliferate and form new blood vessels, a process aimed at compensating for the poor oxygen and nutrient delivery to retinal tissues. While this may initially seem beneficial, the new vessels formed are often fragile and can lead to complications such as leakage, bleeding, and further damage to the retinal structure, contributing to vision loss. This underscores the dual role of VEGF in promoting necessary vascularization while also playing a part in pathological conditions when dysregulated. The other options do not accurately reflect the primary role of VEGF in this condition. While increased blood flow may occur as a result of new vessel formation, it's not the direct effect of VEGF itself. Additionally, VEGF is associated with retinal swelling and does not stabilize existing blood vessels, making it clear why the promotion of new blood vessels is the most relevant effect of VEG

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://medsurgnursing.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE