MODERATE-SEVERE TRAUMATIC BRAIN INJURY (TBI) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://moderateseveretbi.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is the overall goal of ICP management in moderate-to-severe TBI?

A. To prevent secondary brain injury by controlling ICP and maintaining adequate CPP using a stepwise approach of medical and surgical therapies as indicated

B. To maintain ICP at the highest possible level to ensure perfusion

C. To avoid any use of osmotic therapies

D. To eliminate all sedation during management

2. Before initiating pharmacologic DVT prophylaxis in traumatic brain injury, what condition should be confirmed?

A. It is safe to administer (no active bleeding and stable ICP)

B. The patient has fever

C. The patient has normal glucose

D. The patient has elevated blood pressure

3. An External Ventricular Drain is placed into which structure to monitor ICP and drain CSF?

A. Lateral ventricle.

B. Third ventricle.

C. Subarachnoid space.

D. Cerebellar hemisphere.

4. The Rancho Los Amigos Cognitive Scale is used for what purpose?

A. Predicts mortality risk.

B. Classifies cognitive functioning and recovery level after brain injury.

C. Measures blood glucose control.

D. Assesses physical mobility.

5. Which hyponatremia patterns can occur after TBI?

A. SIADH or cerebral salt wasting

B. Hyperkalemia

C. Hypernatremia from dehydration

D. Hyperglycemia

SAMPLE

6. Which statement differentiates cytotoxic edema from vasogenic edema on imaging?

A. Vasogenic edema is extracellular fluid from BBB disruption with T2/FLAIR hyperintensity

B. Enhanced gadolinium uptake

C. Cytotoxic edema is cellular swelling with restricted diffusion on DWI

D. Hyperintense on DWI and T2 equally

7. What is the primary goal of initial resuscitation in moderate-severe TBI?

A. Stabilize airway, breathing, circulation with prevention of hypoxia and hypotension and optimization of cerebral perfusion pressure

B. Achieve normoglycemia

C. Initiate early DVT prophylaxis

D. Begin antibiotics prophylactically

8. How does anemia affect outcomes after traumatic brain injury?

A. Has no effect

B. Reduces oxygen delivery to brain; may worsen outcomes; transfusion considered per protocol

C. Improves oxygen delivery

D. Increases blood viscosity in beneficial way

9. In the context of moderate-severe TBI, what is the purpose of maintaining a CPP target?

A. To ensure adequate cerebral blood flow and oxygen delivery to the brain

B. To optimize systemic vascular resistance

C. To minimize heart rate

D. To maximize CSF production

10. The key components of secondary injury prevention in TBI include preventing which insults?

A. Hypoxia, hypotension, fever, edema, seizures

B. Hyperglycemia

C. Hypothermia

D. None of the above

SAMPLE

Answers

SAMPLE

1. A
2. A
3. A
4. B
5. A
6. C
7. A
8. B
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. What is the overall goal of ICP management in moderate-to-severe TBI?

A. To prevent secondary brain injury by controlling ICP and maintaining adequate CPP using a stepwise approach of medical and surgical therapies as indicated

B. To maintain ICP at the highest possible level to ensure perfusion

C. To avoid any use of osmotic therapies

D. To eliminate all sedation during management

The goal of ICP management in moderate-to-severe TBI is to prevent secondary brain injury by keeping intracranial pressure under control and ensuring adequate cerebral perfusion pressure through a stepwise medical and surgical plan. When ICP rises, cerebral perfusion pressure (CPP = MAP − ICP) falls, which can lead to brain ischemia and further injury. So the priority is to lower or stabilize ICP while maintaining CPP in a range that supports consistent brain blood flow, typically around 60–70 mmHg in adults. Start with less invasive measures—adequate oxygenation and ventilation, appropriate sedation and analgesia, head elevation, and meticulous hemodynamic control—and escalate to medical therapies such as hyperosmolar therapy (hypertonic saline or mannitol) and CSF drainage when indicated. If ICP remains elevated despite these steps, add temporizing measures like controlled hyperventilation and, in refractory cases, consider surgical options such as decompressive craniectomy. The emphasis is on preventing secondary injury by preserving perfusion, not on keeping ICP high. Osmotic therapy and thoughtful sedation are integral parts of this approach, while avoiding them would undermine proper ICP management.

2. Before initiating pharmacologic DVT prophylaxis in traumatic brain injury, what condition should be confirmed?

A. It is safe to administer (no active bleeding and stable ICP)

B. The patient has fever

C. The patient has normal glucose

D. The patient has elevated blood pressure

Before starting pharmacologic DVT prophylaxis in a traumatic brain injury patient, you must confirm there is no active intracranial bleeding and that intracranial pressure is stable. Anticoagulants can worsen an existing bleed or cause hematoma expansion, leading to worse outcomes. If imaging shows ongoing bleeding or ICP is unstable, delaying pharmacologic prophylaxis or using alternative, non-pharmacologic methods is safer. Fever, normal glucose, or elevated blood pressure by themselves do not determine the safety of starting prophylaxis; they require separate management, but do not replace the need to verify that there is no active bleed and that ICP is controlled.

SAMPLE

3. An External Ventricular Drain is placed into which structure to monitor ICP and drain CSF?

A. Lateral ventricle.

B. Third ventricle.

C. Subarachnoid space.

D. Cerebellar hemisphere.

The key idea is that an external ventricular drain provides both continuous ICP monitoring and CSF drainage by entering the ventricular system. The lateral ventricle is the usual entry point because it offers direct access to CSF and allows the catheter tip to sit in the frontal horn near the foramen of Monro, giving an accurate intraventricular pressure reading and a reliable drainage path. Draining from the ventricular system makes the ICP measurement reflect true intracranial pressure and enables effective CSF removal to prevent or treat hydrocephalus. Placing the catheter in the third ventricle is technically possible but much less common and more risky; the subarachnoid space drains CSF but does not provide as precise, reliable ICP monitoring with an external drain; the cerebellar hemisphere is not part of the ventricular CSF pathway and isn't used for EVD.

4. The Rancho Los Amigos Cognitive Scale is used for what purpose?

A. Predicts mortality risk.

B. Classifies cognitive functioning and recovery level after brain injury.

C. Measures blood glucose control.

D. Assesses physical mobility.

The Rancho Los Amigos Cognitive Scale is used to classify cognitive functioning and recovery level after brain injury. It describes a patient's cognitive and behavioral status as they recover, guiding rehabilitation planning and prognosis. The scale focuses on how brain injury affects attention, memory, problem solving, orientation, initiation, and self-monitoring, and it tracks progress from early levels of reduced responsiveness to higher levels of purposeful, independent activity. It helps therapists set realistic goals, determine therapy intensity, and communicate a patient's current cognitive abilities to the care team and family. This is different from mortality risk, blood glucose control, or physical mobility measures, which are assessed with other tools.

5. Which hyponatremia patterns can occur after TBI?

A. SIADH or cerebral salt wasting

B. Hyperkalemia

C. Hypernatremia from dehydration

D. Hyperglycemia

When the brain is injured, its control of fluid balance can go awry, leading to hyponatremia from two main patterns: SIADH and cerebral salt wasting. SIADH is a dilutional hyponatremia caused by excessive release of antidiuretic hormone, which makes the body retain water. This results in low serum sodium but with normal or near-normal blood volume (euvolemia) and urine that is inappropriately concentrated with high sodium excretion. Cerebral salt wasting, on the other hand, involves true loss of salt in the urine plus accompanying water loss, leading to hyponatremia with hypovolemia. Here the kidneys waste sodium, and the patient becomes volume depleted despite the low sodium in the blood. These two patterns are both hyponatremia after brain injury, but they differ in volume status and the underlying mechanism, which is crucial for treatment: SIADH usually calls for fluid restriction, while cerebral salt wasting requires salt and fluid replacement to correct both sodium and volume deficits. Hyperkalemia, hypernatremia from dehydration, and hyperglycemia describe other electrolyte or metabolic issues and do not characterize the hyponatremia patterns seen after TBI.

SAMPLE

6. Which statement differentiates cytotoxic edema from vasogenic edema on imaging?

A. Vasogenic edema is extracellular fluid from BBB disruption with T2/FLAIR hyperintensity

B. Enhanced gadolinium uptake

C. Cytotoxic edema is cellular swelling with restricted diffusion on DWI

D. Hyperintense on DWI and T2 equally

The key idea is diffusion imaging: cytotoxic edema causes cells to swell and water to become trapped inside cells, which restricts how freely water molecules can move. On MRI this shows up as bright signal on diffusion-weighted imaging (DWI) with a corresponding low signal on the ADC map, indicating restricted diffusion. This pattern is typical of cellular swelling from energy failure, often seen in acute ischemia or injury. In contrast, vasogenic edema involves fluid accumulating outside cells due to disruption of the blood-brain barrier, increasing extracellular space. Water diffusion is not restricted; the ADC map shows higher values and the edema often appears as T2/FLAIR hyperintensity. Gadolinium enhancement may occur with BBB breakdown, supporting vasogenic processes, but diffusion characteristics are the distinguishing feature. So, the statement that cytotoxic edema is cellular swelling with restricted diffusion on DWI best captures the differentiating imaging feature.

7. What is the primary goal of initial resuscitation in moderate-severe TBI?

A. Stabilize airway, breathing, circulation with prevention of hypoxia and hypotension and optimization of cerebral perfusion pressure

B. Achieve normoglycemia

C. Initiate early DVT prophylaxis

D. Begin antibiotics prophylactically

The priority in initial resuscitation for moderate-severe TBI is to prevent secondary brain injury by ensuring the brain receives enough oxygen and blood flow. That means securing the airway and ensuring adequate ventilation and oxygenation to avoid hypoxia, and maintaining circulation to prevent hypotension. By keeping oxygen delivery high and blood pressure sufficient, you protect cerebral perfusion pressure (CPP), which is the driving force for blood flow to the injured brain. CPP is roughly the difference between mean arterial pressure and intracranial pressure, so if you keep MAP up and ICP in check, you support the brain's perfusion during the critical early phase. While managing glucose, infections, or clot prevention is important, they are not the main focus of the initial resuscitation itself.

SAMPLE

8. How does anemia affect outcomes after traumatic brain injury?

A. Has no effect

B. Reduces oxygen delivery to brain; may worsen outcomes; transfusion considered per protocol

C. Improves oxygen delivery

D. Increases blood viscosity in beneficial way

Anemia reduces the brain's oxygen delivery, which matters after traumatic brain injury because the injured brain is especially vulnerable to hypoxia. Oxygen delivery to tissue depends on how much oxygen the blood can carry (largely driven by hemoglobin) and how much blood is flowing. When hemoglobin is low, CaO2—the amount of oxygen carried per volume of blood—decreases, so even with good oxygen saturation, the brain receives less O2. In the setting of moderate to severe TBI, metabolic demands rise and the brain's ability to compensate for reduced delivery is limited, so lower oxygen delivery can worsen secondary brain injury and worsen outcomes. That's why transfusion is considered within protocols to maintain adequate oxygen-carrying capacity and support brain oxygenation. The other statements aren't accurate: anemia doesn't improve oxygen delivery, and it doesn't confer a beneficial increase in blood viscosity; in fact, viscosity tends to be lower with anemia, and that isn't protective in this context.

9. In the context of moderate-severe TBI, what is the purpose of maintaining a CPP target?

A. To ensure adequate cerebral blood flow and oxygen delivery to the brain

B. To optimize systemic vascular resistance

C. To minimize heart rate

D. To maximize CSF production

The purpose of a CPP target is to ensure adequate cerebral blood flow and oxygen delivery to the brain when intracranial pressure is elevated after a moderate to severe TBI. CPP represents the driving pressure for blood entering the brain, calculated roughly as mean arterial pressure minus intracranial pressure. If CPP falls too low, cerebral blood flow decreases and brain tissue becomes ischemic, fueling secondary injury. Therefore, the goal is to keep CPP in a range that maintains sufficient cerebral perfusion, often around 60–70 mmHg, individualized to the patient. This is achieved by supporting MAP and actively managing ICP through fluid and vasopressor strategies, osmotic therapies, or drainage as appropriate. The other options don't address perfusion of brain tissue: systemic vascular resistance, heart rate, and CSF production aren't the main levers for ensuring adequate brain blood flow and oxygen delivery.

SAMPLE

10. The key components of secondary injury prevention in TBI include preventing which insults?

A. Hypoxia, hypotension, fever, edema, seizures

B. Hyperglycemia

C. Hypothermia

D. None of the above

Secondary injury after TBI is driven by events that occur after the initial insult, and preventing those insults helps limit further brain damage. The most important modifiable insults to prevent are hypoxia, which deprives the brain of oxygen and can worsen injury when blood flow is already compromised; hypotension, which lowers cerebral perfusion pressure and risks ischemia; fever, which raises metabolic demands and can amplify injury; edema, which increases intracranial pressure and reduces cerebral blood flow; and seizures, which cause excessive metabolic stress and excitotoxic injury. These are the common targets clinicians focus on to protect brain tissue in the acute period. While hyperglycemia and other factors are relevant to outcome, the set listed here represents the familiar bundle of secondary insults targeted for prevention, which is why this option is the best fit.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://moderateseveretbi.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE