

MPAS COMMON LABORATORY TESTS: BLOOD PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS



**VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE**

<https://mpascommonlabtestblood.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

- 1. Leukopenia is defined as which of the following?**
 - A. Increased WBC count
 - B. Normal WBC count
 - C. Decreased WBC count
 - D. Increased RBC count

- 2. Why is fasting recommended for lipid testing, and what is the typical impact of nonfasting on triglyceride levels?**
 - A. To prevent glucose fluctuations; nonfasting triglycerides rise dramatically and invalidate results.
 - B. To reduce postprandial variation; nonfasting triglycerides rise modestly.
 - C. To increase test sensitivity; nonfasting lowers triglycerides.
 - D. Fasting has no impact on accuracy; nonfasting always acceptable.

- 3. Increased amylase is most commonly associated with which condition?**
 - A. Renal failure
 - B. Pancreatitis
 - C. Hepatitis
 - D. Hyperglycemia

- 4. What roles do the BMP play in assessing kidney function and electrolyte balance?**
 - A. Evaluates electrolytes (Na, K, Cl, HCO₃), renal function (BUN, creatinine), and glucose; helps assess acid-base and hydration.
 - B. Measures liver enzymes exclusively.
 - C. Only tests for blood sugar.
 - D. Tests lipid panel.

- 5. What do rising BUN and creatinine indicate and how can you interpret the BUN/creatinine ratio?**
 - A. Indicate reduced kidney function; a high ratio (>20:1) may suggest prerenal azotemia, while a normal/low ratio with elevated creatinine may suggest intrinsic renal disease.
 - B. Indicate liver disease.
 - C. Ratio is irrelevant.
 - D. A low ratio indicates prerenal azotemia.

- 6. Relative to central lab testing, POCT is commonly characterized by being...**
- A. Significantly More Accurate
 - B. Significantly Less Accurate
 - C. Slightly Less Accurate
 - D. Equally Accurate
- 7. Increased alkaline phosphatase levels can be seen during which physiological process?**
- A. Growth
 - B. Liver injury
 - C. Hemolysis
 - D. Dehydration
- 8. Which set lists the components of a BMP?**
- A. Hematology panel
 - B. Sodium, Potassium, Chloride, Bicarbonate, plus BUN, Creatinine, Glucose
 - C. Na, K, Cl, Bicarb, BUN, Creatinine, Glucose, calcium
 - D. Sodium, Potassium, Magnesium, Phosphate
- 9. What does POCT stand for?**
- A. Point of Care Testing
 - B. Point of Contact Test
 - C. Point of Central Therapy
 - D. Point of Clinical Trial
- 10. Why is ferritin considered a marker of iron stores and what is its limitation?**
- A. Ferritin correlates with iron stores, but can be elevated with inflammation or infection, masking iron deficiency
 - B. Ferritin directly measures bone marrow iron
 - C. Ferritin is unaffected by inflammation
 - D. Ferritin decreases with iron deficiency regardless of stores

Answers

SAMPLE

1. C
2. C
3. B
4. A
5. A
6. C
7. A
8. C
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. Leukopenia is defined as which of the following?

- A. Increased WBC count
- B. Normal WBC count
- C. Decreased WBC count**
- D. Increased RBC count

Leukopenia means a low white blood cell count. White blood cells defend the body against infection, so if their number drops below the normal range, infection risk increases and this finding is described as leukopenia. In adults, a typical normal WBC count is roughly 4,000 to 11,000 cells per microliter; counts below this range define leukopenia. The other options describe different situations: a higher WBC count is leukocytosis, a normal count isn't leukopenia, and an increased red blood cell count relates to a different parameter altogether.

2. Why is fasting recommended for lipid testing, and what is the typical impact of nonfasting on triglyceride levels?

- A. To prevent glucose fluctuations; nonfasting triglycerides rise dramatically and invalidate results.
- B. To reduce postprandial variation; nonfasting triglycerides rise modestly.
- C. To increase test sensitivity; nonfasting lowers triglycerides.**
- D. Fasting has no impact on accuracy; nonfasting always acceptable.

When measuring a lipid panel, triglycerides are highly influenced by recent meals because fat absorbed from digestion enters the blood as chylomicrons, causing a postprandial spike. Fasting removes this meal-related variation, giving a stable baseline that makes results more comparable and interpretations more reliable, especially for calculating LDL cholesterol with the Friedewald equation. If a nonfasting sample is used, triglycerides typically rise after eating, but the increase is usually modest rather than dramatic. This post-meal rise can still affect interpretation and is why fasting is the standard practice for lipid testing.

3. Increased amylase is most commonly associated with which condition?

- A. Renal failure
- B. Pancreatitis**
- C. Hepatitis
- D. Hyperglycemia

Amylase elevation is most characteristically tied to pancreatic inflammation. When the pancreas is inflamed, its acinar cells release large amounts of amylase into the blood (and urine), so serum amylase rises early in the course of pancreatitis, often within a few hours and peaking around the first day or so. This makes pancreatitis the classic condition associated with increased amylase. Renal failure can also show higher amylase levels because the kidneys aren't clearing it efficiently, but that isn't the typical or defining association. Hepatitis and hyperglycemia don't usually produce a notable rise in amylase.

4. What roles do the BMP play in assessing kidney function and electrolyte balance?

- A. Evaluates electrolytes (Na, K, Cl, HCO₃), renal function (BUN, creatinine), and glucose; helps assess acid-base and hydration.**
- B. Measures liver enzymes exclusively.
- C. Only tests for blood sugar.
- D. Tests lipid panel.

The basic metabolic panel is focused on the body's chemical balance and waste filtration, giving a snapshot of kidney function and electrolyte status. It measures key electrolytes—sodium, potassium, chloride, and bicarbonate—which are essential for fluid balance, nerve and muscle function, and overall cellular operations. It also includes kidney-function markers—blood urea nitrogen and creatinine—that reflect how well the kidneys are filtering waste from the blood. In many panels, glucose is included as part of the metabolic assessment, providing a quick look at blood sugar status. The bicarbonate level helps gauge acid-base balance, while the BUN/creatinine relationship can give clues about hydration status. Taken together, these components allow you to assess electrolyte balance, kidney function, acid-base status, and hydration—exactly what the basic metabolic panel is designed to evaluate. The other choices focus on liver enzymes, blood sugar alone, or lipid levels, which are not the primary purposes of the BMP.

5. What do rising BUN and creatinine indicate and how can you interpret the BUN/creatinine ratio?

- A. Indicate reduced kidney function; a high ratio (>20:1) may suggest prerenal azotemia, while a normal/low ratio with elevated creatinine may suggest intrinsic renal disease.**
- B. Indicate liver disease.
- C. Ratio is irrelevant.
- D. A low ratio indicates prerenal azotemia.

Rising BUN and creatinine mean the kidneys aren't filtering waste effectively, signaling reduced kidney function and a lower glomerular filtration rate. The BUN/creatinine ratio helps distinguish where the problem lies. A high ratio (greater than about 20:1) points to prerenal azotemia, where decreased renal perfusion or volume status (like dehydration or heart failure) causes the kidney to reabsorb more urea, so BUN climbs disproportionately to creatinine. When both BUN and creatinine rise but the ratio stays normal or low, this suggests intrinsic renal disease (such as acute tubular injury), where the kidney's ability to reabsorb BUN is impaired and the two wastes rise more in parallel. Liver disease can affect BUN production, but rising BUN and creatinine together mainly indicate kidney function issues, not liver pathology. The ratio isn't irrelevant; it provides useful context but isn't a definitive diagnosis on its own.

6. Relative to central lab testing, POCT is commonly characterized by being...

- A. Significantly More Accurate
- B. Significantly Less Accurate
- C. Slightly Less Accurate
- D. Equally Accurate

Point-of-care testing prioritizes rapid results at the patient's side, which often comes with a small trade-off in analytical precision compared with central laboratory testing. Central labs use high-volume, automated analyzers with stringent quality control, calibration, and specially trained staff, yielding high accuracy and reproducibility. Point-of-care devices are typically user-operated and rely on quicker, closer-to-bedside workflows, with samples like capillary blood and more variable pre-analytical handling; calibration and quality checks can be less rigorous, and operator technique can influence results. These factors tend to make POCT slightly less accurate overall, though the results are usually adequate for making timely clinical decisions. There are tests where POCT performance is very close to central labs, but the common characterization is a slight decrease in accuracy.

7. Increased alkaline phosphatase levels can be seen during which physiological process?

- A. Growth
- B. Liver injury
- C. Hemolysis
- D. Dehydration

Alkaline phosphatase rises when bone formation is active. During growth, especially in children and adolescents, bones are rapidly forming as the skeleton lengthens. Osteoblasts produce more ALP to aid mineralization, and this enzyme spills into the bloodstream, elevating serum levels. That normal, increased bone turnover during growth is why ALP is higher in growing individuals. Elevations from liver injury can also occur, but that reflects a pathological process with cholestasis and other liver enzymes involved, not a normal growth-related change. Hemolysis and dehydration don't specifically drive an increase in ALP in the way active bone formation does.

8. Which set lists the components of a BMP?

- A. Hematology panel
- B. Sodium, Potassium, Chloride, Bicarbonate, plus BUN, Creatinine, Glucose
- C. Na, K, Cl, Bicarb, BUN, Creatinine, Glucose, calcium
- D. Sodium, Potassium, Magnesium, Phosphate

The basic metabolic panel measures the body's key electrolytes and metabolic byproducts to assess fluid balance, kidney function, and energy status. The core components include sodium, potassium, chloride, bicarbonate, plus the kidney function markers BUN and creatinine, and a glucose level. Calcium is often included in BMPs depending on the lab, so a set that lists Na, K, Cl, bicarbonate, BUN, creatinine, glucose, and calcium aligns with how many panels are reported. The other options either describe a different type of panel (hematology) or omit calcium, or add minerals not typically part of a BMP (like magnesium and phosphate), making them not representative of the BMP as commonly defined.

9. What does POCT stand for?

- A. Point of Care Testing**
- B. Point of Contact Test
- C. Point of Central Therapy
- D. Point of Clinical Trial

POCT stands for testing done at or near the patient, not in a central laboratory. This approach brings the test results quickly to the point of care, enabling faster clinical decisions and often improving patient flow in clinics, emergency departments, or during home health visits. Examples include bedside blood glucose meters, rapid infectious disease tests, and portable coagulation testing for anticoagulation management. The main advantage is speed, but it requires solid quality control and proper operator training to ensure accuracy. The other options aren't standard terms used in laboratory medicine, so they don't describe this practical testing approach.

10. Why is ferritin considered a marker of iron stores and what is its limitation?

- A. Ferritin correlates with iron stores, but can be elevated with inflammation or infection, masking iron deficiency**
- B. Ferritin directly measures bone marrow iron
- C. Ferritin is unaffected by inflammation
- D. Ferritin decreases with iron deficiency regardless of stores

Ferritin is the storage form of iron in the body, so serum ferritin generally mirrors the amount of iron stored in tissues. When iron stores are low, ferritin levels fall; when stores are adequate or high, ferritin tends to be higher. The big limitation is that ferritin is an acute-phase reactant—its levels rise in the presence of inflammation or infection regardless of how much iron is actually stored. This means someone could have depleted iron stores but normal or even elevated ferritin if they have an inflammatory state, masking iron deficiency. In practice, ferritin has to be interpreted with inflammation markers (like CRP) and often alongside other iron tests, such as transferrin saturation or soluble transferrin receptor, to get a clearer picture of iron status.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://mpascommonlabtestblood.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE