MPS: VENIPUNCTURE, INJECTIONS, IVS, AND ABGS PROCEDURES PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://mpsvenipuncture.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1
Table of Contents 2
Introduction 3
How to Use This Guide 4
Questions 5
Answers 8
Explanations 10
Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What are the primary adult anatomical sites for intramuscular injections?

A. Anterior thighs, upper outer quadrant of buttocks, and deltoid.

B. Anterior thigh only.

C. Dorsogluteal muscle exclusively.

D. Vastus lateralis and gluteus medius only.

2. If Staphylococcus epidermidis is detected in only one of several blood culture bottles, what is the most likely interpretation?

A. Contamination from skin flora

B. Definitive bacteremia with this organism

C. Contamination across all bottles

D. Laboratory error with sample mislabeling

3. In a patient with a left mastectomy and a functioning IV in the opposite arm, what is the correct approach to blood draw?

A. Turn off IV for 2 minutes, draw distally to IV, and discard the first 5 mL

B. Draw from the mastectomy side

C. Draw from the arm with the IV without modification

D. Use a leg vein

4. Which statement describes a contraindication to venipuncture due to venous fibrosis?

A. Venous fibrosis.

B. Varicose veins.

C. Small veins.

D. Normal skin.

5. Capillary puncture is typically performed at which sites?

A. Fingerstick or heel.

B. Arm vein.

C. Neck vein.

D. Ankle.

SAMPLE

6. Why is it important to keep the venipuncture site clean and dry after collection?

A. To reduce infection risk and promote hemostasis.

B. To speed up blood flow into the vein.

C. To prevent the patient from feeling faint.

D. To avoid contrast in future testing.

7. Which specialized needle device is recommended when performing venipuncture on hand veins or fragile veins?

A. A 25-gauge winged butterfly device.

B. A 21-gauge winged butterfly device.

C. A 23-gauge winged butterfly device.

D. A 23-gauge straight needle.

8. What constitutes a complete blood culture bottle collection kit?

A. One aerobic culture bottle and one anaerobic culture bottle.

B. Two aerobic culture bottles.

C. One anaerobic culture bottle only.

D. A bottle that contains both aerobic and anaerobic media.

9. What is a recommended prevention for phlebitis?

A. Aseptic technique and site rotation

B. Frequent IV flush

C. Avoid cleaning around site

D. Increase IV rate

10. What are the typical needle gauge and length for subcutaneous injections in adults, and at what angle?

A. 22–24 gauge; 1 inch; 60-degree angle

B. 25–27 gauge; 3/16 to 5/8 inch; 45-degree angle

C. 27–29 gauge; 1/4 to 5/8 inch; 30-degree angle

D. 20–22 gauge; 1 inch; 90-degree angle

SAMPLE

Answers

SAMPLE

1. A
2. A
3. A
4. A
5. A
6. A
7. C
8. D
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. What are the primary adult anatomical sites for intramuscular injections?

A. Anterior thighs, upper outer quadrant of buttocks, and deltoid.

B. Anterior thigh only.

C. Dorsogluteal muscle exclusively.

D. Vastus lateralis and gluteus medius only.

The main idea is that adults have several reliable intramuscular sites that provide good absorption and a lower risk of hitting nerves or vessels. The three primary sites used are the deltoid muscle of the upper arm, the gluteal region in the upper outer quadrant of the buttock, and the anterior (outer) portion of the thigh. The deltoid is good for small-volume injections, the upper outer buttock (gluteus maximus) handles larger volumes, and the anterior thigh region offers a large, accessible muscle mass for injections when needed. This combination explains why those areas are listed together as primary sites. The other options limit sites unnecessarily or point to a single site, which isn't how IM injections are typically approached in adults.

2. If Staphylococcus epidermidis is detected in only one of several blood culture bottles, what is the most likely interpretation?

A. Contamination from skin flora

B. Definitive bacteremia with this organism

C. Contamination across all bottles

D. Laboratory error with sample mislabeling

When evaluating blood cultures, the pattern of which bottles turn positive helps distinguish true infection from a sample mistake. Staphylococcus epidermidis is a common skin flora and a frequent contaminant when blood is drawn. If it appears in only one of several bottles, the finding is most consistent with contamination at the time of collection rather than a true bloodstream infection. If this organism were causing definitive bacteremia, you'd typically expect it to be isolated from multiple bottles or sets, and often alongside clinical signs of infection. Seeing the organism in a single bottle makes contamination far more likely. Options suggesting contamination across all bottles or a laboratory mislabeling would not align with the pattern of a solitary positive bottle, and while lab errors can occur, the more fitting interpretation given this pattern is contamination from skin flora.

3. In a patient with a left mastectomy and a functioning IV in the opposite arm, what is the correct approach to blood draw?

A. Turn off IV for 2 minutes, draw distally to IV, and discard the first 5 mL

B. Draw from the mastectomy side

C. Draw from the arm with the IV without modification

D. Use a leg vein

When you collect blood from a patient with an IV, the goal is to prevent IV fluids or medications from contaminating the blood sample. The best way to do this is to stop the IV for a short period, draw from a vein distal to the IV site, and discard the first amount drawn to flush out any IV fluid. In this scenario, with a left mastectomy (higher lymphedema risk on that side) and an IV in the opposite arm, the method described—turning the IV off for about two minutes, drawing from a vein distal to the IV, and discarding the first ~5 mL—achieves sample integrity if the draw would involve the arm with the IV. It reduces the chances that IV fluid adulterates the specimen. Avoiding the mastectomy side for phlebotomy remains important to minimize lymphedema risk, and drawing from a leg vein is generally a last resort.

SAMPLE

4. Which statement describes a contraindication to venipuncture due to venous fibrosis?

A. Venous fibrosis.

B. Varicose veins.

C. Small veins.

D. Normal skin.

Fibrotic veins are rigid and scarred, losing their elasticity and normal lumen structure. This makes needle entry unreliable, increases the risk of unsuccessful blood draw or vessel injury, and can lead to poor venous access or complications. Because the vein's fibrotic state compromises safety and effectiveness of venipuncture at that site, it should be avoided, and an alternative vein or access method should be used. Varicose veins are enlarged and tortuous and can be more difficult to cannulate, but they are not automatically a contraindication; sometimes they can still be used if no better option exists. Small veins pose technical challenges, yet without fibrosis they aren't a contraindication. Normal skin presents no barrier to venipuncture.

5. Capillary puncture is typically performed at which sites?

A. Fingerstick or heel.

B. Arm vein.

C. Neck vein.

D. Ankle.

Capillary puncture uses the tiny vessels just under the skin, so the best sites are where capillaries are closest to the surface. In adults, the fingertip is the standard site because it provides easy access to a dense capillary network and yields enough blood with a quick lancet puncture. In newborns and infants, the heel is used for the same reason, offering a safe, reliable capillary bed. The other sites—arm, neck, or ankle—are associated with larger veins and require venipuncture rather than capillary sampling, so they aren't used for capillary puncture.

6. Why is it important to keep the venipuncture site clean and dry after collection?

A. To reduce infection risk and promote hemostasis.

B. To speed up blood flow into the vein.

C. To prevent the patient from feeling faint.

D. To avoid contrast in future testing.

Keeping the venipuncture site clean and dry after collection reduces infection risk and supports hemostasis. Cleaning helps remove skin bacteria that could enter the puncture wound, while keeping the area dry allows the platelet plug and coagulation cascade to seal the puncture more effectively, minimizing ongoing bleeding. A dry, clean site also helps the dressing stay in place and reduces contamination if the bandage is disturbed. The other options don't fit because they deal with effects that aren't influenced by post-collection site cleanliness or dryness, such as blood flow into the vein, fainting, or future testing with contrast.

SAMPLE

7. Which specialized needle device is recommended when performing venipuncture on hand veins or fragile veins?

A. A 25-gauge winged butterfly device.

B. A 21-gauge winged butterfly device.

C. A 23-gauge winged butterfly device.

D. A 23-gauge straight needle.

When veins are small or fragile, you want a puncture that minimizes trauma and gives you solid control. A winged butterfly device provides stability and gentle handling for superficial veins, so you can puncture with a steadier, shallower entry. The 23-gauge size strikes a good balance: small enough to be gentle yet sturdy enough to maintain patency and allow enough blood flow for a successful draw. A larger gauge like 21 would be more traumatic for delicate hand veins, and a straight needle lacks the stabilizing wings that help guide the needle with precision. A 25-gauge butterfly might be too flimsy and slow to yield blood in a fragile vein. So the 23-gauge winged butterfly setup is the best choice for hand or fragile veins.

8. What constitutes a complete blood culture bottle collection kit?

A. One aerobic culture bottle and one anaerobic culture bottle.

B. Two aerobic culture bottles.

C. One anaerobic culture bottle only.

D. A bottle that contains both aerobic and anaerobic media.

A blood culture collection kit is meant to detect a broad range of bacteria, including those that grow with oxygen and those that don't. A bottle that contains both aerobic and anaerobic media provides the right environment for organisms with different oxygen needs in a single container. This design increases the likelihood of recovering pathogens from a sample, reduces the number of bottles handling, and minimizes the chance of contamination during transfer or transport. If you used only an aerobic bottle, anaerobic organisms might not grow well or at all. An anaerobic bottle alone would miss aerobes. Using two aerobic bottles would fail to detect anaerobes, limiting the test's sensitivity. So, a single bottle combining both media offers the most comprehensive, efficient approach for complete collection.

9. What is a recommended prevention for phlebitis?

A. Aseptic technique and site rotation

B. Frequent IV flush

C. Avoid cleaning around site

D. Increase IV rate

Phlebitis prevention focuses on keeping the IV site clean and minimizing irritation to the vein. Using aseptic technique during insertion and when handling the line reduces infection risk and chemical irritation from the IV fluids. Rotating IV sites limits repeated mechanical trauma to the same vein, lowering the chance of inflammation from ongoing cannulation. The other ideas don't address inflammation risk: frequent flushing keeps the line patent but doesn't prevent vein irritation; avoiding cleaning around the site raises infection risk; increasing the IV rate can increase irritation and vein damage.

SAMPLE

10. What are the typical needle gauge and length for subcutaneous injections in adults, and at what angle?

A. 22–24 gauge; 1 inch; 60-degree angle

B. 25–27 gauge; 3/16 to 5/8 inch; 45-degree angle

C. 27–29 gauge; 1/4 to 5/8 inch; 30-degree angle

D. 20–22 gauge; 1 inch; 90-degree angle

Subcutaneous injections are deposited into the fatty tissue just beneath the skin, so you want a short, fine needle and a shallow angle to stay in that layer. In adults, the typical combination is a fine gauge around 25–27 with a length of about 3/16 to 5/8 inch, inserted at roughly 45 degrees. This strikes a balance between delivering the medication into subcutaneous tissue and avoiding the muscle. Longer needles or a steeper angle (like 60 or 90 degrees) increase the chance of intramuscular delivery and more discomfort. Conversely, using a much thicker gauge or an angle suited for intramuscular or intradermal injections would not fit the standard subcutaneous technique.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://mpsvenipuncture.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE