NATIONAL PHYSICAL THERAPY EXAMINATION (NPTE) MODALITIES PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nptemodalities.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **For reducing patient pain during treatment of patellar tendonitis, which solution is most appropriate for use with iontophoresis?**

   A. Acetic acid

   B. Lidocaine

   C. Sodium chloride

   D. Zinc oxide

2. **Which ion is best to promote tissue healing when using iontophoresis for a lower extremity ulceration?**

   A. Acetate

   B. Magnesium

   C. Lidocaine

   D. Zinc

3. **What is the best choice for wheelchair prescription for a patient recovering from a right cerebrovascular accident resulting in severe left hemiplegia and a diabetic ulcer?**

   A. Hemiplegic chair with elevating leg rest on the left

   B. Powered wheelchair with joystick and elevating leg rests

   C. Lightweight active duty wheelchair with elevating leg rests

   D. One-arm drive chair with elevating leg rest on the left

4. **A patient with an ankle sprain exhibits worsening swelling. What is the best intervention at this time?**

   A. Intermittent compression followed by elevation

   B. Contrast baths followed by limb elevation

   C. Cold whirlpool followed by massage

   D. Cold/intermittent compression combination with the limb elevated

5. **A patient with spastic hemiplegia is referred for ambulation training. What biofeedback protocol should be used for knee extension?**

   A. High-detection sensitivity, with electrodes placed close together

   B. Low-detection sensitivity, with electrodes placed far apart

   C. High-detection sensitivity, with electrodes placed far apart

   D. Low-detection sensitivity, with electrodes placed close together

SAMPLE

6. What action would best minimize the impact of sudden force transition during neuromuscular electrical stimulation?

A. Decrease the treatment duration

B. Modify the waveform

C. Utilize a ramp

D. Increase the phase duration

7. Which ultrasound parameters produce purely nonthermal effects?

A. Low intensity, low duty cycle

B. Low intensity, high duty cycle

C. High intensity, low duty cycle

D. High intensity, high duty cycle

8. When using biphasic pulsed current for knee extension, what should the therapist adjust to make the current more tolerable while maintaining therapeutic effect?

A. Current polarity

B. Pulse duration

C. Current intensity

D. Pulse rate

9. If a burn from iontophoresis is an acidic reaction, what is the most likely pH level of the skin?

A. 2

B. 4

C. 6

D. 8

10. What action would best minimize the risk of a burn when using iontophoresis?

A. Increase the size of the cathode relative to the anode.

B. Decrease the space between the electrodes.

C. Increase the current density.

D. Decrease the moisture of the electrodes.

SAMPLE

Answers

SAMPLE

1. B
2. D
3. A
4. D
5. D
6. C
7. A
8. B
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. For reducing patient pain during treatment of patellar tendonitis, which solution is most appropriate for use with iontophoresis?

A. Acetic acid

B. Lidocaine

C. Sodium chloride

D. Zinc oxide

Lidocaine is an appropriate choice for iontophoresis in the treatment of patellar tendonitis because it is a local anesthetic that provides pain relief. When delivered through iontophoresis, lidocaine can penetrate the skin and create a temporary numbing effect in the localized area of treatment. This is particularly beneficial in managing pain associated with conditions like patellar tendonitis, as it helps improve patient comfort during physical therapy interventions. Other solutions mentioned in the options serve different therapeutic purposes. Acetic acid is typically used for conditions related to calcific deposits or tissue calcification; its role in pain reduction is not as direct as that of lidocaine. Sodium chloride is often used in acute management situations for hydrating tissues or in saline solutions but does not have analgesic properties. Zinc oxide is sometimes applied for its skin protective and healing properties but does not effectively alleviate pain in the way lidocaine does. Therefore, when specifically targeting pain reduction with iontophoresis, lidocaine is the most effective and appropriate choice.

2. Which ion is best to promote tissue healing when using iontophoresis for a lower extremity ulceration?

A. Acetate

B. Magnesium

C. Lidocaine

D. Zinc

Using iontophoresis to promote tissue healing, particularly for lower extremity ulcerations, has a specific application of certain ions due to their biological effects and interactions with tissues. Zinc is a crucial mineral for tissue regeneration and repair processes, making it particularly effective in promoting healing in wounds and ulcerations. It plays a vital role in protein synthesis, cell proliferation, and the inflammatory response, all of which are important in the healing process. When using iontophoresis, zinc can be delivered transdermally to the targeted area, enhancing local tissue repair mechanisms and accelerating the healing of the ulcer. Its use is supported by evidence that indicates a positive effect on wound healing due to its role in collagen synthesis and immune function. In contrast, while acetate, magnesium, and lidocaine have their own therapeutic benefits, they do not have the same direct and established role in enhancing tissue repair processes as zinc. Acetate is often used for its potential in breaking down calcium deposits, magnesium is involved in various body functions but not specifically for enhancing wound healing, and lidocaine serves primarily as a local anesthetic without direct benefits for tissue regeneration. Thus, zinc is recognized as the most effective ion for promoting tissue healing in this context.

SAMPLE

3. What is the best choice for wheelchair prescription for a patient recovering from a right cerebrovascular accident resulting in severe left hemiplegia and a diabetic ulcer?

A. Hemiplegic chair with elevating leg rest on the left

B. Powered wheelchair with joystick and elevating leg rests

C. Lightweight active duty wheelchair with elevating leg rests

D. One-arm drive chair with elevating leg rest on the left

A hemiplegic chair with elevating leg rest on the left is the most appropriate choice for a patient recovering from a right cerebrovascular accident with severe left hemiplegia and a diabetic ulcer. This type of wheelchair is specifically designed for individuals with hemiplegia, offering features that accommodate their needs. The chair's specific orientation allows the patient to efficiently utilize their functional right side for steering and propulsion, while providing necessary support and positioning to their left side, which is affected by paralysis. The elevating leg rest on the left helps manage and relieve pressure on the diabetic ulcer, promoting healing by reducing weight and friction on the affected area. Other options may not provide the specialized support or the balance needed for this patient's unique condition. For example, a powered wheelchair, while offering independence, might not be necessary given the patient's current recovery phase and could be more complex than needed. Lightweight active duty chairs or one-arm drive chairs may not adequately address hemiplegia-specific needs and could compromise safety and comfort during recovery. Selecting a chair that directly addresses the challenges of hemiplegia and pressure management while supporting recovery is critical for this patient.

4. A patient with an ankle sprain exhibits worsening swelling. What is the best intervention at this time?

A. Intermittent compression followed by elevation

B. Contrast baths followed by limb elevation

C. Cold whirlpool followed by massage

D. Cold/intermittent compression combination with the limb elevated

The most effective intervention for a patient with an ankle sprain that is exhibiting worsening swelling involves using a combination of cold therapy and intermittent compression, along with elevation of the limb. This approach is beneficial for several reasons. Cold therapy is instrumental in reducing inflammation and pain immediately following an injury. The cold works to constrict blood vessels, which minimizes blood flow to the affected area, thus attending to the acute phase of swelling. Intermittent compression provides mechanical support that helps prevent the accumulation of excess fluid in the tissues, further controlling swelling. By applying pressure to the swollen area and using a rhythmic release, intermittent compression can effectively assist in promoting circulation and lymphatic drainage. Elevating the limb is critical as it utilizes gravity to decrease hydrostatic pressure in the venous system, facilitating venous return and reducing swelling. Combining these strategies—cold for pain and inflammation reduction, intermittent compression for mechanical support and fluid control, and elevation for enhanced venous drainage—addresses the swelling holistically and is particularly suited for an acute injury like an ankle sprain. This multifaceted approach is especially advantageous when a patient presents with worsening symptoms, indicating a need for comprehensive intervention to manage and mitigate swelling effectively.

SAMPLE

5. A patient with spastic hemiplegia is referred for ambulation training. What biofeedback protocol should be used for knee extension?

A. High-detection sensitivity, with electrodes placed close together

B. Low-detection sensitivity, with electrodes placed far apart

C. High-detection sensitivity, with electrodes placed far apart

D. Low-detection sensitivity, with electrodes placed close together

In the context of biofeedback protocols for knee extension in a patient with spastic hemiplegia, using low-detection sensitivity with electrodes placed close together is the most appropriate choice. This approach allows for enhanced control by providing meaningful feedback to the patient. When utilizing low-detection sensitivity, the biofeedback system is designed to register and respond to larger muscle contractions, which can be particularly beneficial in a patient with spasticity. The close placement of electrodes facilitates the capture of targeted muscle activity and increases the chances of detecting relevant muscle contractions effectively. This setup can improve the precision of the feedback, subsequently aiding in the patient's ability to gain better control over knee extension during ambulation training. Additionally, the use of low-detection sensitivity can help manage the irregular and sometimes overly strong contractions associated with spasticity, as it reduces the likelihood of eliciting overwhelming feedback that might occur with high-detection settings. This balance is significant in promoting the engagement of the appropriate muscles for knee extension while avoiding additional complications from spastic reflexes.

6. What action would best minimize the impact of sudden force transition during neuromuscular electrical stimulation?

A. Decrease the treatment duration

B. Modify the waveform

C. Utilize a ramp

D. Increase the phase duration

Utilizing a ramp during neuromuscular electrical stimulation (NMES) is essential for minimizing the impact of sudden force transitions. A ramp function gradually increases the intensity of the stimulation over a specified period, allowing the muscle to contract more naturally and avoiding the jarring effect that a sudden onset of stimulation would cause. This gradual increase aids in the prevention of discomfort and allows the muscle fibers to engage more cooperatively, which can enhance the efficacy of the treatment and ensure a smoother transition into and out of contraction. The other options do not effectively address the sudden change in force. Decreasing the treatment duration may limit the overall effectiveness of the stimulation, as there is less time for the muscle to respond appropriately. Modifying the waveform can alter the characteristics of the electrical current but does not inherently reduce the abruptness of muscle activation. Increasing the phase duration might enhance muscle contraction but would not prevent sudden force transitions, as it may still elicit a rapid increase in force that can be uncomfortable or damaging. Thus, incorporating a ramp effectively smooths the transition and is the best choice for minimizing the effects of abrupt changes in stimulation.

SAMPLE

7. Which ultrasound parameters produce purely nonthermal effects?

A. Low intensity, low duty cycle

B. Low intensity, high duty cycle

C. High intensity, low duty cycle

D. High intensity, high duty cycle

The parameters that produce purely nonthermal effects in ultrasound therapy are characterized by low intensity combined with a low duty cycle. When using ultrasound in a therapeutic context, nonthermal effects are primarily associated with mechanical and biochemical changes at the cellular level rather than an increase in temperature. Low intensity minimizes the heating effects caused by ultrasound waves, while a low duty cycle indicates that the ultrasound is on for a smaller percentage of the treatment time compared to being off. This combination allows for the mechanical effects of ultrasound, such as cavitation and microstreaming, to influence tissue healing without generating significant heat. In contrast, a high intensity setting typically increases the likelihood of thermal effects occurring, as it delivers more energy to the tissues, raising their temperature. Additionally, a high duty cycle keeps the ultrasound on for a larger portion of the treatment time, further promoting heat production. Therefore, options that involve either high intensity or high duty cycle are less likely to result in purely nonthermal outcomes. Thus, low intensity and a low duty cycle are specifically designed to achieve the desired nonthermal effects in therapeutic ultrasound.

8. When using biphasic pulsed current for knee extension, what should the therapist adjust to make the current more tolerable while maintaining therapeutic effect?

A. Current polarity

B. Pulse duration

C. Current intensity

D. Pulse rate

Adjusting the pulse duration when using biphasic pulsed current can make the current more tolerable while still maintaining the therapeutic effect. Pulse duration refers to the length of time each pulse lasts. Shorter pulse durations can often be perceived as less intense by the patient, reducing discomfort while still targeting the desired muscle groups for stimulation during rehabilitation, such as knee extension. Maintaining an effective pulse duration is crucial because it influences both the strength of muscle contraction and patient comfort. Too long of a pulse duration can lead to increased discomfort without necessarily enhancing the therapeutic benefits. By fine-tuning this variable, the therapist can ensure that the treatment remains effective in achieving the desired muscle response while preventing excessive discomfort for the patient, thereby enhancing overall adherence to the treatment protocol. Adjusting the current intensity might risk either reducing the effectiveness of the treatment or causing discomfort if set too high. Current polarity typically does not affect tolerability in the same way and is more applicable in specific clinical scenarios. Pulse rate influences the frequency of muscle contractions but does not directly correlate to comfort when maintaining therapeutic goals.

SAMPLE

9. If a burn from iontophoresis is an acidic reaction, what is the most likely pH level of the skin?

A. 2

B. 4

C. 6

D. 8

In the context of iontophoresis, when a burn occurs as a result of an acidic reaction, it indicates an increase in hydrogen ions (H+) at the electrode site. This process generally leads to a lower pH, which means the skin becomes more acidic. The pH scale ranges from 0 to 14, where values less than 7 indicate an acidic environment. A pH of 2, being significantly lower than 7, represents a strongly acidic condition. This aligns with the understanding that injuries or reactions characterized by acidity can lead to very low pH levels. In this case, the most likely pH level of the skin that experiences this acidic reaction from iontophoresis is indeed 2, confirming that the environment on the skin has become markedly acidic. Other pH options like 4 and 6, while still on the acidic side, do not reflect the severity of the acidic reaction typically associated with a burn from this modality. A pH of 8, on the other hand, is alkaline and contrasts with the acidic nature suggested by the reaction. Therefore, understanding the implications of an acidic reaction is crucial in assessing the pH level resulting from a burn due to iontophoresis.

10. What action would best minimize the risk of a burn when using iontophoresis?

A. Increase the size of the cathode relative to the anode.

B. Decrease the space between the electrodes.

C. Increase the current density.

D. Decrease the moisture of the electrodes.

To minimize the risk of a burn when using iontophoresis, increasing the size of the cathode relative to the anode is a sound choice. This is because a larger cathode disperses the current over a greater surface area, which reduces the current density at any given point on the skin. A lower current density helps to minimize thermal effects, such as burns, as less electrical energy is concentrated in any one area of the skin. Using a larger cathode can help ensure that the skin does not become overly heated during treatment because it reduces the likelihood of generating excessive heat at the site of application. Additionally, the cathode's configuration can influence the distribution of ions and the overall effectiveness of the treatment. Thus, selecting a cathode with a larger surface area compared to the anode not only enhances safety but can also improve therapeutic outcomes in iontophoresis. The other actions listed, such as reducing the distance between electrodes or increasing current density, would likely increase the risk of burns instead of minimizing it. Additionally, decreasing the moisture of the electrodes could reduce conductivity, potentially leading to inefficient treatment rather than improved safety.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nptemodalities.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE