NATIONAL PHYSICAL THERAPY EXAMINATION (NPTE) PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://npte.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What does 4+ on the pitting edema scale indicate?

A. No pitting

B. A slight indentation that returns to normal after 30 seconds

C. A deep indentation that remains for more than 30 seconds

D. A negligible indentation

2. Which clinical finding is indicative of an unstable fracture site in the context of a femoral fracture?

A. Intact range of motion

B. Severe swelling

C. Increased pain with weight-bearing

D. A visible deformity

3. What is a key benefit of fluidotherapy for patients recovering from fractures?

A. Provides heat and moisture

B. Promotes deep tissue massage

C. Encourages muscle strengthening

D. Aids in immediate pain relief

4. In physical therapy, which approach is appropriate for managing a patient with hip osteoarthritis?

A. Immediate surgical intervention

B. Manual therapy and exercise prescription

C. Complete rest without any movement

D. Application of a hot pack on the hip only

5. If a patient demonstrates an increase in heart rate of 25 beats/min during exercise, what should the physical therapist assistant consider?

A. Normal reaction to exercise

B. Possible cardiovascular distress

C. Dehydration

D. Fatigue

SAMPLE

6. In managing a patient, which condition could cause swelling and fluid retention in a home care setting?

A. Asthma exacerbation

B. Congestive heart failure

C. Chronic Kidney Disease

D. Pneumonia

7. For a patient taking thiazide medication, which combination of signs may indicate an adverse effect?

A. Agitation, dizziness, dyspnea, and hyperventilation

B. Abdominal cramps, agitation, dyspnea, and numbness

C. Clammy skin, dizziness, hyperventilation, and hypotension

D. Abdominal cramps, clammy skin, hypotension, and numbness

8. Which location of a cerebrovascular accident (CVA) is most likely for a patient with sensory loss in the right face and arm?

A. Left MCA inferior division

B. Left MCA superior division

C. Right MCA superior division

D. Right MCA inferior division

9. In the context of postural control, which position is typically recommended for individuals with balance deficits?

A. Sitting

B. Standing

C. Kneeling

D. Supine

10. What is the best predictor of altered sensation in a patient recovering from an upper extremity burn?

A. The percentage of the body surface affected

B. The extent of any hypertrophic scarring

C. The presence of a skin graft

D. The depth of the burn injury

SAMPLE

Answers

SAMPLE

1. C
2. C
3. A
4. B
5. B
6. B
7. C
8. A
9. B
10. D

SAMPLE

Explanations

SAMPLE

1. What does 4+ on the pitting edema scale indicate?

A. No pitting

B. A slight indentation that returns to normal after 30 seconds

C. A deep indentation that remains for more than 30 seconds

D. A negligible indentation

A rating of 4+ on the pitting edema scale indicates a deep indentation that remains for more than 30 seconds. This level of edema is the most severe category on the scale, reflecting significant fluid retention in the interstitial spaces. The sustained indentation demonstrates that there is a substantial accumulation of fluid, which the tissues are unable to disperse quickly. Understanding the pitting edema scale is crucial for clinicians, as it helps assess the severity of edema. A rating of 4+ suggests significant concern, often requiring further evaluation of the underlying causes, such as heart failure, kidney issues, or other systemic conditions. The persistent nature of the indentation indicates a considerable impact on the body's ability to manage and regulate fluid levels effectively.

2. Which clinical finding is indicative of an unstable fracture site in the context of a femoral fracture?

A. Intact range of motion

B. Severe swelling

C. Increased pain with weight-bearing

D. A visible deformity

In the context of a femoral fracture, increased pain with weight-bearing is a strong indicator of an unstable fracture site. When a fracture site is unstable, applying weight or elevating the mechanical load to the affected area often results in heightened pain. This occurs because unstable fractures typically involve significant disruption of bone integrity and the surrounding soft tissues, leading to instability that is exacerbated by movement or bearing weight. The increased stress on the injury from weight-bearing can further irritate the fracture site and the surrounding structures, causing the patient's pain to worsen. Other clinical findings, such as severe swelling, may be present in various injury types, including stable fractures or soft tissue injuries, making them less specific. Intact range of motion could sometimes be observed even with a fractured bone, particularly if there is no associated soft tissue damage or if the fracture is stable. A visible deformity can also signal a fracture, but not all fractures, especially stable ones, will present with noticeable deformities. Therefore, the presence of increased pain specifically correlates to the instability of the fracture site, making it the most indicative clinical finding in this scenario.

SAMPLE

3. What is a key benefit of fluidotherapy for patients recovering from fractures?

A. Provides heat and moisture

B. Promotes deep tissue massage

C. Encourages muscle strengthening

D. Aids in immediate pain relief

Fluidotherapy is a therapeutic modality that utilizes circulating warm, dry particles to provide treatment, particularly beneficial in rehabilitation settings. One of its key benefits is that it provides heat and moisture. The circulating air in the fluidotherapy chamber warms the particles, leading to an increase in the local temperature of the tissues, which can accelerate the healing process following fractures. The warmth facilitates increased blood flow to the affected area, promoting tissue relaxation and reducing stiffness, which is crucial for recovery after a fracture. Additionally, the moisture component, which can also be adjusted to suit patient needs, helps to hydrate the skin and underlying tissues, making it particularly useful for individuals who may have compromised skin integrity due to immobilization or other factors. In the context of recovery from fractures, the combination of heat and moisture can alleviate discomfort and enhance mobility, making it an effective therapeutic option in rehabilitative care.

4. In physical therapy, which approach is appropriate for managing a patient with hip osteoarthritis?

A. Immediate surgical intervention

B. Manual therapy and exercise prescription

C. Complete rest without any movement

D. Application of a hot pack on the hip only

The appropriate approach for managing a patient with hip osteoarthritis is centered around manual therapy and exercise prescription. This method focuses on improving joint mobility, reducing pain, and enhancing function. Manual therapy techniques can include soft tissue mobilization and joint mobilization, which aim to alleviate stiffness and improve the range of motion in the affected hip joint. Exercise prescription is crucial as it helps strengthen the surrounding musculature, improve stability, and maintain joint function. These exercises can be tailored to the individual's level of pain and function, making them a vital component of an effective treatment plan for osteoarthritis. Both of these interventions empower patients to take an active role in their rehabilitation, promote long-term joint health, and can lead to better management of symptoms compared to the other options listed.

SAMPLE

5. If a patient demonstrates an increase in heart rate of 25 beats/min during exercise, what should the physical therapist assistant consider?

A. Normal reaction to exercise

B. Possible cardiovascular distress

C. Dehydration

D. Fatigue

An increase in heart rate during exercise is a typical response as the body requires more oxygen-rich blood to be delivered to the working muscles. However, a significant increase of 25 beats per minute can indicate that the patient's cardiovascular system may be responding improperly to the demands of exercise. The normal physiological response usually involves a heart rate increase that aligns with the intensity of the activity performed. An exaggerated response, especially if it occurs alongside other symptoms, might suggest possible cardiovascular distress, which needs to be monitored closely. This is particularly crucial for patients with existing cardiovascular conditions or risk factors, as they might not tolerate exertion as well as others. It's vital for healthcare professionals to assess whether the increased heart rate is appropriate based on the patient's baseline and the intensity of the exercise being performed. If it appears excessive or is accompanied by other symptoms such as chest pain, dizziness, or shortness of breath, it may warrant further investigation to ensure the patient's safety.

6. In managing a patient, which condition could cause swelling and fluid retention in a home care setting?

A. Asthma exacerbation

B. Congestive heart failure

C. Chronic Kidney Disease

D. Pneumonia

Congestive heart failure can lead to swelling and fluid retention in a home care setting due to the heart's impaired ability to pump blood effectively. When the heart is unable to maintain adequate circulation, fluid can back up in the veins, leading to increased pressure that forces fluid out of the blood vessels and into surrounding tissues. This often results in edema, particularly in the lower extremities and sometimes in the abdomen. Fluid retention can also occur because the kidneys attempt to compensate for decreased cardiac output by retaining sodium and water, further contributing to the swelling. In home care environments, patients with congestive heart failure might require monitoring for weight changes, edema, and other symptoms related to fluid overload, making it a significant condition to be aware of for managing overall health and quality of life. In contrast, while asthma exacerbation can lead to breathing difficulties, it is not primarily associated with fluid retention. Chronic Kidney Disease also causes fluid retention, but it is primarily a renal-related issue, and the question implies a more direct relationship with heart function. Pneumonia can cause respiratory symptoms and systemic effects but does not typically result in localized swelling and fluid retention in the manner seen with congestive heart failure.

SAMPLE

7. For a patient taking thiazide medication, which combination of signs may indicate an adverse effect?

A. Agitation, dizziness, dyspnea, and hyperventilation

B. Abdominal cramps, agitation, dyspnea, and numbness

C. Clammy skin, dizziness, hyperventilation, and hypotension

D. Abdominal cramps, clammy skin, hypotension, and numbness

Thiazide diuretics are commonly prescribed to manage conditions such as hypertension and heart failure, but they can lead to several adverse effects. The combination of clammy skin, dizziness, hyperventilation, and hypotension suggests a state of dehydration or electrolyte imbalance, which are known risks associated with thiazide use. Clammy skin can indicate poor peripheral perfusion, often related to hypotension or low blood volume. Dizziness signals that the patient's blood pressure may be too low, leading to inadequate cerebral perfusion. Hyperventilation can occur as a physiological response to metabolic changes or anxiety but may also manifest in the context of electrolyte disturbances. Hypotension, or low blood pressure, is a direct consequence of diuretic use, especially if the patient has experienced excessive fluid loss. In this case, the combination of these signs suggests a serious adverse reaction to the thiazide medication, particularly due to possible volume depletion and associated electrolyte imbalances, which are critical to monitor in patients taking these drugs. This reinforces the necessity for clinicians to carefully observe patients on thiazide diuretics for these potential complications.

8. Which location of a cerebrovascular accident (CVA) is most likely for a patient with sensory loss in the right face and arm?

A. Left MCA inferior division

B. Left MCA superior division

C. Right MCA superior division

D. Right MCA inferior division

The left middle cerebral artery (MCA) inferior division is responsible for supplying blood to areas of the brain that control sensory and motor functions for the right side of the body, specifically the right face and arm, due to the contralateral organization of the nervous system. If a cerebrovascular accident (CVA) occurs in this region, it can lead to sensory loss on the opposite side of the body, which in this case would present as sensory loss in the right face and arm. In the context of the other options, each potential division of the MCA can impact different functional areas of the brain. The left MCA superior division primarily supplies the regions responsible for motor functions and higher cognitive functions, leading to motor deficits or language issues rather than isolated sensory loss. The right MCA divisions similarly control the motor and sensory functions for the left side of the body. Thus, they would not typically cause sensory loss in the right face and arm. Understanding the vascular supply and functional areas of the brain is crucial in identifying the cause of specific deficits following a CVA.

SAMPLE

9. In the context of postural control, which position is typically recommended for individuals with balance deficits?

A. Sitting

B. Standing

C. Kneeling

D. Supine

The recommended position for individuals with balance deficits is standing because it promotes active engagement of the postural control mechanisms that are crucial for maintaining stability. When standing, individuals can utilize their sensory inputs—including visual, vestibular, and somatosensory systems—to maintain balance. Standing allows for an automatic adjustment of the body's center of mass over the base of support, which is critical in improving postural stability. Moreover, standing can facilitate functional mobility and readiness for movement, encouraging the individual to practice and integrate balance strategies they may need in everyday activities. Sitting, while stable and providing support, can limit the activation of the muscles involved in standing balance control and does not challenge postural stability effectively. Kneeling can also be beneficial for some individuals but may not provide the same level of engagement and practice with balance strategies as standing allows. The supine position, being the most stable and supportive, typically does not engage balance systems adequately, as it does not simulate the challenges faced during upright activities. Thus, standing is the position that fosters the necessary skills and strength required for improving balance in individuals with deficits.

10. What is the best predictor of altered sensation in a patient recovering from an upper extremity burn?

A. The percentage of the body surface affected

B. The extent of any hypertrophic scarring

C. The presence of a skin graft

D. The depth of the burn injury

The depth of the burn injury is the best predictor of altered sensation in a patient recovering from an upper extremity burn because it directly impacts the integrity of nerve endings and sensory pathways. Burns are classified into various depths—such as superficial, partial-thickness, and full-thickness—and each type affects the skin layers and associated structures differently. A superficial burn may only involve the epidermis, preserving most sensory function, while deeper burns can damage nerve endings and disrupt normal sensation, leading to increased risks of numbness, tingling, or hypersensitivity in the affected area during the healing process. The depth of the burn is, therefore, crucial in understanding potential sensory deficits and guiding rehabilitation efforts to address these challenges. While factors like the percentage of body surface affected, extent of hypertrophic scarring, and presence of skin grafts can influence recovery and sensation to some degree, they do not have the same direct relationship with sensory nerve involvement as the depth of the burn injury does. Hence, depth serves as the primary factor in predicting altered sensory perception during the recovery phase.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://npte.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE