NATIONAL SAFETY COUNCIL (NSC) CPR CERTIFICATION PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nsccpr.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What specific technique is used for infants during CPR?

A. Use the palm of the hand for compressions

B. Use a single finger for compressions

C. Use 2 fingers for compressions at a depth of about 1.5 inches

D. Use an adult compression method

2. A sign of an internal abdominal injury is?

A. Sweating

B. Seizures

C. A swollen or rigid abdomen

D. All of the above

3. Why is CPR considered a valuable skill?

A. It can only be used in medical settings

B. It may save someone's life during cardiac emergencies

C. It requires extensive training to master

D. It is rarely used in real-life situations

4. When should an Automated External Defibrillator (AED) be used?

A. When the victim is breathing normally

B. As soon as it is available after calling 911

C. Only after 5 minutes of CPR

D. After all breathing techniques have been attempted

5. What is the primary goal of first aid?

A. To make a diagnosis of the injury

B. To stabilize the victim's condition until professional help arrives

C. To alleviate pain as much as possible

D. To provide advanced medical treatment

6. DOTS stands for?

A. Deformities, open wounds, tenderness, swelling

B. Dislocations, open injuries, trauma, soreness

C. Deformities, obstructions, tenderness, severity

D. Discoloration, open wounds, trauma, swelling

SAMPLE

7. True or False: In any emergency, the best thing to do is to move the victim to your car and rush to an emergency department.

A. True

B. False

8. Which sign indicates you should continue CPR?

A. The victim is not moving

B. You feel tired

C. You hear bystanders discussing other matters

D. A medical professional arrives

9. What should be the first step if you suspect someone is choking?

A. Perform abdominal thrusts immediately

B. Encourage them to cough

C. Give them water

D. Call 911

10. When performing CPR on an infant or child, compress to a depth of at least what fraction of the chest depth?

A. 1/4

B. 1/3

C. 1/2

D. 2/3

SAMPLE

Answers

SAMPLE

1. C
2. C
3. B
4. B
5. B
6. A
7. B
8. A
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What specific technique is used for infants during CPR?

A. Use the palm of the hand for compressions

B. Use a single finger for compressions

C. Use 2 fingers for compressions at a depth of about 1.5 inches

D. Use an adult compression method

When performing CPR on infants, the technique involves using two fingers for chest compressions. This method is designed to provide adequate pressure while being gentle enough to prevent injury to the fragile ribcage of an infant. The compressions should be delivered at a depth of about 1.5 inches, which is appropriate for the infant's anatomy and ensures that the heart is effectively compressed to circulate blood. Using two fingers helps to target the correct area on the infant's chest, which is just below the nipple line in the center of the chest. This technique not only promotes effective blood circulation but also minimizes the risk of trauma to the infant's soft bones. The specific depth of 1.5 inches is critical for achieving the necessary compression while maintaining safety. The other options present techniques that are either inappropriate for infants or do not follow established guidelines for infant CPR. The method chosen is essential for ensuring the safety and effectiveness of CPR in young children, where anatomical considerations are significantly different from those of adults.

2. A sign of an internal abdominal injury is?

A. Sweating

B. Seizures

C. A swollen or rigid abdomen

D. All of the above

A swollen or rigid abdomen is a key sign of an internal abdominal injury because it indicates that there may be bleeding or fluid accumulation within the abdominal cavity. When injuries occur, such as a rupture of organs, the body often responds by tightening the abdominal muscles in an attempt to protect the affected area. This rigidity can be felt during a physical examination. Furthermore, a swollen abdomen may suggest the presence of internal bleeding, which is a critical situation requiring immediate medical attention. The body's inflammatory response to injury can also lead to swelling, causing the abdomen to appear distended. Recognizing this symptom is crucial for effective assessment and intervention in emergency situations. While sweating and seizures might be symptoms present in various medical emergencies, they are not as directly indicative of internal abdominal injuries as rigidity and swelling of the abdomen.

3. Why is CPR considered a valuable skill?

A. It can only be used in medical settings

B. It may save someone's life during cardiac emergencies

C. It requires extensive training to master

D. It is rarely used in real-life situations

CPR, or cardiopulmonary resuscitation, is considered a valuable skill primarily because it has the potential to save lives in critical situations, particularly during cardiac emergencies. When a person experiences cardiac arrest, their heart stops beating effectively, and blood flow to the brain and other vital organs ceases. Performing CPR can maintain blood circulation and oxygenation until professional medical help arrives, significantly increasing the chances of survival and recovery for the victim. This life-saving technique is crucial in various scenarios, such as choking incidents, drowning, or heart attacks, where immediate action can make all the difference. Knowing how to perform CPR empowers individuals to act confidently and effectively in emergencies, ultimately contributing to better outcomes for those affected by cardiac events.

SAMPLE

4. When should an Automated External Defibrillator (AED) be used?

A. When the victim is breathing normally

B. As soon as it is available after calling 911

C. Only after 5 minutes of CPR

D. After all breathing techniques have been attempted

The appropriate time to use an Automated External Defibrillator (AED) is as soon as it is available after calling 911. This is essential because early defibrillation is a critical component of the chain of survival for someone experiencing a cardiac arrest. The AED is designed to quickly analyze the heart's rhythm and determine if a shock is necessary to restore a normal heart rhythm. Using the AED promptly can significantly increase the chance of survival for the victim, as every minute without defibrillation decreases the chances of survival. It is important to note that if the individual is showing signs of life such as normal breathing, then the use of an AED is not necessary at that moment. The AED should not be delayed by waiting for prolonged CPR to be completed or to try additional breathing techniques, as these actions can divert critical attention from restoring the heart rhythm. Instead, rapid access to the AED, combined with high-quality CPR, is the best course of action in such emergency situations.

5. What is the primary goal of first aid?

A. To make a diagnosis of the injury

B. To stabilize the victim's condition until professional help arrives

C. To alleviate pain as much as possible

D. To provide advanced medical treatment

The primary goal of first aid is to stabilize the victim's condition until professional help arrives. This encompasses providing immediate care that can prevent the situation from worsening and can be crucial in preserving life and promoting recovery. First aid involves assessing the scene and the individual, managing life-threatening conditions, and providing basic emergency care, which may include CPR, controlling bleeding, or assisting with choking. First aid acts as a bridge between the moment an injury occurs and the arrival of further medical assistance, emphasizing the importance of acting quickly and effectively. By stabilizing the individual's condition, you can help ensure that they remain as safe as possible until more comprehensive care is available. Making a diagnosis of the injury is not a primary goal of first aid, as this typically requires professional training and tools. Alleviating pain is also important, but it is secondary to the critical aim of maintaining life and preventing further harm. Advanced medical treatment is beyond the scope of first aid and should only be handled by trained medical personnel. Thus, focusing on stabilization reflects the essential principles of first aid effectively.

SAMPLE

6. DOTS stands for?

A. Deformities, open wounds, tenderness, swelling

B. Dislocations, open injuries, trauma, soreness

C. Deformities, obstructions, tenderness, severity

D. Discoloration, open wounds, trauma, swelling

The correct answer is based on the acronym DOTS, which is a systematic approach used in first aid assessments to help responders identify potential injuries in a patient. Specifically, DOTS stands for Deformities, Open wounds, Tenderness, and Swelling. - Deformities refer to any unnatural shapes or alignment in bones or body parts, suggesting fractures or dislocations. - Open wounds indicate breaches in the skin that may require attention to prevent infection and control bleeding. - Tenderness is assessed by palpating the injured area; tenderness can indicate underlying injuries that may not be visible. - Swelling often occurs following injury and can signal inflammation or trauma, which is important for treatment considerations. By using this acronym, first responders can quickly and efficiently gather critical information about a patient's condition. Understanding each component of DOTS allows responders to prioritize care effectively and communicate clearly with medical personnel once they arrive. This comprehensive method of assessment is essential for ensuring that all aspects of a patient's injuries are evaluated timely.

7. True or False: In any emergency, the best thing to do is to move the victim to your car and rush to an emergency department.

A. True

B. False

In emergency situations, the priority is to ensure the safety and well-being of the victim while also minimizing further harm. Moving a victim, especially one who may have sustained serious injuries, can lead to additional complications such as worsening a spinal injury, causing internal bleeding to worsen, or jeopardizing their airway. Proper emergency protocol generally dictates that the victim should remain still and should only be moved if there is an immediate danger to their safety, such as a fire or an unstable structure. It's crucial to call for professional medical help and provide care on-site until trained responders arrive. Staying with the victim, monitoring their condition, and providing appropriate first aid is typically more beneficial than attempting to transport them personally, which may create additional risks. Thus, the response should affirm the understanding of emergency care by emphasizing the importance of keeping victims stable and awaiting professional medical assistance, rather than moving them unnecessarily.

8. Which sign indicates you should continue CPR?

A. The victim is not moving

B. You feel tired

C. You hear bystanders discussing other matters

D. A medical professional arrives

Continuing CPR is essential in maintaining circulation and providing oxygen to a person who is unresponsive and not breathing. The sign that indicates you should continue performing CPR is when the victim is not moving. This lack of movement is a critical indicator that the person's heart is likely not beating effectively, and they are unresponsive, meaning immediate and continuous CPR is required until professional help arrives or the victim shows signs of life, such as movement or breathing. In contrast, feeling tired or fatigued during CPR does not mean you should stop. While it's important to switch rescuers if possible, remaining exhausted shouldn't be a sole reason to discontinue care. Hearing bystanders discussing other matters is irrelevant to the need for continued CPR; their conversation does not impact the medical condition of the victim. The arrival of a medical professional indeed indicates that you may cease CPR, but this is contingent upon their assessment or interventions being begun. Therefore, the most direct and clear sign that continues the urgency for CPR is the lack of movement from the victim.

SAMPLE

9. What should be the first step if you suspect someone is choking?

A. Perform abdominal thrusts immediately

B. Encourage them to cough

C. Give them water

D. Call 911

When someone is suspected of choking, encouraging them to cough is considered the first appropriate step because coughing is the body's natural reflex to clear an obstruction from the airway. Coughing can help to expel the object causing the blockage if it is lodged in the throat but not completely obstructing airflow. This approach allows the individual to attempt to clear their own airway with a method that is non-invasive and generally safe. If their coughing is productive and they are still able to speak, breathe, or make sounds, it indicates that the airway is not completely blocked, and you should allow them to continue coughing while monitoring their condition closely. In instances where the individual is unable to cough effectively, has difficulty breathing, or loses consciousness, then more forceful interventions, such as abdominal thrusts, become necessary. However, starting with encouraging a cough is a natural and helpful initial response before escalating to more aggressive measures.

10. When performing CPR on an infant or child, compress to a depth of at least what fraction of the chest depth?

A. 1/4

B. 1/3

C. 1/2

D. 2/3

When performing CPR on an infant or child, the recommended compression depth is at least one-third of the chest depth. For infants, this typically translates to approximately 1.5 inches, while for children, it can range from 2 inches to a little more, depending on the size of the child. Using this depth is crucial because it ensures that sufficient pressure is applied to circulate blood effectively, thus maintaining blood flow to vital organs during cardiac arrest. Compressing to a depth less than one-third may not generate adequate blood flow, reducing the effectiveness of the compressions. This understanding reinforces the importance of proper technique in CPR, especially with smaller patients, where the risk of causing injury from overly aggressive compressions must be balanced with the need for effective resuscitation.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nsccpr.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE