NBEO OCULAR MOTILITY PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nbeoocularmotility.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which movement is a conjugate movement?

A. Vergences

B. Incyclotorsion

C. Dextrocycloversion

D. Versions

2. Which muscle has excyclotorsion as its secondary action?

A. Lateral Rectus

B. Inferior Rectus

C. Superior Oblique

D. Superior Rectus

3. NOT true regarding the vestibulo-ocular reflex (VOR)?

A. It is stimulated by the movement of endolymph within the semicircular canals

B. It stabilizes images during head movements

C. It requires a visual stimulus

D. VOR typically only compensates for up to 30 second periods at a time

4. Prism adaptation is used in ocular motility management for which scenario?

A. It is used for large strabismus requiring surgical management.

B. It permanently cures diplopia.

C. Prism adaptation refers to gradual improvement in alignment with wearing prisms over time; used for convergence insufficiency or other small misalignments with residual diplopia.

D. It has no clinical use.

5. Which reflex is primarily responsible for stabilizing vision during head rotation by producing compensatory eye movements?

A. Optokinetic reflex (OKR)

B. Optokinetic nystagmus (OKN)

C. Vestibulo-ocular reflex (VOR)

D. Pursuit

SAMPLE

6. Which test mainly evaluates pursuits?

A. King-Devick

B. DEM

C. Pierce Saccades Tests

D. Groffman Tracings

7. Oculocephalic testing is also known as which maneuver?

A. Head thrust test

B. Skew deviation test

C. Doll's head maneuver

D. Cover test

8. Duane's Retraction syndrome Type 2 is defined by which limitation?

A. Limited abduction

B. Limited adduction

C. Limited abduction and adduction

D. Normal

9. End-point nystagmus is physiological.

A. Pathological

B. Physiological

C. Both

D. Neither

10. Describe the near point of convergence (NPC) test and typical normal values.

A. Move a small target toward the nose along the midline; typical NPC 6–8 cm in adults

B. Move a large target toward the nose; typical NPC 20–30 cm

C. Move target away from the nose; typical NPC 12–15 cm

D. NPC is not used in adults

SAMPLE

Answers

SAMPLE

1. D
2. B
3. C
4. C
5. C
6. D
7. C
8. B
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. Which movement is a conjugate movement?

A. Vergences

B. Incyclotorsion

C. Dextrocycloversion

D. Versions

Conjugate movements are eye movements where both eyes move in the same direction by the same amount, keeping the gaze aligned. This is exactly what versions describe—looking up, down, left, or right with both eyes moving together. Vergences, on the other hand, are disjunctive: the eyes move in opposite directions to adjust for depth (convergence for near, divergence for far). Incyclotorsion is rotation around the eye's visual axis of one eye, a torsional movement rather than a coordinated binocular shift. Dextrocycloversion implies a rightward gaze with a torsional component, but the straightforward binoc ular movement that fits the common definition of conjugate gaze shifts is versions. So the movement that represents a conjugate shift of both eyes in the same direction is versions.

2. Which muscle has excyclotorsion as its secondary action?

A. Lateral Rectus

B. Inferior Rectus

C. Superior Oblique

D. Superior Rectus

Excyclotorsion is the outward rotation of the top of the eye around the visual axis. Each extraocular muscle has a primary action (the main movement) and secondary actions that arise from its pull around the eye's axes. The inferior rectus mainly depresses the eye, but because of where it attaches on the globe, its pull also causes the eye to rotate outward around the optic axis. That outward rotation is excyclotorsion, making it the secondary torsional action of this muscle. Other muscles contribute torsion in different directions (for example, the superior rectus mainly intorts, and the superior oblique is a primary intorter and depressor), so the inferior rectus is the one whose secondary action includes excyclotorsion.

3. NOT true regarding the vestibulo-ocular reflex (VOR)?

A. It is stimulated by the movement of endolymph within the semicircular canals

B. It stabilizes images during head movements

C. It requires a visual stimulus

D. VOR typically only compensates for up to 30 second periods at a time

The vestibulo-ocular reflex is a vestibular-driven mechanism that keeps your gaze steady during head movement by producing eye movements in the opposite direction of where you're moving your head. When your head turns, the endolymph in the semicircular canals moves, bending the hair cells and sending signals to eye muscles to generate a counter-rotation of the eyes. This reflex works even without any visual input, so it can stabilize images in the dark. Vision can enhance the overall stability, but the VOR itself does not require a visual stimulus to function. That's why the statement saying it requires a visual stimulus isn't true. The VOR does stabilize images during head movements, which is accurate. And there isn't a strict 30-second limit to its compensation—the reflex continues as long as head movement occurs (though performance can be affected by fatigue or pathology).

SAMPLE

4. Prism adaptation is used in ocular motility management for which scenario?

A. It is used for large strabismus requiring surgical management.

B. It permanently cures diplopia.

C. Prism adaptation refers to gradual improvement in alignment with wearing prisms over time; used for convergence insufficiency or other small misalignments with residual diplopia.

D. It has no clinical use.

Prism adaptation relies on the eye-brain system gradually adjusting to the optical shift caused by wearing prisms. When a small prism is worn, the vergence and fusional systems can recalibrate, slowly aligning the eyes and reducing or eliminating diplopia for the misalignment it addresses. This approach is especially useful in convergence insufficiency and other small misalignments where there is residual diplopia after initial evaluation or therapy. It's not a permanent cure—remove the prisms and the diplopia can return if the underlying motor alignment hasn't changed. It's also not suited for large strabismus, which typically needs surgical or other interventions. So prism adaptation serves as a non-surgical option to improve alignment and guide management in small-angle deviations with residual diplopia.

5. Which reflex is primarily responsible for stabilizing vision during head rotation by producing compensatory eye movements?

A. Optokinetic reflex (OKR)

B. Optokinetic nystagmus (OKN)

C. Vestibulo-ocular reflex (VOR)

D. Pursuit

The vestibulo-ocular reflex stabilizes vision during head rotation by producing eye movements opposite to the direction of head movement. When the head turns, the vestibular apparatus (especially the semicircular canals) detects angular velocity and sends signals to the eyes to move in the opposite direction with roughly the same magnitude. This counter-rotation keeps the image stable on the retina, preserving a clear, steady gaze even as the head moves. Other reflexes respond to different stimuli. The optokinetic reflex and its eye movements (optokinetic nystagmus) are driven by movement of the visual scene across the retina, helping stabilize vision when the environment itself moves. Pursuit is a voluntary, cortical-driven ability to smoothly track a moving target, not a reflex that automatically stabilizes gaze during head motion.

SAMPLE

6. Which test mainly evaluates pursuits?

A. King-Devick

B. DEM

C. Pierce Saccades Tests

D. Groffman Tracings

Smooth pursuit is the ability to smoothly follow a moving target with the eyes. A test that targets this skill presents a moving target the patient tracks, and the clinician looks for how well the eyes maintain continuous tracking, without jumping to catch the target. Groffman Tracings is specifically designed to assess this ability: the patient follows a moving line or outline as it shifts across the page, so you can observe pursuit smoothness, any catch up saccades, and any loss of fixation. This direct focus on maintaining continuous, coordinated eye movement makes it the best choice for evaluating pursuits. The other tests concentrate more on saccades, the quick jumps the eyes make between fixed targets. King-Devick requires rapid ocular jumps as numbers are read aloud, highlighting saccadic speed and accuracy. The Pierce Saccades Tests isolate saccades themselves, and the Developmental Eye Movement test emphasizes saccadic efficiency and visuomotor sequencing, with only incidental involvement of pursuit. So while they provide useful oculomotor information, they're not primarily about smooth pursuit.

7. Oculocephalic testing is also known as which maneuver?

A. Head thrust test

B. Skew deviation test

C. Doll's head maneuver

D. Cover test

Oculocephalic testing assesses the brainstem-mediated vestibulo-ocular reflex by rapidly turning the head to elicit eye movements. The classic name for this maneuver is the Doll's head maneuver. When the head is moved quickly from side to side, the eyes should move in the opposite direction to maintain fixation on a target. This opposite movement indicates an intact reflex pathway. If the eyes do not move opposite the head or appear to move with the head, the reflex is impaired, suggesting brainstem dysfunction. The other tests listed serve different purposes and involve different reflexes or alignment checks: the head thrust test evaluates the vestibulo-ocular reflex for peripheral vestibular function; the skew deviation test assesses vertical misalignment due to central pathology; and the cover test checks ocular alignment and phorias rather than eliciting the oculocephalic reflex.

8. Duane's Retraction syndrome Type 2 is defined by which limitation?

A. Limited abduction

B. Limited adduction

C. Limited abduction and adduction

D. Normal

In Duane retraction syndrome, the pattern of eye movement limitation defines the type. The hallmark across all types is abnormal innervation of the horizontal recti leading to globe retraction on attempted adduction due to co-contraction of the medial and lateral rectus muscles. For this type, the limited movement is in adduction while abduction remains relatively intact. That means the eye struggles to move inward toward the nose, but can move outward normally. This reflects the specific miswiring pattern that predominantly affects the adducting motion, with abduction preserved. So the correct description is limited adduction. Limiting abduction describes the other type, limiting both directions describes another type, and normal movement would not fit this syndrome.

SAMPLE

9. End-point nystagmus is physiological.

A. Pathological

B. Physiological

C. Both

D. Neither

End-point nystagmus is a normal, physiological response that shows up when the eyes are held at the extreme edge of their range. At far gaze, the eye-holding system and the vestibulo-ocular reflex interact in a way that can't perfectly stabilize the eye at that position. A tiny drift toward the center is followed by a quick corrective eye movement, producing a small, rapid nystagmus. This happens in healthy individuals and disappears when gaze is moved away from the end point, so it's not a sign of disease. Pathological nystagmus, by contrast, would not be confined to the extreme gaze and wouldn't be considered a normal finding.

10. Describe the near point of convergence (NPC) test and typical normal values.

A. Move a small target toward the nose along the midline; typical NPC 6–8 cm in adults

B. Move a large target toward the nose; typical NPC 20–30 cm

C. Move target away from the nose; typical NPC 12–15 cm

D. NPC is not used in adults

The near point of convergence tests how well the eyes can turn inward together to keep a near object fused into a single image. A small target is moved along the midline toward the patient, and you note the distance from the nose to the target at which double vision first occurs (the break point). Then you continue moving the target away until single vision returns (the recovery point). In adults, a typical normal NPC is about 6–8 centimeters from the nose. Values longer than this can indicate reduced convergence ability, as seen with convergence insufficiency, while shorter distances are generally normal. This test uses a small target because fine convergence is being measured; a large target or testing divergence (moving away) serves different purposes, so they don't reflect NPC. NPC is relevant in adults and children as part of evaluating binocular vision and comfort with near work.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nbeoocularmotility.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE