NCC NEONATAL INTENSIVE CARE NURSING (RNC-NIC) CERTIFICATION PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nccrncnic.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is true about cephalohematoma in a neonate?

A. It extends across cranial sutures

B. It is a subperitoneal collection of fluid

C. Overlying scalp may not show discoloration

D. It appears immediately after birth

2. Type II pneumocytes are primarily responsible for which essential function in the lungs?

A. Gas exchange

B. Surfactant production

C. Vascular maintenance

D. Alveolar structural integrity

3. What kind of meconium is typically associated with abnormal gut motility?

A. Soft and watery

B. Thin and liquid

C. Thick and tenacious

D. Oily and fragrant

4. Which factor causes the oxygen-hemoglobin dissociation curve to shift to the left?

A. Increased temperature

B. Decreased pH

C. Decreased $PaCO_2$

D. Higher levels of carbon dioxide

5. What type of encephalopathy presents with lethargy and hypotonia?

A. Mild encephalopathy

B. Moderate encephalopathy

C. Severe encephalopathy

D. Mild hypoglycemia

SAMPLE

6. What characterizes disseminated intravascular coagulation (DIC) in infants?

A. Acute increase in coagulation factors

B. Deficiency in coagulation factors and platelets

C. Independently occurring hemorrhagic disease

D. Hypotension without associated illness

7. What does a Lecithin/Sphingomyelin (L/S) ratio of less than 2 typically suggest?

A. Fetal lung maturity

B. Immature lungs

C. Normal fetal development

D. High risk of congenital anomalies

8. Which type of neural tube defect is visible outside the body?

A. Spina bifida occulta

B. Meningocele

C. Myelomeningocele

D. Myeloschisis

9. When does hypernatremia become a concern during the diuretic phase?

A. With too rapid intake of fluids

B. With too rapid reduction in body water

C. With excessive sodium intake

D. With increased GFR

10. What anatomical feature does a cephalohematoma typically not cross?

A. Cranial suture lines

B. Meningeal layers

C. Midline of the skull

D. Soft tissue boundaries

SAMPLE

Answers

SAMPLE

1. C
2. B
3. C
4. C
5. B
6. B
7. B
8. C
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What is true about cephalohematoma in a neonate?

A. It extends across cranial sutures

B. It is a subperitoneal collection of fluid

C. Overlying scalp may not show discoloration

D. It appears immediately after birth

Cephalohematoma is a localized collection of blood between the periosteum and the skull bone in neonates, typically resulting from birth trauma. The statement that overlying scalp may not show discoloration is accurate because, in cases of cephalohematoma, the blood accumulates beneath the periosteum, and the overlying skin may appear normal or show minimal discoloration, unlike other types of head injuries such as caput succedaneum where swelling and discoloration are visible over the scalp. Understanding the characteristics of cephalohematoma helps differentiate it from other head presentations in newborns, which is crucial for proper diagnosis and management. The fine differentiation in presentation can impact monitoring and parental education about what to expect regarding the baby's condition following delivery.

2. Type II pneumocytes are primarily responsible for which essential function in the lungs?

A. Gas exchange

B. Surfactant production

C. Vascular maintenance

D. Alveolar structural integrity

Type II pneumocytes play a crucial role in the lungs by producing surfactant, a substance that reduces surface tension within the alveoli. This surfactant is essential for maintaining the stability of the alveoli during the breathing cycle, preventing their collapse (atelectasis) and ensuring efficient gas exchange. By lowering surface tension, surfactant allows for easier inflation of the alveoli, thus facilitating proper lung function and improving compliance during ventilation. The production of surfactant is particularly important in neonates, where its deficiency can lead to respiratory complications such as neonatal respiratory distress syndrome (NRDS). The other functions listed, while relevant to lung physiology, are not the primary responsibility of Type II pneumocytes. For instance, gas exchange primarily occurs at the level of Type I pneumocytes, which constitute the alveolar epithelium and provide a large surface area for gas diffusion. Vascular maintenance and alveolar structural integrity are more related to other cell types and components within the lung architecture. Therefore, the production of surfactant is indeed the essential and defining function of Type II pneumocytes.

3. What kind of meconium is typically associated with abnormal gut motility?

A. Soft and watery

B. Thin and liquid

C. Thick and tenacious

D. Oily and fragrant

Thick and tenacious meconium is typically associated with abnormal gut motility. This type of meconium suggests a delay in the passage through the intestines, which can occur in conditions such as meconium ileus or other forms of intestinal obstruction. When there is abnormal gut motility, the transit of meconium can be slowed down, resulting in it becoming thick and sticky, indicating that it has not been expelled in a timely manner post-birth. In contrast, soft and watery meconium, thin and liquid meconium, or oily and fragrant meconium would indicate a different state of gut motility and are not typically associated with the complications seen in abnormal motility. For example, soft and watery meconium may indicate a normal passage, while oily meconium might suggest the presence of certain factors like breastfeeding or dietary changes. Each of these other types of meconium texture can reflect a normal physiological response or an alteration due to other factors unrelated to mechanical obstruction or delayed motility.

SAMPLE

4. Which factor causes the oxygen-hemoglobin dissociation curve to shift to the left?

A. Increased temperature

B. Decreased pH

C. Decreased PaCO2

D. Higher levels of carbon dioxide

The oxygen-hemoglobin dissociation curve shifts to the left under conditions that enhance hemoglobin's affinity for oxygen. A key factor that contributes to this shift is a decrease in carbon dioxide levels (PaCO2). When the concentration of carbon dioxide decreases, it results in an increase in blood pH (more alkaline conditions). This increase in pH leads to a higher affinity of hemoglobin for oxygen, meaning hemoglobin holds on to oxygen more tightly. In the context of neonatal care, understanding this relationship is crucial because it can influence oxygen delivery to tissues. For instance, if a neonate is in a state where carbon dioxide levels are low, this could indicate hyperventilation or other metabolic adjustments. Recognizing these physiological responses helps nurses manage appropriate oxygen therapy and monitor the respiratory status of neonates effectively. In contrast, other factors leading to a rightward shift, such as increased temperature, decreased pH (more acidic conditions), or higher levels of carbon dioxide, would indicate a scenario where hemoglobin releases oxygen more readily, which is not the focus of this question.

5. What type of encephalopathy presents with lethargy and hypotonia?

A. Mild encephalopathy

B. Moderate encephalopathy

C. Severe encephalopathy

D. Mild hypoglycemia

Moderate encephalopathy is characterized by symptoms such as lethargy and hypotonia. In this stage of encephalopathy, the infant may appear less alert and exhibit decreased muscle tone. The lethargy suggests a compromised level of consciousness, where the infant is more difficult to arouse than usual, and hypotonia indicates there is reduced muscle strength or resistance to passive movement. These signs reflect a moderate level of brain dysfunction, which can arise from various underlying issues, including metabolic disturbances or hypoxia. In contrast, mild encephalopathy would typically present with only subtle changes in behavior or tone, while severe encephalopathy might be accompanied by more profound neurological deficits, such as seizures or significant unresponsiveness. Mild hypoglycemia is not classified as encephalopathy and generally presents with symptoms like weakness or irritability rather than the specific findings of lethargy and hypotonia associated with moderate encephalopathy.

SAMPLE

6. What characterizes disseminated intravascular coagulation (DIC) in infants?

A. Acute increase in coagulation factors

B. Deficiency in coagulation factors and platelets

C. Independently occurring hemorrhagic disease

D. Hypotension without associated illness

Disseminated intravascular coagulation (DIC) in infants is characterized by a deficiency in coagulation factors and platelets. In DIC, there is an abnormal activation of the coagulation cascade, which leads to widespread clotting in small blood vessels. This increased clotting consumes platelets and clotting factors, resulting in a paradoxical increased risk of bleeding as these components become depleted. Thus, the clinical picture often includes thrombocytopenia (low platelet count) and a deficiency of coagulation factors, which contribute to the hemorrhagic complications commonly associated with DIC. Understanding this deficiency is crucial in managing DIC effectively, as it guides treatment approaches and monitoring in affected infants.

7. What does a Lecithin/Sphingomyelin (L/S) ratio of less than 2 typically suggest?

A. Fetal lung maturity

B. Immature lungs

C. Normal fetal development

D. High risk of congenital anomalies

A Lecithin/Sphingomyelin (L/S) ratio of less than 2 is indicative of immature lungs. In the context of fetal development, this ratio is used to assess lung maturity, particularly in at-risk pregnancies. Lecithin is a phospholipid component of surfactant, which is produced by type II alveolar cells in the lungs. As the fetus approaches term, the production of lecithin increases relative to sphingomyelin, leading to a higher L/S ratio. When the ratio is less than 2, it suggests that there is insufficient production of surfactant, which is essential for keeping the alveoli open and preventing atelectasis (collapse of the lung). This immature lung condition can be a precursor to respiratory distress syndrome (RDS), a common complication in preterm infants due to inadequate surfactant levels. In summary, a low L/S ratio is a key indicator of lung immaturity, highlighting the need for monitoring and potential interventions to support respiratory function in the neonate.

8. Which type of neural tube defect is visible outside the body?

A. Spina bifida occulta

B. Meningocele

C. Myelomeningocele

D. Myeloschisis

Myelomeningocele is a type of neural tube defect that is characterized by the herniation of the spinal cord and its protective covering (the meninges) through a defect in the bony encasement of the spinal column. This defect leads to a notable protrusion that is visible outside the body, distinguishing it from other types of neural tube defects. In myelomeningocele, the spinal cord is not only exposed but also often contains nerve tissue, which can lead to significant disability and complications. The visible sac contains both the meninges and the neural elements, making this condition evident upon physical examination. In contrast, spina bifida occulta is a mild form of spina bifida where there is an incomplete closure of the spine, but it does not result in any external protrusion, making it not visible from the outside. Meningocele involves the protrusion of the meninges without neural tissue involvement, which can also result in a visible sac, though less frequently compared to myelomeningocele. Myeloschisis is a more severe defect where the spinal cord is completely open and exposed, but it does not typically present as a defined sac that is visible externally.

SAMPLE

9. When does hypernatremia become a concern during the diuretic phase?

A. With too rapid intake of fluids

B. With too rapid reduction in body water

C. With excessive sodium intake

D. With increased GFR

Hypernatremia becomes a concern during the diuretic phase primarily due to the rapid reduction in body water. During this phase, there is significant diuresis, leading to the loss of water from the body. If fluid loss occurs without appropriate fluid replacement, it can result in a higher concentration of sodium in the bloodstream, thus elevating serum sodium levels. This is particularly critical in neonatal care, where maintaining proper fluid balance is essential for preventing complications. In this context, while factors such as increased sodium intake or changes in glomerular filtration rate (GFR) can impact sodium balance, they are not direct triggers for hypernatremia during the diuretic phase. Instead, the focus is on the body's ability to retain water, and when that ability is impaired or overwhelmed by rapid fluid loss, hypernatremia becomes a significant concern. Additionally, in this phase, careful monitoring of fluid intake and electrolyte balance is crucial to avoid adverse effects associated with elevated sodium levels.

10. What anatomical feature does a cephalohematoma typically not cross?

A. Cranial suture lines

B. Meningeal layers

C. Midline of the skull

D. Soft tissue boundaries

A cephalohematoma is a collection of blood that forms between the periosteum and the skull in a neonate, typically resulting from birth trauma. One key characteristic of a cephalohematoma is its confined nature, which means it does not cross cranial suture lines. This is due to the anatomical structure of the skull, where the periosteum is tightly adhered to the bones at the sutures. As a result, any blood accumulation remains localized within the area bounded by the sutures. The recognition of this feature is important in clinical practice because it helps differentiate a cephalohematoma from other types of hematomas, such as a caput succedaneum, which can cross suture lines as it is located above the periosteum in the subcutaneous tissue. Understanding this distinction aids neonatal intensive care nurses and other healthcare providers in making accurate assessments and determining the appropriate management for infants presenting with head swelling after delivery.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nccrncnic.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE