
NCCAA RE-CERTIFICATION PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nccaarecertification.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. **Which of the following is NOT a sign of hypovolemia?**

   A. Increased heart rate

   B. Decreased blood pressure

   C. Increased central venous pressure (CVP)

   D. Decreased urine output

2. **Which condition is characterized by decreased RV compliance as seen on a CVP tracing?**

   A. Atrial flutter

   B. Acute coronary syndrome

   C. Tricuspid or pulmonic stenosis

   D. Decreased cardiac output

3. **What type of nerve fibers are primarily involved in proprioception?**

   A. A-beta type and C-type fibers

   B. A-delta type and A-alpha type

   C. A-alpha type and A-beta type

   D. C-type and A-delta type

4. **Cannon A waves on a CVP tracing are characteristic of which condition?**

   A. Tricuspid stenosis

   B. Atrial fibrillation

   C. Tricuspid regurgitation

   D. Pulmonary hypertension

5. **Which change would NOT increase the rate of inhalational induction?**

   A. Increased alveolar ventilation

   B. Utilizing an inotropic infusion

   C. Substitution of Sevoflurane for Isoflurane

   D. Delivering anesthesia in different altitudes

6. **What individuals would most likely experience stridor after extubation?**

   A. Patients with obesity

   B. Patients with BP fluctuations

   C. Patients with bilateral RLN damage

   D. Patients with obstructive sleep apnea

SAMPLE

7. What could be a possible cause for a sudden decrease in ETCO2 during a total hip replacement?

A. Hypothermia

B. Tourniquet release

C. Venous air emboli

D. Endobronchial migration of the ETT

8. Which description about alveoli distribution is correct?

A. Alveoli are largest at the base of the lung.

B. The smallest alveoli are found at the apex of the lung.

C. The largest alveoli are found throughout the lung.

D. Alveoli are smallest at the base of the lung.

9. What vital sign is most likely to indicate the presence of malignant hyperthermia?

A. Decreased ETCO2

B. Increased muscle rigidity

C. Increased ETCO2

D. Increased blood pressure

10. Which agent is known for having the greatest antimuscarinic/vagolytic actions?

A. Gallamine

B. Succinylcholine

C. Pancuronium

D. D-Tubocurarine

SAMPLE

Answers

SAMPLE

1. C
2. C
3. C
4. D
5. B
6. C
7. C
8. D
9. C
10. A

SAMPLE

Explanations

SAMPLE

1. Which of the following is NOT a sign of hypovolemia?

A. Increased heart rate

B. Decreased blood pressure

C. Increased central venous pressure (CVP)

D. Decreased urine output

The correct answer is that increased central venous pressure (CVP) is not a sign of hypovolemia. Hypovolemia, which refers to a state of decreased blood volume, typically leads to physiological responses that aim to maintain blood circulation and compensate for the reduced volume. When the body experiences hypovolemia, compensatory mechanisms are triggered, often resulting in increased heart rate and decreased blood pressure. These changes are part of a response to maintain adequate perfusion to vital organs. Concurrently, kidney function may be affected due to reduced blood volume, often resulting in decreased urine output as the body attempts to conserve fluid. In contrast, elevated CVP usually indicates that there is an excess of fluid in the circulatory system, often related to conditions such as heart failure, fluid overload, or increased venous return. Thus, in cases of hypovolemia, we would expect the CVP to be low rather than high, supporting the assertion that increased CVP is not a hallmark of hypovolemia.

2. Which condition is characterized by decreased RV compliance as seen on a CVP tracing?

A. Atrial flutter

B. Acute coronary syndrome

C. Tricuspid or pulmonic stenosis

D. Decreased cardiac output

The correct answer is tricuspid or pulmonic stenosis. This condition is characterized by an obstruction to blood flow across the right heart valves, resulting in decreased right ventricular (RV) compliance. When the right ventricle cannot adequately fill due to the narrowed valve opening, it leads to increased pressure in the right atrium and subsequently influences the central venous pressure (CVP) tracing. In CVP monitoring, this will manifest as a plateau or a more pronounced "wedge" appearance on the tracing due to the elevated pressures. When assessing how tricuspid or pulmonic stenosis impacts the right ventricle, it is important to recognize that the heart's ability to expand and contract effectively is compromised. The resulting back pressure can lead to systemic venous congestion as blood backs up into the systemic circulation, reflecting abnormal compliance on CVP tracing. The other options presented do not specifically tie to a direct decrease in RV compliance as seen in the CVP readings. For example, atrial flutter primarily affects the atrial rhythm without necessarily impacting ventricular compliance. Acute coronary syndrome involves myocardial ischemia that affects the heart muscle itself but does not inherently describe the hemodynamic profile seen in CVP tracing caused by valvular stenosis. De

SAMPLE

3. What type of nerve fibers are primarily involved in proprioception?

A. A-beta type and C-type fibers

B. A-delta type and A-alpha type

C. A-alpha type and A-beta type

D. C-type and A-delta type

Proprioception is the body's ability to sense its position and movement in space, which is crucial for coordination and balance. The primary nerve fibers involved in proprioception are A-alpha and A-beta type fibers. A-alpha fibers are large-diameter myelinated fibers that primarily transmit signals from muscle spindles and Golgi tendon organs. These receptors provide information about muscle stretch and tension, which are essential for maintaining posture and executing movements. A-beta fibers, also myelinated but slightly smaller than A-alpha, carry information from mechanoreceptors that sense touch and pressure, and they play a role in proprioception by transmitting signals about the position of skin and joints. Together, these fiber types effectively relay proprioceptive information to the central nervous system, allowing for immediate response and adjustment of muscle activity necessary for balance and coordination. Thus, the combination of A-alpha and A-beta fibers is crucial for the proprioceptive feedback mechanism that helps maintain body awareness and control during movement activities.

4. Cannon A waves on a CVP tracing are characteristic of which condition?

A. Tricuspid stenosis

B. Atrial fibrillation

C. Tricuspid regurgitation

D. Pulmonary hypertension

Cannon A waves on a central venous pressure (CVP) tracing indicate the presence of a significant atrioventricular (AV) dissociation, which is often associated with conditions affecting the heart's rhythm and function. The correct answer reflects that cannon A waves can occur in the context of pulmonary hypertension due to the increased pressure in the pulmonary arteries, which leads to changes in the dynamics of right atrial contraction. In conditions like pulmonary hypertension, the right atrium may contract against a higher resistance, and when there is dissociation between the atria and ventricles (as can occur in certain arrhythmias), this will result in cannon A waves, which are prominent and occur simultaneously with the ventricular contraction rather than following it. The other choices reflect different pathophysiological mechanisms that do not specifically lead to the occurrence of cannon A waves in the same way. Tricuspid stenosis, for example, can cause elevated right atrial pressure but is more likely to lead to a different wave form profile, primarily steep A waves due to impaired filling of the right ventricle. Atrial fibrillation also typically presents with irregularly timed and absent A waves on CVP tracing, rather than the pronounced cannon A waves associated with AV

SAMPLE

5. Which change would NOT increase the rate of inhalational induction?

A. Increased alveolar ventilation

B. Utilizing an inotropic infusion

C. Substitution of Sevoflurane for Isoflurane

D. Delivering anesthesia in different altitudes

The choice that would not increase the rate of inhalational induction is the utilization of an inotropic infusion. Inotropic agents primarily influence the heart's contractility and cardiac output rather than the dynamics of inhalational anesthetic uptake and distribution. Inhalational induction refers to the process of administering anesthesia agents through inhalation. Factors that typically increase the rate of induction include enhancing the agent's delivery to the alveoli and improving its absorption into the bloodstream. Increased alveolar ventilation, for example, raises the concentration gradient between the alveoli and the blood, leading to faster absorption of the anesthetic agent. The substitution of Sevoflurane for Isoflurane is also relevant because Sevoflurane has a lower blood-gas partition coefficient, which means it is less soluble in blood compared to Isoflurane. This characteristic results in quicker delivery to the brain and, therefore, a faster induction of anesthesia. Delivering anesthesia at different altitudes can affect the partial pressures of gases and potentially alter the anesthetic induction time, but these factors are more complex and variable. Thus, using an inotropic infusion does not directly influence pulmonary ventilation or gas exchange and does not enhance inhalational induction rates like the other options do.

6. What individuals would most likely experience stridor after extubation?

A. Patients with obesity

B. Patients with BP fluctuations

C. Patients with bilateral RLN damage

D. Patients with obstructive sleep apnea

Stridor is a high-pitched wheezing sound that occurs when there is a narrowing or obstruction of the airway. After extubation, patients may experience stridor due to various factors related to airway edema or injury. Individuals with bilateral recurrent laryngeal nerve (RLN) damage are most likely to experience stridor after extubation because this nerve is crucial for the control of the vocal cords. Damage to both sides of the RLN can lead to the inability of the vocal cords to adduct properly, which is necessary for maintaining airway patency. This can result in significant airway obstruction, causing stridor as the patient breathes. In contrast, while patients with obesity and obstructive sleep apnea may have airway management challenges, their risk factors do not directly relate to the type of airway compromise indicated by stridor following extubation. Patients with blood pressure fluctuations may also experience a variety of complications, but stridor is specifically linked to airway mechanics and vocal cord function, which is primarily affected by RLN integrity. Thus, individuals with bilateral RLN damage present a clear pathway to developing stridor after the removal of an endotracheal tube.

SAMPLE

7. What could be a possible cause for a sudden decrease in ETCO2 during a total hip replacement?

A. Hypothermia

B. Tourniquet release

C. Venous air emboli

D. Endobronchial migration of the ETT

The chosen answer points to a significant concern in the context of anesthesia and surgical procedures, particularly during a total hip replacement. A sudden decrease in end-tidal carbon dioxide (ETCO2) can indicate a rapid change in the patient's physiology, particularly in the lungs' ability to eliminate carbon dioxide. When venous air emboli occur, air enters the venous bloodstream, leading to an obstruction in the pulmonary circulation. This can drastically reduce the amount of blood reaching the lungs and, consequently, the carbon dioxide being expelled from the body during exhalation. The result is a sharp drop in the ETCO2 levels since there is less carbon dioxide being eliminated from the bloodstream. This condition can lead to serious complications, including hypotension, hypoxemia, and potential cardiovascular collapse, making it a critical situation that requires immediate attention. Other options, while relevant, do not account for a sudden decrease in ETCO2 in the same way. Hypothermia can affect metabolic rates and ventilation but does not typically cause a rapid drop in carbon dioxide levels. Tourniquet release could lead to a temporary increase in ETCO2 due to the sudden return of blood flow from the extremity, rather than a decrease. Endobronchial migration of

8. Which description about alveoli distribution is correct?

A. Alveoli are largest at the base of the lung.

B. The smallest alveoli are found at the apex of the lung.

C. The largest alveoli are found throughout the lung.

D. Alveoli are smallest at the base of the lung.

The claim that alveoli are smallest at the base of the lung is supported by the understanding of lung anatomy and physiology. Alveoli, the tiny air sacs where gas exchange occurs, vary in size depending on their location within the lungs. At the apex of the lungs, the alveoli are generally larger. This larger size is due to the lower volume of air pressure and distribution at higher altitudes of the lung tissue. Conversely, as we move toward the base of the lungs, the alveoli become smaller; this is because they are compressed by the weight of the lung tissue above them as well as the greater blood flow and expansion of lung volume that occurs during inhalation at the base. This anatomical distribution allows for optimal gas exchange, as the larger alveoli can accommodate greater volumes of air and facilitate gas exchange more effectively in regions of the lung that receive more ventilation. Understanding this distribution is essential for grasping how the respiratory system functions efficiently.

SAMPLE

9. What vital sign is most likely to indicate the presence of malignant hyperthermia?

A. Decreased ETCO2

B. Increased muscle rigidity

C. Increased ETCO2

D. Increased blood pressure

Malignant hyperthermia is a severe reaction typically triggered by certain general anesthetics and neuromuscular blockers, leading to a hypermetabolic state in skeletal muscle. One of the key indicators of this condition is an increase in end-tidal carbon dioxide (ETCO2) levels. During malignant hyperthermia, the body's metabolism accelerates, resulting in increased carbon dioxide production as muscles consume more energy. As the body's ability to eliminate CO2 through ventilation is overwhelmed by the excessive production, the ETCO2 levels rise significantly. This is a critical sign that helps clinicians recognize the onset of malignant hyperthermia, prompting immediate treatment interventions to mitigate the condition. The other options, while they may be present in various clinical scenarios, do not specifically signify malignant hyperthermia in the same way that increased ETCO2 does. Increased muscle rigidity is a common sign of the condition, but it does not serve as a direct measure of the metabolic changes that are occurring. Therefore, the increase in ETCO2 is directly linked to the hypermetabolic state and is a vital sign healthcare providers monitor closely during anesthesia to identify and manage malignant hyperthermia effectively.

10. Which agent is known for having the greatest antimuscarinic/vagolytic actions?

A. Gallamine

B. Succinylcholine

C. Pancuronium

D. D-Tubocurarine

Gallamine is recognized for having the most significant antimuscarinic or vagolytic actions among the agents listed. Antimuscarinic effects refer to the ability of a drug to block the action of acetylcholine at muscarinic receptors, which are a type of cholinergic receptor found in many tissues throughout the body, including the heart and smooth muscle. By inhibiting these receptors, gallamine can lead to increased heart rate and reduced glandular secretions, which is beneficial in certain clinical scenarios where bradycardia or excessive secretions need to be managed during anesthesia. In contrast, while succinylcholine is a neuromuscular blocker, it primarily functions as a depolarizing muscle relaxant and does not have significant antimuscarinic effects. Pancuronium is another neuromuscular blocker but has less pronounced vagolytic actions compared to gallamine. D-Tubocurarine, though a neuromuscular blocker as well, is also associated with a lower degree of antimuscarinic activity. Therefore, gallamine stands out for its ability to not only block neuromuscular transmission but also to exert pronounced vagolytic effects, making it the correct choice in this context.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nccaarecertification.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE