
NCFE DENTAL NURSING PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://ncfedentalnursing.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which sign of inflammation typically occurs last?

A. Heat

B. Loss of function

C. Pain

D. Swelling

2. What personal protective equipment is essential to wear during procedures that can generate aerosols?

A. Face mask

B. Gloves

C. Goggles

D. All of the above

3. Which one of the following materials is not used to cement a molar band to a tooth as part of a fixed orthodontic appliance?

A. Composite filler

B. Glass ionomer

C. Zinc phosphate

D. Zinc polycarboxylate

4. What is the recommended storage practice for sterilised instruments?

A. Store in open containers

B. Keep in a dry, clean area until needed

C. Place in a wet environment for cooling

D. Wrap in plastic before sterilisation

5. What is the correct classification of a mesio-palatal cavity in the upper right lateral incisor?

A. Class I

B. Class II

C. Class III

D. Class IV

SAMPLE

6. Which one of the following actions should be the final stage of a risk assessment in the dental workplace?

A. Control the risk

B. Evaluate the risk

C. Identify the risk

D. Identify those at risk

7. What is the fundamental principle of standard (universal) precautions in healthcare?

A. All dental personnel will be free from disease

B. All patients will have received vaccinations

C. Any patient may be infected with a pathogen at any time

D. Any patient may have natural immunity

8. What is the correct term for a yellow-coloured, hard deposit often visible in the oral cavity due to poor hygiene?

A. Biofilm

B. Plaque

C. Subgingival calculus

D. Supragingival calculus

9. Which medical emergency drug may be administered to assist a casualty who is suffering an anaphylactic reaction?

A. Adrenaline

B. Aspirin

C. Midazolam

D. Salbutamol

10. Which one of the following is an effect that often occurs in older patients due to age-related changes to their teeth?

A. Decreased tooth sensitivity

B. Occlusal caries

C. Root curvature

D. Whitened teeth

SAMPLE

Answers

SAMPLE

1. B
2. D
3. A
4. B
5. C
6. A
7. C
8. D
9. A
10. A

SAMPLE

Explanations

SAMPLE

1. Which sign of inflammation typically occurs last?

A. Heat

B. Loss of function

C. Pain

D. Swelling

Loss of function is considered to be the last sign of inflammation due to its development in response to the preceding signs of inflammation, such as heat, pain, and swelling. Inflammation is the body's immune response to injury or infection, and these early signs function to alert the body to a problem and to promote healing. Heat occurs as blood vessels dilate and more blood flows to the injured area, causing warmth. Pain arises from the release of inflammatory chemicals and the direct effect of swelling on nearby nerves. Swelling, or edema, results from increased vascular permeability, leading to the accumulation of fluid in tissues. Loss of function, however, often results as a cumulative effect of these preceding signs. When pain and swelling limit movement or function in a particular area, it ultimately contributes to the functional impairment observed. Therefore, loss of function emerges after the other signs have already developed and is often the last symptom to manifest during the inflammatory response.

2. What personal protective equipment is essential to wear during procedures that can generate aerosols?

A. Face mask

B. Gloves

C. Goggles

D. All of the above

Wearing personal protective equipment (PPE) during procedures that generate aerosols is crucial for ensuring the safety of dental professionals and patients alike. The correct choice emphasizes the importance of a comprehensive approach to PPE, which includes a combination of face masks, gloves, and goggles. Face masks are essential in preventing the inhalation of airborne particles and aerosols that may contain infectious agents. Masks act as a barrier to protect the respiratory system, which is particularly vital during dental procedures where aerosols are produced. Gloves serve a different yet equally important purpose by providing a protective barrier for the hands against blood, saliva, and other potentially infectious materials. This protects both the clinician and the patient, ensuring that the risk of cross-contamination is minimized. Goggles protect the eyes from splashes and potential contact with infectious aerosols. Eye protection is important as mucous membranes of the eyes can be a portal of entry for pathogens. Thus, the correct answer underscores the necessity of utilizing all three types of personal protective equipment during aerosol-generating procedures to provide comprehensive protection and adhere to best practices in infection control within dental settings.

SAMPLE

3. Which one of the following materials is not used to cement a molar band to a tooth as part of a fixed orthodontic appliance?

A. Composite filler

B. Glass ionomer

C. Zinc phosphate

D. Zinc polycarboxylate

Composite filler is not typically used to cement a molar band to a tooth as part of a fixed orthodontic appliance. The primary reason for this is that composite materials are generally used for bonding or filling purposes rather than for cementing orthodontic bands, which require materials that provide a strong and durable bond capable of withstanding significant forces during orthodontic treatment. In orthodontics, the materials commonly used for cementing molar bands include glass ionomer, zinc phosphate, and zinc polycarboxylate. Glass ionomer cement bonds chemically to both the tooth structure and the metal band, offering good retention, fluoride release, and ease of handling. Zinc phosphate cement has a long history of use, providing excellent strength and a reliable bond. Zinc polycarboxylate cement, while slightly less strong than zinc phosphate, is still effective and offers a degree of bonding to tooth structure. Therefore, while composite filler can have its applications in dentistry, particularly for restorative purposes, it is not the material of choice for cementing molar bands in an orthodontic setting.

4. What is the recommended storage practice for sterilised instruments?

A. Store in open containers

B. Keep in a dry, clean area until needed

C. Place in a wet environment for cooling

D. Wrap in plastic before sterilisation

Storing sterilised instruments in a dry, clean area until they are needed is the best practice because it helps maintain their sterility. A dry environment minimizes the risk of contamination from moisture, which can promote the growth of bacteria or other pathogens. Additionally, keeping the instruments clean and away from contaminants in a designated storage area ensures that the value of the sterilisation process is preserved until the instruments are required for use in dental procedures. Proper storage is critical in maintaining the integrity and safety of sterilised instruments, emphasizing the need for a controlled environment that protects them from potential sources of contamination. This practice helps prevent any factors that could compromise the sterility of the instruments, ensuring that they are safe and effective when needed.

5. What is the correct classification of a mesio-palatal cavity in the upper right lateral incisor?

A. Class I

B. Class II

C. Class III

D. Class IV

A mesio-palatal cavity in the upper right lateral incisor is classified as Class III. This classification is appropriate because Class III cavities involve caries that affect the interproximal area of an anterior tooth without extending to the incisal edge. Since the cavity in question is located on the mesial (toward the midline of the face) and palatal (the side facing the palate) surfaces of the lateral incisor, it fits the criteria for a Class III lesion. Class I cavities involve the occlusal surfaces of posterior teeth, Class II cavities refer to caries that affect the proximal surfaces of posterior teeth, and both Class IV and Class III are associated with anterior teeth, but Class IV specifically involves the incisal edge, which isn't applicable here. Therefore, understanding the specifics of cavity classification helps in accurate diagnosis and treatment planning in dental practice.

SAMPLE

6. Which one of the following actions should be the final stage of a risk assessment in the dental workplace?

A. Control the risk

B. Evaluate the risk

C. Identify the risk

D. Identify those at risk

The final stage of a risk assessment in the dental workplace is to control the risk. After a thorough assessment process that includes identifying the risk, evaluating its significance, and determining who may be affected, the focus shifts to implementing measures to eliminate or mitigate those risks. Controlling the risk is essential because it translates the assessment into actionable steps that enhance safety for both patients and staff. This can include implementing safety protocols, using personal protective equipment, or modifying procedures to reduce the likelihood of an incident. Effectively controlling the risk ensures that the identified hazards are managed appropriately, resulting in a safer dental environment. This is the crucial final step that brings the risk assessment process to fruition by addressing the potential dangers identified earlier.

7. What is the fundamental principle of standard (universal) precautions in healthcare?

A. All dental personnel will be free from disease

B. All patients will have received vaccinations

C. Any patient may be infected with a pathogen at any time

D. Any patient may have natural immunity

The fundamental principle of standard (universal) precautions in healthcare is that any patient may be infected with a pathogen at any time. This principle underscores the importance of treating all patients as potentially infectious, regardless of their known medical history or appearance. By adopting this approach, healthcare providers, including dental personnel, can minimize the risk of transmission of infections. Standard precautions encompass a range of practices aimed at protecting both patients and healthcare workers. These practices include the use of personal protective equipment (PPE), proper hand hygiene, and safe handling of instruments and waste. By operating under the assumption that all patients may harbor infectious agents, dental professionals are better equipped to prevent cross-contamination and ensure a safe environment for everyone. Other options may suggest ideal scenarios regarding disease-free personnel or vaccinated patients, but they do not reflect the reality of clinical practice where infections can be present without visible symptoms. The option about natural immunity also does not align with the precautionary principle, as it does not address the potential risks associated with unknowing carriers of infections.

SAMPLE

8. What is the correct term for a yellow-coloured, hard deposit often visible in the oral cavity due to poor hygiene?

A. Biofilm

B. Plaque

C. Subgingival calculus

D. Supragingival calculus

The term "supragingival calculus" refers to the hard yellowish deposit that often forms above the gum line due to inadequate oral hygiene. This type of calculus develops from mineralized dental plaque, which is composed of bacteria, food particles, and saliva. When plaque is not regularly removed through brushing and flossing, it can harden and become calculus. Supragingival calculus is typically found on the teeth, especially on the buccal surfaces of the molars and the lingual surfaces of the lower incisors, where saliva flows. Its presence is a visible indicator of poor oral hygiene practices, as it becomes more prominent when brushing and flossing are neglected. In contrast, biofilm refers to a thin, sticky film of bacteria that forms on teeth, but it is not specifically a hard deposit and can describe a wider range of microbial growth. Plaque is a soft, sticky accumulation of bacteria that can lead to the formation of calculus if not cleaned properly; however, it is not the hard deposit itself. Subgingival calculus forms below the gum line and may not be visible without clinical examination, distinguishing it from the supragingival type that is readily seen.

9. Which medical emergency drug may be administered to assist a casualty who is suffering an anaphylactic reaction?

A. Adrenaline

B. Aspirin

C. Midazolam

D. Salbutamol

The administration of adrenaline is crucial during an anaphylactic reaction because it acts as a powerful bronchodilator and vasoconstrictor. Anaphylaxis can lead to severe respiratory distress and cardiovascular collapse, and adrenaline quickly counteracts these symptoms. By stimulating alpha-1 adrenergic receptors, adrenaline causes vasoconstriction, which increases blood pressure and reduces swelling, while stimulation of beta-2 adrenergic receptors helps to relax the bronchial smooth muscles, easing breathing difficulties. While other medications such as aspirin serve different purposes, primarily for pain relief or anti-inflammatory effects, they do not address the immediate life-threatening aspects of anaphylaxis. Midazolam is a sedative used primarily for seizure management or anxiety but does not have any role in treating anaphylaxis. Salbutamol is a bronchodilator that can help with breathing issues but is not a first-line treatment for the systemic effects of anaphylaxis. Thus, adrenaline remains the definitive treatment for this medical emergency.

SAMPLE

10. Which one of the following is an effect that often occurs in older patients due to age-related changes to their teeth?

A. Decreased tooth sensitivity

B. Occlusal caries

C. Root curvature

D. Whitened teeth

The correct alternative highlights a common phenomenon among older individuals that leads to a decrease in tooth sensitivity. As people age, several factors contribute to this reduction. One significant factor is the natural wear and tear on the teeth, including the thickening of the enamel and the secondary dentin formation. This process can provide a sort of protective barrier, making the nerves less responsive to stimuli such as temperature changes or certain foods. In contrast, other options represent conditions that may not typically decrease with age. Occlusal caries, for instance, are a type of cavity commonly found in older adults due to various factors, including prolonged exposure to plaque and a decrease in oral hygiene effectiveness. Root curvature and the whitening of teeth are also not universally accepted outcomes of aging, with root curvature being more associated with individual anatomical variations rather than a direct consequence of aging. Whitening of teeth generally becomes less common as staining occurs and enamel thins, resulting in a yellower appearance instead. Thus, understanding the biological changes occurring in the dentition as patients age is crucial, and the decrease in tooth sensitivity is a notable effect often observed in older demographics.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://ncfedentalnursing.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE