
# NCLEX HEART FAILURE PRACTICE TEST (SAMPLE)

## STUDY GUIDE



## SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

**VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE**

https://nclexheartfailure.examzify.com

**Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.**

**ALL RIGHTS RESERVED.**

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

# Table of Contents


Copyright ..... 1

Table of Contents ..... 2

Introduction ..... 3

How to Use This Guide ..... 4

Questions ..... 5

Answers ..... 8

Explanations ..... 10

Next Steps ..... 15


SAMPLE

# Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

# How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

### 1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

### 2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

### 3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

### 4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

### 5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

### 6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

# Questions

SAMPLE

**1. How do NSAIDs affect heart failure management?**

A. They promote fluid retention, worsen hypertension, and impair renal function.

B. They improve diuresis and lower blood pressure.

C. They have no effect on HF.

D. They improve renal function.

**2. Which medication increases the patient's risk of Digoxin toxicity by promoting potassium loss?**

A. Furosemide

B. Metformin

C. Nitroglycerin

D. Coumadin

**3. Which device therapy is indicated for reducing mortality in some heart failure patients and may require biventricular pacing?**

A. Implantable cardioverter-defibrillator

B. Cardiac resynchronization therapy

C. Left ventricular assist device

D. Pacemaker

**4. What emerging HF therapy class provides mortality and hospitalization benefits beyond glucose control, even in non-diabetics?**

A. ACE inhibitors

B. Beta blockers

C. SGLT2 inhibitors

D. ARBs

**5. Which electrolyte requires monitoring due to diuretic and RAAS therapy?**

A. Sodium

B. Potassium

C. Calcium

D. Magnesium

SAMPLE

**6. Which agent is used to acutely reduce preload in pulmonary edema?**

A. IV Nitroglycerin (or other IV nitrates)

B. IV Furosemide

C. Morphine

D. Nifedipine

**7. Which symptom is most characteristic of left-sided heart failure?**

A. Jugular venous distention

B. Persistent cough with crackles in the lungs

C. Peripheral edema

D. Ascites

**8. A patient taking Digoxin has severe bradycardia, nausea, and vomiting, with a Digoxin level of 4 ng/mL. What medication would the physician order?**

A. Narcan

B. Aminophylline

C. Digibind

D. No medication because this is a normal level.

**9. Which of the following actions is most appropriate if a patient on Digoxin reports vision changes?**

A. Ignore as it is unlikely related to Digoxin

B. Advise the patient to rest and retake the dose later

C. Instruct to report immediately to the clinician

D. Resume dosing only after Potassium normalizes

**10. In the hospital management of heart failure, inotropes are used to:**

A. Cure HF

B. Improve cardiac output in patients with low output despite optimized therapy or in shock

C. Increase preload

D. Worsen heart failure

SAMPLE

# Answers

SAMPLE

1. A
2. A
3. B
4. C
5. B
6. A
7. B
8. C
9. C
10. B

SAMPLE

# Explanations

SAMPLE

## 1. How do NSAIDs affect heart failure management?

A. They promote fluid retention, worsen hypertension, and impair renal function.

B. They improve diuresis and lower blood pressure.

C. They have no effect on HF.

D. They improve renal function.

*NSAIDs can worsen heart failure by disrupting kidney function and fluid balance. They block prostaglandin synthesis, which reduces the ability of the kidneys to maintain adequate blood flow, especially when perfusion is already challenged in heart failure. Prostaglandins help dilate the afferent arteriole and support glomerular filtration; without them, the kidney's filtration rate can drop, leading to fluid and salt retention and higher blood pressure. This fluid buildup increases preload and can worsen edema and symptoms of HF, sometimes precipitating decompensation. In addition, reduced renal function makes diuretics less effective, complicating management further. For these reasons, NSAIDs are generally avoided in heart failure; if pain relief is needed, acetaminophen is preferred, and if NSAIDs must be used, they should be used at the lowest dose for the shortest time with close monitoring of renal function, weight, and blood pressure.*

## 2. Which medication increases the patient's risk of Digoxin toxicity by promoting potassium loss?

A. Furosemide

B. Metformin

C. Nitroglycerin

D. Coumadin

*The key idea is that low potassium levels increase the risk of digoxin toxicity. Loop diuretics like furosemide promote potassium loss in the distal tubule, which can lead to hypokalemia. Digoxin works by inhibiting the Na+/K+ ATPase pump in cardiac cells; when potassium is low, digoxin binds more readily to this pump, amplifying its effects. That heightened inhibition causes more calcium to accumulate inside heart cells, which can trigger dangerous arrhythmias and other toxicity symptoms. So a medication that causes potassium wasting directly raises the chance of digoxin toxicity, making furosemide the correct choice. The other drugs listed don't promote potassium loss or directly increase digoxin toxicity through this mechanism. Metformin, nitroglycerin, and warfarin do not affect potassium or the digoxin–pump interaction in the way loop diuretics do.*

## 3. Which device therapy is indicated for reducing mortality in some heart failure patients and may require biventricular pacing?

A. Implantable cardioverter-defibrillator

B. Cardiac resynchronization therapy

C. Left ventricular assist device

D. Pacemaker

*Cardiac resynchronization therapy targets the problem of ventricular dyssynchrony in heart failure. By delivering pacing stimuli to both ventricles (biventricular pacing), it synchronizes their contractions, which improves the heart's pumping efficiency and can reduce mortality and hospitalizations in patients with reduced ejection fraction and a wide QRS complex, especially with left bundle branch block. Some CRT devices also include defibrillation capabilities (CRT-D), adding protection against sudden arrhythmic death while still providing resynchronization. Other devices don't address dyssynchrony: an implantable cardioverter-defibrillator reduces sudden death risk through shocks but isn't primarily about coordinating ventricular timing; a left ventricular assist device provides mechanical support for advanced heart failure; a pacemaker alone paces the heart but doesn't correct dyssynchrony to the same extent.*

SAMPLE

**4. What emerging HF therapy class provides mortality and hospitalization benefits beyond glucose control, even in non-diabetics?**

A. ACE inhibitors

B. Beta blockers

C. SGLT2 inhibitors

D. ARBs

*SGLT2 inhibitors are the class that fit this description. These drugs, originally developed to lower blood glucose in diabetes, have shown clear heart failure benefits that extend beyond glycemic control and apply even to people without diabetes. Key trials in heart failure demonstrated meaningful reductions in death and hospitalization for heart failure with agents like dapagliflozin and empagliflozin in patients with reduced ejection fraction, and similar benefits in many with preserved ejection fraction. Importantly, these benefits occurred regardless of diabetes status, highlighting effects beyond glucose lowering. The benefits come from multiple mechanisms beyond lowering glucose: they promote natriuresis and diuresis, reduce preload and afterload, improve renal function and kidney-heart crosstalk, and may enhance myocardial energy efficiency and reduce inflammation. This combination translates into lower mortality and fewer heart failure hospitalizations, which is why this class is viewed as an important and relatively new advance in heart failure therapy for diverse patient populations.*

**5. Which electrolyte requires monitoring due to diuretic and RAAS therapy?**

A. Sodium

B. Potassium

C. Calcium

D. Magnesium

*Potassium is the electrolyte to monitor because diuretics, especially loop and thiazide types, increase potassium loss, while RAAS therapies (ACE inhibitors, ARBs, and particularly aldosterone antagonists like spironolactone or eplerenone) promote potassium retention. In heart failure care, patients are commonly on both diuretics to relieve fluid overload and RAAS inhibitors to improve outcomes, which can push potassium levels in opposite directions. This interplay makes potassium balance the most critical to watch to prevent dangerous arrhythmias or severe electrolyte disturbances. Sodium, calcium, and magnesium can be affected too, but the combination of diuretic-induced loss and RAAS-mediated retention most directly targets potassium. Regular monitoring of potassium helps guide dose adjustments and safeguard the patient.*

SAMPLE

## 6. Which agent is used to acutely reduce preload in pulmonary edema?

A. IV Nitroglycerin (or other IV nitrates)

B. IV Furosemide

C. Morphine

D. Nifedipine

*The key idea is to rapidly lower the amount of blood returning to the heart, which reduces left-sided filling pressures and pulmonary congestion in acute pulmonary edema. The quickest way to achieve this is venodilation, which decreases venous return (preload) to the heart. IV nitroglycerin acts as a potent venodilator, so it rapidly lowers preload and left ventricular end-diastolic pressure, which directly reduces pulmonary capillary wedge pressure and edema. This helps relieve dyspnea quickly and improves oxygenation. Diuretics like IV furosemide also reduce preload by removing fluid, but their effect takes longer to develop, so they're not as immediate for acute relief. Morphine can cause venodilation and anxiolysis, but its routine use for preload reduction is less favored today due to potential adverse effects and inconsistent outcomes. Calcium channel blockers such as nifedipine can lower BP and don't provide the targeted, rapid preload reduction needed in this setting; they're not the ideal choice for acutely reducing preload in pulmonary edema. So, the agent best for acutely reducing preload is IV nitroglycerin (or other IV nitrates).*

## 7. Which symptom is most characteristic of left-sided heart failure?

A. Jugular venous distention

B. Persistent cough with crackles in the lungs

C. Peripheral edema

D. Ascites

*Left-sided heart failure leads to a backup of blood into the lungs, causing pulmonary congestion. This fluid overload in the lungs irritates airways and impairs gas exchange, producing a persistent cough and crackles heard on auscultation. These lung findings are the hallmark of left-sided failure because the problem starts in the heart's left side and shows up primarily in the lungs. Jugular venous distention, peripheral edema, and ascites are more commonly associated with right-sided failure or advanced biventricular failure, reflecting systemic venous congestion rather than pulmonary edema. Therefore, the cough with lung crackles best fits the left-sided pattern.*

## 8. A patient taking Digoxin has severe bradycardia, nausea, and vomiting, with a Digoxin level of 4 ng/mL. What medication would the physician order?

A. Narcan

B. Aminophylline

C. Digibind

D. No medication because this is a normal level.

*Digoxin toxicity is suspected when there are prominent GI symptoms like nausea and vomiting along with bradycardia, and a serum digoxin level well above the therapeutic range. The treatment of choice in this situation is digoxin-specific antibody fragments (Digibind). These antibody fragments bind free digoxin in the bloodstream, forming inactive complexes that are cleared by the kidneys. By neutralizing digoxin, they rapidly reverse the drug's effects on the heart and other tissues, which is crucial when severe bradycardia and toxicity symptoms are present. Knowing what each option would do helps connect the concept: Narcan reverses opioids, not digoxin. Aminophylline has limited or historical use in certain arrhythmias but is not the standard antidote for digoxin toxicity. Not giving any medication would leave the toxicity unmanaged despite the high risk. The key idea is that the antidote directly targets and neutralizes digoxin, which is essential when toxicity is evident and the level is elevated.*

SAMPLE

**9. Which of the following actions is most appropriate if a patient on Digoxin reports vision changes?**

A. Ignore as it is unlikely related to Digoxin

B. Advise the patient to rest and retake the dose later

C. Instruct to report immediately to the clinician

D. Resume dosing only after Potassium normalizes

*Digoxin has a very narrow therapeutic window, so signs of toxicity can appear quickly. Vision disturbances, such as blurred vision or seeing yellow or green halos, are a classic warning sign of digoxin toxicity. When a patient on digoxin reports vision changes, the correct course is to have them contact a clinician immediately. Prompt evaluation is essential because the clinician may hold the medication, check the digoxin level, assess kidney function and electrolyte status (especially potassium), and determine the safest next steps. This reduces the risk of progression to more serious toxicity and dangerous heart rhythms. Ignoring the symptom or delaying evaluation is not appropriate, and simply resting or retaking the dose later could allow toxicity to worsen. Resuming dosing after potassium normalizes assumes a safe correction without medical guidance, which is not reliable enough when vision changes are involved.*

**10. In the hospital management of heart failure, inotropes are used to:**

A. Cure HF

B. Improve cardiac output in patients with low output despite optimized therapy or in shock

C. Increase preload

D. Worsen heart failure

*Inotropes are used to improve forward flow by increasing the heart's contractility when cardiac output is too low, such as in acute decompensated heart failure or cardiogenic shock. By boosting contractility, they raise stroke volume and overall cardiac output, which improves perfusion to vital organs and helps stabilize blood pressure. This is temporary support while other treatments address the underlying problem; they are not a cure for heart failure. Inotropes do not increase preload, and if misused they can raise heart rate and oxygen demand or trigger arrhythmias, so they're used with careful monitoring. Common agents like dobutamine or milrinone are chosen based on the patient's hemodynamic needs.*

SAMPLE

# Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nclexheartfailure.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE