NCLEX NEUROLOGIC AND SENSORY SYSTEMS PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nclexneurologicsensorysys.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What role does the somatosensory cortex play in the nervous system?

A. Processing auditory information

B. Regulating emotional responses

C. Processing tactile information

D. Controlling voluntary movement

2. Which symptom might indicate a neurological emergency such as herniation?

A. Persistent headache

B. Slight confusion

C. Sudden change in consciousness or pupillary responses

D. Nausea only

3. Which instructions should the nurse give to a client at risk of corneal damage?

A. Use saline drops.

B. Improve lighting at home.

C. Increase humidity at home.

D. Wear prescribed lens for best vision.

4. Which cranial nerve is responsible for the sense of smell?

A. Cranial nerve II

B. Cranial nerve V

C. Cranial nerve I

D. Cranial nerve VIII

5. How does hydrocephalus affect neurological function?

A. It enhances cognitive abilities

B. It leads to decreased intracranial pressure

C. It can lead to increased intracranial pressure and neurological impairments

D. It only affects motor function

SAMPLE

6. How does a hemorrhagic stroke differ from an ischemic stroke?

A. Hemorrhagic stroke involves a blockage in blood flow

B. Hemorrhagic stroke occurs due to bleeding in the brain

C. Ischemic stroke occurs due to swelling in the brain

D. Ischemic stroke is caused by high blood pressure

7. Which condition is primarily characterized by damage to the myelin sheath?

A. Parkinson's disease

B. Multiple sclerosis

C. Alzheimer's disease

D. Huntington's disease

8. Which cranial nerve primarily controls eye movements?

A. Trigeminal nerve (CN V)

B. Olfactory nerve (CN I)

C. Oculomotor nerve (CN III)

D. Facial nerve (CN VII)

9. Which symptom is associated with a lesion in the occipital lobe?

A. Hearing loss

B. Vision problems

C. Balance issues

D. Memory impairment

10. Which situation would suggest that a patient with increased ICP might benefit from sedation?

A. Patient exhibits combativeness

B. Patient shows decreased respiratory rate

C. Patient remains alert and oriented

D. Patient is unresponsive to verbal stimuli

SAMPLE

Answers

SAMPLE

1. C
2. C
3. A
4. C
5. C
6. B
7. B
8. C
9. B
10. A

SAMPLE

Explanations

SAMPLE

1. What role does the somatosensory cortex play in the nervous system?

A. Processing auditory information

B. Regulating emotional responses

C. Processing tactile information

D. Controlling voluntary movement

The somatosensory cortex is primarily responsible for processing tactile information, which includes sensations such as touch, temperature, pain, and body position. This area of the brain is located in the parietal lobe and receives input from sensory receptors throughout the body. When a stimulus is detected, such as pressure on the skin, the somatosensory cortex interprets this information to allow the individual to perceive sensations and respond accordingly. This processing is crucial for the integration of sensory information, enabling effective interaction with the environment. Other functions mentioned, such as processing auditory information or regulating emotional responses, are managed by separate regions of the brain, such as the temporal lobe for auditory processing and the limbic system for emotional regulation. Additionally, the control of voluntary movement is primarily managed by the motor cortex, which is located in the frontal lobe. Thus, the distinct role of the somatosensory cortex in processing tactile information highlights its critical function in sensory perception within the nervous system.

2. Which symptom might indicate a neurological emergency such as herniation?

A. Persistent headache

B. Slight confusion

C. Sudden change in consciousness or pupillary responses

D. Nausea only

A sudden change in consciousness or pupillary responses is a critical symptom that can indicate a neurological emergency such as herniation. This occurs when brain tissue shifts from its normal position due to increased intracranial pressure (ICP). In cases of herniation, rapid and severe pressure on the brain can lead to alterations in consciousness, as the brain's ability to process information and maintain alertness is compromised. Additionally, changes in pupillary responses, such as dilated or unequal pupils, can reflect increased ICP or brainstem involvement, signaling that immediate medical intervention is necessary. Recognizing these signs is essential for prompt diagnosis and treatment to prevent further neurological damage or deterioration of the patient's condition.

SAMPLE

3. Which instructions should the nurse give to a client at risk of corneal damage?

A. Use saline drops.

B. Improve lighting at home.

C. Increase humidity at home.

D. Wear prescribed lens for best vision.

Using saline drops is an appropriate instruction for a client at risk of corneal damage because these drops help to keep the eyes moist and can provide relief from dryness, which is a common issue that can lead to corneal irritation or damage. Saline drops assist in maintaining the natural moisture of the cornea, thereby preventing further complications that can arise from dryness, such as abrasions or infections. Improving lighting at home may enhance visibility but does not directly address the moisture needs of the cornea. When a client is at risk for corneal damage, maintaining the integrity of the ocular surface is more crucial than simply improving environmental lighting conditions. While increasing humidity can be beneficial in certain contexts, it is not as immediate or targeted in providing relief to the cornea as saline drops. Humidity affects the overall environment but does not provide direct lubrication or hydration to the eyes. Wearing prescribed lenses is important for achieving the best vision, yet it does not necessarily protect against corneal damage. Some lenses may not be suitable for individuals with corneal vulnerabilities, especially if they contribute to dryness or irritation. Thus, while lens wear is essential for vision correction, it does not prioritize the immediate protection and hydration needed for corneal health.

4. Which cranial nerve is responsible for the sense of smell?

A. Cranial nerve II

B. Cranial nerve V

C. Cranial nerve I

D. Cranial nerve VIII

Cranial nerve I, also known as the olfactory nerve, is specifically responsible for the sense of smell. It functions by transmitting sensory information from the olfactory receptors in the nasal cavity to the olfactory bulb, which processes the signals and connects them to higher brain regions involved in smell perception. Cranial nerve II, or the optic nerve, is associated with vision rather than olfactory senses. Cranial nerve V, the trigeminal nerve, primarily carries sensory information from the face and is involved in sensations such as touch and pain rather than smell. Lastly, cranial nerve VIII, known as the vestibulocochlear nerve, is concerned with hearing and balance, making it unrelated to the sense of smell. Thus, the correct answer focuses entirely on the olfactory function provided by cranial nerve I.

SAMPLE

5. How does hydrocephalus affect neurological function?

A. It enhances cognitive abilities

B. It leads to decreased intracranial pressure

C. It can lead to increased intracranial pressure and neurological impairments

D. It only affects motor function

Hydrocephalus is a condition characterized by an accumulation of cerebrospinal fluid (CSF) in the ventricles of the brain, which can significantly affect neurological function. This condition often leads to increased intracranial pressure, as the excessive fluid creates pressure within the skull. The elevated pressure can disrupt normal brain function and lead to various neurological impairments. Individuals with hydrocephalus may experience symptoms such as cognitive deficits, changes in personality, difficulties with balance and coordination, and problems with vision. The pressure can also interfere with normal blood flow and the functioning of brain cells, compounding the risk of neurological impairment. Management of hydrocephalus typically involves addressing the increased intracranial pressure, often through the surgical placement of a shunt to drain excess fluid. While hydrocephalus can affect motor functions, it is not limited to just motor function disturbances; it can have a broad impact on cognitive abilities and other neurological functions as well. Therefore, recognizing that increased intracranial pressure can lead to a variety of neurological issues is essential for understanding the implications of hydrocephalus on overall brain health.

6. How does a hemorrhagic stroke differ from an ischemic stroke?

A. Hemorrhagic stroke involves a blockage in blood flow

B. Hemorrhagic stroke occurs due to bleeding in the brain

C. Ischemic stroke occurs due to swelling in the brain

D. Ischemic stroke is caused by high blood pressure

A hemorrhagic stroke is characterized by bleeding in the brain, which arises when a blood vessel ruptures, causing blood to spill into the surrounding brain tissue. This can lead to increased intracranial pressure and damage to the brain cells, as the blood can irritate the tissues and disrupt normal brain function. Consequently, the primary distinction between hemorrhagic and ischemic strokes lies in the nature of the vascular event; hemorrhagic strokes result from a rupture, while ischemic strokes result from blockage of blood supply due to a clot or narrowing of the arteries. This distinction is crucial for determining the appropriate treatment plan. For example, hemorrhagic strokes may require different interventions compared to ischemic strokes, which are often treated with medications aimed at dissolving the clot or restoring blood flow. Understanding the mechanisms behind these two types of strokes is essential for both prevention and management of stroke patients.

SAMPLE

7. Which condition is primarily characterized by damage to the myelin sheath?

A. Parkinson's disease

B. Multiple sclerosis

C. Alzheimer's disease

D. Huntington's disease

Multiple sclerosis is primarily characterized by damage to the myelin sheath, which is essential for the proper functioning of the nervous system. The myelin sheath is a protective covering that surrounds nerve fibers, allowing for the efficient transmission of electrical impulses between nerve cells. In multiple sclerosis, the immune system mistakenly attacks and destroys this myelin sheath, leading to a range of neurological symptoms. The damage disrupts communication between the brain and other parts of the body, resulting in symptoms such as muscle weakness, coordination issues, visual disturbances, and cognitive changes. The presence of lesions in the central nervous system due to the immune-mediated damage is a hallmark of multiple sclerosis, making it distinct among the conditions listed. Understanding the role of myelin in neurological function highlights why multiple sclerosis is uniquely associated with myelin impairment, distinguishing it from other neurological disorders such as Parkinson's disease, Alzheimer's disease, and Huntington's disease, which have different underlying pathologies.

8. Which cranial nerve primarily controls eye movements?

A. Trigeminal nerve (CN V)

B. Olfactory nerve (CN I)

C. Oculomotor nerve (CN III)

D. Facial nerve (CN VII)

The cranial nerve that primarily controls eye movements is the oculomotor nerve, which is the third cranial nerve. This nerve innervates several extraocular muscles that are responsible for most of the eye's movements, including the superior, medial, and inferior rectus muscles, as well as the inferior oblique muscle. Additionally, the oculomotor nerve provides parasympathetic innervation for pupil constriction and lens accommodation, which are crucial for focusing on nearby objects. For a deeper understanding, the other cranial nerves listed do not primarily govern eye movements. The trigeminal nerve is responsible for sensation in the face and motor functions such as biting and chewing, while the olfactory nerve is involved in the sense of smell. The facial nerve primarily controls facial expressions, taste sensations from the anterior two-thirds of the tongue, and other functions related to facial muscles. Therefore, the unique role of the oculomotor nerve in controlling eye movements distinctly sets it apart from the functions of the other cranial nerves mentioned.

SAMPLE

9. Which symptom is associated with a lesion in the occipital lobe?

A. Hearing loss

B. Vision problems

C. Balance issues

D. Memory impairment

Vision problems are associated with a lesion in the occipital lobe because the occipital lobe is primarily responsible for processing visual information. This part of the brain receives and interprets signals sent from the eyes, allowing us to perceive shapes, colors, motion, and depth. When a lesion occurs in this area, it can lead to a variety of visual disturbances, such as visual field deficits, difficulties in recognizing familiar faces, or experiencing visual hallucinations. Other symptoms related to hearing loss, balance issues, or memory impairment are not typically linked to the occipital lobe. Hearing loss is more directly associated with lesions in auditory pathways or structures in the temporal lobe. Balance issues often relate to problems in the cerebellum or other components of the vestibular system. Memory impairment is typically linked to the temporal lobe and structures like the hippocampus, which are crucial for memory formation and retrieval. Thus, the correct identification of vision problems reflects the essential role of the occipital lobe in visual processing.

10. Which situation would suggest that a patient with increased ICP might benefit from sedation?

A. Patient exhibits combativeness

B. Patient shows decreased respiratory rate

C. Patient remains alert and oriented

D. Patient is unresponsive to verbal stimuli

Increased intracranial pressure (ICP) can lead to various neurological complications, and managing a patient's condition effectively is crucial. Sedation may be indicated in situations where the patient's behavior puts them at risk or complicates their care. A patient exhibiting combativeness can be challenging to manage, as their agitation can lead to increased ICP due to factors such as increased blood flow to the brain caused by elevated heart rate and blood pressure during episodes of agitation. Additionally, combativeness may hinder the ability to perform necessary assessments, interventions, or therapeutic procedures. Sedation in this scenario helps create a more stable environment, reduces metabolic demands on the brain, lowers stress response, and can facilitate more effective management of the patient's condition. The other scenarios mentioned either reflect a stable neurological status or do not indicate an acute response that would necessitate sedation for the patient's safety and care. Therefore, addressing combativeness through sedation is a logical and beneficial approach for a patient with increased ICP.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nclexneurologicsensorysys.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE