NCLEX OB POSTPARTUM PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nclexobpostpartum.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. Which intervention is used if pharmacologic measures fail to control postpartum hemorrhage due to uterine atony?

A. Immediate hysterectomy.

B. Balloon tamponade.

C. Observation only.

D. Antibiotics.

2. Which factor is NOT associated with increased afterpains?

A. Bottle-feeding

B. Diabetes

C. Multiple gestation

D. Primiparity

3. Which symptom might indicate a thromboembolic event postpartum?

A. Severe chest pain.

B. Mild headaches.

C. Clear breath sounds and no pain.

D. No cough.

4. Which statement best describes the taking-in stage after delivery?

A. The mother immediately begins full involvement in infant care.

B. The mother is uninterested in the infant and asks others to respond to cues.

C. The mother focuses on her own needs and rest.

D. The mother experiences postpartum depression.

5. What is the key nursing action to prevent uterine atony-related hemorrhage after delivery?

A. Administer uterotonic medications as prescribed and perform fundal massage as needed.

B. Elevate legs and restrict fluids to prevent edema.

C. Apply cold packs to the abdomen to reduce cramping.

D. Delay breastfeeding to prevent stimulation of the uterus.

SAMPLE

6. Which statement best reflects taking-in stage behavior in the postpartum period?

A. The mother is preoccupied with her own needs and rest.

B. The mother demonstrates strong early bonding by initiating infant care.

C. The mother experiences postpartum psychosis.

D. The mother refuses to hold infant.

7. Which hormone stimulates milk production in the postpartum period?

A. Oxytocin stimulates milk production.

B. Prolactin stimulates milk production.

C. Estrogen stimulates milk production.

D. Progesterone stimulates milk production.

8. Which postpartum condition commonly requires antibiotic therapy and infection control precautions?

A. Postpartum depression.

B. Uterine atony.

C. Endometritis.

D. Physiologic jaundice.

9. Within the BUBBLE-HE postpartum assessment framework, what does the Emotions/Homans sign component assess?

A. Assessing emotional status and screening for deep vein thrombosis

B. Assessing appetite and hormonal changes

C. Assessing urinary continence

D. Assessing wound approximation

10. Which finding would indicate hypovolemia in a postpartum client?

A. Warm, dry skin

B. Decreased thirst

C. Rapid, weak pulse

D. Normal blood pressure

SAMPLE

Answers

SAMPLE

1. B
2. A
3. A
4. C
5. A
6. A
7. B
8. C
9. A
10. C

SAMPLE

Explanations

SAMPLE

1. Which intervention is used if pharmacologic measures fail to control postpartum hemorrhage due to uterine atony?

A. Immediate hysterectomy.

B. Balloon tamponade.

C. Observation only.

D. Antibiotics.

When postpartum hemorrhage is caused by uterine atony, the goal is to rapidly stop the bleeding by helping the uterus contract and by applying direct pressure to bleeding sites. If medications that contract the uterus don't work, inserting a balloon tamponade device (a balloon inserted into the uterus and then inflated with saline) provides internal pressure that compresses bleeding vessels and helps control the hemorrhage. This approach is less invasive than surgery and can stabilize the patient while arranging definitive therapy, such as uterine artery embolization if needed or, in extreme cases, hysterectomy. Observation would not address the active bleeding, and antibiotics don't stop hemorrhage. Immediate hysterectomy is more definitive but is avoided when tamponade can control the bleed and preserve fertility.

2. Which factor is NOT associated with increased afterpains?

A. Bottle-feeding

B. Diabetes

C. Multiple gestation

D. Primiparity

Afterpains are cramps caused by the uterus contracting as it involutes back to its pre-pregnancy size. The strength of these contractions is driven by oxytocin. Breastfeeding stimulates oxytocin release, which increases uterine contractions and, therefore, afterpains. Bottle-feeding avoids nipple stimulation, so oxytocin release is reduced and afterpains are not increased—often they're milder. The other factors listed are associated with more intense involution and stronger contractions (for example, a uterus that has been more stretched in multiple gestation), so they tend to be linked with increased afterpains. Diabetes isn't a primary driver of increased afterpains.

3. Which symptom might indicate a thromboembolic event postpartum?

A. Severe chest pain.

B. Mild headaches.

C. Clear breath sounds and no pain.

D. No cough.

Postpartum is a time when the blood is more likely to clot, due to a combination of a hypercoagulable state, venous stasis, and vascular changes from delivery. A symptom that strongly suggests a thromboembolic event is severe chest pain, especially if it comes with shortness of breath, rapid heartbeat, or sweating. This pattern points to a potential pulmonary embolism, where a clot travels to the lungs and disrupts circulation. Mild headaches are not typical signs of a clot and could reflect other postpartum issues like tension or hormonal changes. Clear breath sounds with no pain and no cough do not point toward a thromboembolic event; they suggest the lungs and airways aren't showing signs of PE at that moment. So the presence of severe chest pain in the postpartum period is the red flag that warrants urgent evaluation for a thromboembolism.

SAMPLE

4. Which statement best describes the taking-in stage after delivery?

A. The mother immediately begins full involvement in infant care.

B. The mother is uninterested in the infant and asks others to respond to cues.

C. The mother focuses on her own needs and rest.

D. The mother experiences postpartum depression.

The taking-in stage after birth centers on recovery and meeting the mother's own needs. During this time she often feels exhausted and focused on rest, fluids, and nourishment, relying on others to help with basic care. She may need time to process the birth and is less focused on actively caring for the newborn yet. This is why describing the mother as concentrating on her own needs and rest fits this stage best. In contrast, immediate full involvement in infant care aligns with a later stage where the mother begins to take a more active role, and statements about being uninterested or experiencing postpartum depression describe other conditions or phases.

5. What is the key nursing action to prevent uterine atony-related hemorrhage after delivery?

A. Administer uterotonic medications as prescribed and perform fundal massage as needed.

B. Elevate legs and restrict fluids to prevent edema.

C. Apply cold packs to the abdomen to reduce cramping.

D. Delay breastfeeding to prevent stimulation of the uterus.

Maintaining uterine tone to prevent postpartum hemorrhage from uterine atony is the key. The best nursing action is to administer uterotonic medications as prescribed and perform fundal massage as needed. Uterotonic drugs stimulate the uterus to contract, which compresses the blood vessels and reduces bleeding. Fundal massage physically prompts the uterus to firm up, helping to regain Tone and control hemorrhage if the uterus feels boggy. After delivery, the uterus should be midline and firm. If it's soft or boggy, gently massage the uterus to encourage contraction and re-check its firmness. Along with medications, ensuring the bladder is empty helps the uterus stay in the correct position and contract effectively. Monitor vital signs, lochia, and fundal height regularly, and watch for signs of excessive bleeding or adverse medication effects. Why the other actions aren't appropriate for preventing atony-related bleeding: leg elevation with fluid restriction does not address uterine contraction or bleeding control; applying cold to the abdomen targets cramping rather than uterine tone; delaying breastfeeding removes a natural stimulus that promotes uterine contraction and can worsen atony.

6. Which statement best reflects taking-in stage behavior in the postpartum period?

A. The mother is preoccupied with her own needs and rest.

B. The mother demonstrates strong early bonding by initiating infant care.

C. The mother experiences postpartum psychosis.

D. The mother refuses to hold infant.

Taking-in is the first phase of maternal adjustment after birth, typically lasting about the first 24 to 48 hours. During this time the mother is often dependent and focused on her own recovery—rest, nutrition, comfort, and physical recovery from labor. She may speak about the birth experience and rely on others to care for the newborn while she recuperates. Because of this emphasis on self-care and fatigue, she is less likely to initiate infant care. This makes the statement about being preoccupied with her own needs and rest the best reflection of taking-in behavior. Bonding and active caregiving tend to emerge in the next phase (taking-hold), whereas postpartum psychosis or outright refusal to hold the infant would indicate pathology or a non-normative response rather than a normal stage.

SAMPLE

7. Which hormone stimulates milk production in the postpartum period?

A. Oxytocin stimulates milk production.

B. Prolactin stimulates milk production.

C. Estrogen stimulates milk production.

D. Progesterone stimulates milk production.

Milk production after birth is driven by prolactin, a hormone from the anterior pituitary. After delivery, the placenta is gone, and estrogen and progesterone levels fall, removing their inhibitory effect on the mammary glands. When the baby suckles, nerve signals to the brain reduce dopamine, leading to increased prolactin release and synthesis of milk in the alveolar cells. Oxytocin, while essential for the let down reflex to eject milk, does not stimulate its production. Estrogen and progesterone support breast tissue during pregnancy but don't drive milk production postpartum. Thus, prolactin is the hormone that stimulates milk production in the postpartum period.

8. Which postpartum condition commonly requires antibiotic therapy and infection control precautions?

A. Postpartum depression.

B. Uterine atony.

C. Endometritis.

D. Physiologic jaundice.

Endometritis is an infection of the uterine lining that can develop after delivery, especially after a cesarean birth or with prolonged rupture of membranes. It commonly requires antibiotic therapy to eradicate the infection and prevent progression to sepsis. In the hospital setting, infection control precautions are used to reduce transmission risk, emphasizing strict hand hygiene and standard precautions, with additional measures as indicated by the patient's status. The other postpartum conditions listed do not involve infection requiring antibiotics: postpartum depression is a mood disorder treated with support and therapy (not antibiotics), uterine atony causes hemorrhage and is managed with uterotonics and massage rather than antibiotics, and physiologic jaundice is a benign, noninfectious condition of newborn bilirubin metabolism.

9. Within the BUBBLE-HE postpartum assessment framework, what does the Emotions/Homans sign component assess?

A. Assessing emotional status and screening for deep vein thrombosis

B. Assessing appetite and hormonal changes

C. Assessing urinary continence

D. Assessing wound approximation

The Emotions/Homans sign part of the BUBBLE-HE postpartum assessment covers two areas: how the mother is emotionally and psychologically after birth, and whether there are signs suggesting a risk for deep vein thrombosis. Emotionally, you're looking at mood, affect, coping, bonding with the newborn, and any symptoms that might indicate postpartum mood changes or depression. This helps ensure the mother's mental health is supported during a time when mood fluctuations are common but when more serious issues can develop. The Homans sign portion screens for DVT risk by assessing calf pain or tenderness with dorsiflexion, along with signs like swelling or warmth in the legs. Postpartum women have an increased risk of clots due to a hypercoagulable state, so including this screen in the emotional/homans pairing makes sense. Other options don't fit this component because appetite and hormonal changes relate to endocrine or general physiological changes, urinary continence falls under the bladder assessment, and wound approximation relates to the episiotomy/incision area.

SAMPLE

10. Which finding would indicate hypovolemia in a postpartum client?

A. Warm, dry skin

B. Decreased thirst

C. Rapid, weak pulse

D. Normal blood pressure

Hypovolemia means there is less blood effectively circulating, so the heart tries to compensate by beating faster, but the force of each beat diminishes. A rapid, weak pulse is the clearest reflection of this reduced volume and decreased preload; it often appears early in postpartum hemorrhage as the body attempts to maintain perfusion to vital organs. In contrast, skin that is warm and dry doesn't fit hypovolemia caused by blood loss—the typical response is cool, clammy skin from peripheral vasoconstriction. Decreased thirst isn't a reliable early sign of reduced circulating volume, and normal blood pressure can occur with early compensation, so it doesn't signal hypovolemia by itself.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nclexobpostpartum.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE