


NEURO MUSCLE TONE PRACTICE TEST (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR
THE FULL VERSION STUDY GUIDE

<https://neuromuscletone.examzify.com>



Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright	1
Table of Contents	2
Introduction	3
How to Use This Guide	4
Questions	5
Answers	8
Explanations	10
Next Steps	15

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

Questions

SAMPLE

1. What is the purpose of dynamic ROM assessment in spasticity?

- A. To measure the effect of passive movement latency on rehabilitation planning.
- B. To identify contracture severity independent of neural input.
- C. To assess static range of motion only.
- D. To separate neural resistance from fixed contractures and quantify the dynamic component of tone for targeted treatment planning.

2. Why do you see abnormal limb posturing with dyscontrol?

- A. Gravitational effects
- B. Loss of mobility
- C. Connective tissue changes
- D. Abnormal response to stretch

3. Rigidity is often associated with which type of disorders?

- A. Peripheral neuropathies
- B. Cerebellar diseases
- C. Spinal cord injuries
- D. Basal ganglia disorders

4. Which statement describes a mechanism at the level of the spinal cord contributing to clonus?

- A. It is a mechanism at the level of the spinal cord contributing to clonus
- B. It is not involved in clonus
- C. Only peripheral nerves contribute to clonus
- D. Clonus is not related to central mechanisms

5. What is a common post-stroke pattern difference between upper and lower limbs?

- A. Upper limb flexor-dominant spasticity; lower limb plantarflexors and knee extensors/flexors.
- B. Upper limb plantarflexors; lower limb elbow flexors.
- C. Both limbs show identical patterns.
- D. No difference between limbs.

6. Which statement about spasticity is true?

- A. It is velocity-dependent.
- B. It is not velocity-dependent.
- C. It only occurs during active movement.
- D. It is unrelated to reflexes.

7. Which term describes the normal plantar response?

- A. Toe spreading
- B. Inversion of the foot
- C. Great toe extension
- D. Great toe flexion

8. Which of the following is NOT typically associated with an upper motor neuron lesion?

- A. Babinski sign present
- B. Hyperreflexia
- C. Muscle atrophy due to denervation
- D. Spasticity

9. What are important observation pieces in examination of spasticity?

- A. Sustained posturing
- B. Associated reactions
- C. Gait pattern
- D. Muscle strength

10. Describe the role of dystonia in motor disorders associated with tone.

- A. Dystonia reduces tone.
- B. Dystonia contributes abnormal, often patterned muscle contractions and postures that can exacerbate functional impairment and interact with spasticity.
- C. Dystonia only affects speech.
- D. Dystonia is unrelated to tone.

Answers

SAMPLE

1. D
2. D
3. D
4. A
5. A
6. A
7. D
8. C
9. B
10. B

SAMPLE

Explanations

SAMPLE

1. What is the purpose of dynamic ROM assessment in spasticity?

- A. To measure the effect of passive movement latency on rehabilitation planning.
- B. To identify contracture severity independent of neural input.
- C. To assess static range of motion only.
- D. To separate neural resistance from fixed contractures and quantify the dynamic component of tone for targeted treatment planning.**

The main idea is to separate how much of the movement resistance comes from neural factors that change with speed from the fixed, non-neural restrictions of the muscle and joint. In spasticity, tone increases when you move faster because the stretch reflex is more excitable. By testing range of motion at different speeds, you can see how much resistance is due to this dynamic, neural component versus a hard, fixed contracture. This distinction lets you tailor treatment to what's driving the problem—neural-focused options (like medications or muscle-relaxing injections) when the dynamic neural component is large, or approaches aimed at correcting fixed shortening when the non-neural portion dominates. It's not just about static ROM or about movement latency; it's about how resistance changes with speed and what that implies for targeted therapy.

2. Why do you see abnormal limb posturing with dyscontrol?

- A. Gravitational effects
- B. Loss of mobility
- C. Connective tissue changes
- D. Abnormal response to stretch**

Abnormal limb posturing with dyscontrol comes from an abnormal response to stretch caused by hyperexcitable stretch reflexes after damage to motor pathways. When the nervous system loses proper inhibition from higher centers, the muscle spindle reflexes become exaggerated, producing involuntary, sustained contractions that push the limb into unusual positions. This reflex-driven pattern explains why limbs assume abnormal postures as part of dyscontrol, and why the resistance to passive movement can be velocity dependent. Gravitational effects would shift posture due to body position but don't reflect an exaggerated reflex response. Loss of mobility describes stiffness from reduced movement, not the active, reflex-mediated posturing. Connective tissue changes lead to structural contractures over time, which are positional rather than reflex-driven.

3. Rigidity is often associated with which type of disorders?

- A. Peripheral neuropathies
- B. Cerebellar diseases
- C. Spinal cord injuries
- D. Basal ganglia disorders**

Rigidity arises from dysfunction in the basal ganglia circuits that regulate movement, producing a constant, baseline increase in muscle tone. In basal ganglia disorders, such as Parkinsonian syndromes, the loss of dopaminergic input disrupts the balance of motor pathways, leading to resistance to passive movement that is not dependent on movement speed—often described as lead-pipe rigidity, sometimes with a cogwheel component when tremor is present. This contrasts with spasticity, which is velocity-dependent and typically results from upper motor neuron/corticospinal tract injuries. Peripheral neuropathies mainly cause weakness and reduced reflexes, not sustained increased tone, and cerebellar diseases usually present with hypotonia and coordination problems rather than rigidity. Therefore, rigidity is most characteristic of basal ganglia disorders.

4. Which statement describes a mechanism at the level of the spinal cord contributing to clonus?

- A. It is a mechanism at the level of the spinal cord contributing to clonus**
- B. It is not involved in clonus
- C. Only peripheral nerves contribute to clonus
- D. Clonus is not related to central mechanisms

Clonus arises from spinal cord circuitry that becomes disinhibited when descending control from the brain is reduced, such as after an upper motor neuron injury. In the spinal cord, the muscle spindle's Ia afferents repeatedly excite alpha motor neurons, and without enough supraspinal inhibition, this reflex loop can enter a self-sustaining, rhythmic cycle. Each stretch activates the loop, which then drives further contractions, producing the characteristic clonus. This is a spinal mechanism—the same spinal reflex networks are generating the rhythmic activity—rather than being solely due to peripheral nerves or purely cortical processes.

5. What is a common post-stroke pattern difference between upper and lower limbs?

- A. Upper limb flexor-dominant spasticity; lower limb plantarflexors and knee extensors/flexors.**
- B. Upper limb plantarflexors; lower limb elbow flexors.
- C. Both limbs show identical patterns.
- D. No difference between limbs.

After a stroke, motor impairment often settles into limb-specific spasticity patterns. The upper limb tends to develop a flexor-dominant pattern, meaning the elbow, forearm, and wrist/finger flexors become more spastic and pull the limb into a flexed position. In contrast, the lower limb commonly shows an extensor-dominant pattern, with plantarflexors and knee extensors (and associated hip extensors/adductors) contributing to a stiff, extended posture and a tendency toward pushing the foot downward. So the described pattern—upper limb flexor-dominant spasticity and lower limb plantarflexors with knee extensors—best reflects these typical post-stroke differences. The other options describe opposite, identical, or no differences between limbs, which don't fit the well-established pattern.

6. Which statement about spasticity is true?

- A. It is velocity-dependent.**
- B. It is not velocity-dependent.
- C. It only occurs during active movement.
- D. It is unrelated to reflexes.

Spasticity is a velocity-dependent increase in resistance to passive stretch caused by an overactive stretch reflex after upper motor neuron injury. Because the reflex response grows with the speed of the stretch, moving a limb quickly elicits a stronger resistance than moving it slowly. This speed-related change in tone is what sets spasticity apart from other forms of increased muscle tone, like rigidity, where resistance does not depend on how fast you move. Spasticity can be observed during passive movement, not only during active effort, and it stems from hyperexcitable reflex pathways due to loss of supraspinal inhibition on the spinal cord. So claims that spasticity isn't velocity-dependent, or that it occurs only with active movement, or that it's unrelated to reflexes, don't fit the characteristic pattern of velocity-driven stretch reflexes in upper motor neuron syndromes.

7. Which term describes the normal plantar response?

- A. Toe spreading
- B. Inversion of the foot
- C. Great toe extension
- D. Great toe flexion**

When the sole is stimulated, the normal plantar response is the toes curling downward, especially the toes flexing toward the bottom of the foot. This is known as great toe flexion, with the other toes typically following in a curling motion. It reflects intact corticospinal control and a typical, healthy reflex. In contrast, an upward extension of the big toe with spreading of the other toes (the Babinski sign) is an abnormal response seen with corticospinal tract dysfunction. Toe spreading in response to plantar stimulation is essentially that Babinski pattern, not the normal reflex. Turning the foot inward (inversion) or extending the big toe upward are not the standard toe curling response described here. So the normal plantar response is great toe flexion—the curling of the toes toward the sole.

8. Which of the following is NOT typically associated with an upper motor neuron lesion?

- A. Babinski sign present
- B. Hyperreflexia
- C. Muscle atrophy due to denervation**
- D. Spasticity

Upper motor neuron lesions produce an upper motor neuron syndrome, where the brain's inhibitory control over reflex pathways is lost. This leads to weakness with increased muscle tone (spasticity), brisk reflexes (hyperreflexia), and an abnormal Babinski sign. The muscle changes you'd expect come from disuse or other factors, not from loss of motor neurons themselves. Denervation atrophy, with true muscle fiber loss and fasciculations, is a hallmark of lower motor neuron lesions where the motor units are actually damaged. So the feature described as muscle atrophy due to denervation isn't typical of an upper motor neuron lesion, making it the best choice.

9. What are important observation pieces in examination of spasticity?

- A. Sustained posturing
- B. Associated reactions**
- C. Gait pattern
- D. Muscle strength

Inspiring signs of spasticity show up when you watch how movement in one part of the body affects others and how the muscles respond during tasks. The most informative observation is associated reactions—involuntary movements that appear in a limb when you perform a movement with another limb or apply a maneuver. This phenomenon reflects the abnormal coupling and reflex activation that come with upper motor neuron involvement and helps reveal how spasticity influences motor control beyond simple resistance. Sustained posturing can be seen with muscle contractures or dystonia and isn't specific to spasticity itself. A gait pattern shows how spasticity and other impairments affect walking, but gait is a broader, functional outcome rather than a direct observation of spasticity. Muscle strength measures the ability to generate force and can be reduced for many reasons, not only due to spasticity, so it doesn't isolate the spastic component as clearly. So, observing associated reactions provides the clearest window into how spasticity manifests during movement.

10. Describe the role of dystonia in motor disorders associated with tone.

- A. Dystonia reduces tone.
- B. Dystonia contributes abnormal, often patterned muscle contractions and postures that can exacerbate functional impairment and interact with spasticity.**
- C. Dystonia only affects speech.
- D. Dystonia is unrelated to tone.

Dystonia in motor disorders with abnormal tone shows up as sustained or intermittent muscle contractions that create twisting movements or abnormal postures. This means it changes how muscles are activated, producing patterned, often involuntary contractions, rather than simply changing the overall level of tone. In practice, dystonia can occur alongside spasticity and other tone abnormalities, and during dystonic activity the muscles may co-contract or become markedly hypertonic in specific postures. This can worsen functional ability because the abnormal postures and movements disrupt smooth, voluntary tasks, and the effects are often task- or position-specific and triggered by effort, fatigue, or stress. Dystonia can affect limbs, the trunk, or face and is not limited to speech, nor is it just a reduction of tone.

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

<https://neuromuscletone.examzify.com>

We wish you the very best on your exam journey. You've got this!

SAMPLE