NEW SOUTH WALES AMBULANCE PHARMACOLOGY PRACTICE EXAM (SAMPLE)

STUDY GUIDE



SAMPLE STUDY GUIDE WITH LIMITED QUESTIONS

VISIT US ONLINE FOR THE FULL VERSION STUDY GUIDE

https://nswambpharmacology.examzify.com

Copyright © 2026 by Examzify - A Kaluba Technologies Inc. product.

ALL RIGHTS RESERVED.

No part of this book may be reproduced or transferred in any form or by any means, graphic, electronic, or mechanical, including photocopying, recording, web distribution, taping, or by any information storage retrieval system, without the written permission of the author.

Notice: Examzify makes every reasonable effort to obtain accurate, complete, and timely information about this product from reliable sources.

SAMPLE

Table of Contents

Copyright 1

Table of Contents 2

Introduction 3

How to Use This Guide 4

Questions 5

Answers 8

Explanations 10

Next Steps 16

SAMPLE

Introduction

Preparing for a certification exam can feel overwhelming, but with the right tools, it becomes an opportunity to build confidence, sharpen your skills, and move one step closer to your goals. At Examzify, we believe that effective exam preparation isn't just about memorization, it's about understanding the material, identifying knowledge gaps, and building the test-taking strategies that lead to success.

This guide was designed to help you do exactly that.

Whether you're preparing for a licensing exam, professional certification, or entry-level qualification, this book offers structured practice to reinforce key concepts. You'll find a wide range of multiple-choice questions, each followed by clear explanations to help you understand not just the right answer, but why it's correct.

The content in this guide is based on real-world exam objectives and aligned with the types of questions and topics commonly found on official tests. It's ideal for learners who want to:

- Practice answering questions under realistic conditions,
- Improve accuracy and speed,
- Review explanations to strengthen weak areas, and
- Approach the exam with greater confidence.

We recommend using this book not as a stand-alone study tool, but alongside other resources like flashcards, textbooks, or hands-on training. For best results, we recommend working through each question, reflecting on the explanation provided, and revisiting the topics that challenge you most.

Remember: successful test preparation isn't about getting every question right the first time, it's about learning from your mistakes and improving over time. Stay focused, trust the process, and know that every page you turn brings you closer to success.

Let's begin.

SAMPLE

How to Use This Guide

This guide is designed to help you study more effectively and approach your exam with confidence. Whether you're reviewing for the first time or doing a final refresh, here's how to get the most out of your Examzify study guide:

1. Start with a Diagnostic Review

Skim through the questions to get a sense of what you know and what you need to focus on. Your goal is to identify knowledge gaps early.

2. Study in Short, Focused Sessions

Break your study time into manageable blocks (e.g. 30 – 45 minutes). Review a handful of questions, reflect on the explanations.

3. Learn from the Explanations

After answering a question, always read the explanation, even if you got it right. It reinforces key points, corrects misunderstandings, and teaches subtle distinctions between similar answers.

4. Track Your Progress

Use bookmarks or notes (if reading digitally) to mark difficult questions. Revisit these regularly and track improvements over time.

5. Simulate the Real Exam

Once you're comfortable, try taking a full set of questions without pausing. Set a timer and simulate test-day conditions to build confidence and time management skills.

6. Repeat and Review

Don't just study once, repetition builds retention. Re-attempt questions after a few days and revisit explanations to reinforce learning. Pair this guide with other Examzify tools like flashcards, and digital practice tests to strengthen your preparation across formats.

There's no single right way to study, but consistent, thoughtful effort always wins. Use this guide flexibly, adapt the tips above to fit your pace and learning style. You've got this!

SAMPLE

Questions

SAMPLE

1. What is one of the primary actions of Compound Sodium Lactate?

A. Inhibits blood clotting

B. Distributes throughout the extracellular fluid space

C. Acts as a vasoconstrictor

D. Increases blood pH levels

2. Which antidepressant class does Sodium Bicarbonate interact with by altering protein binding?

A. SSRIs

B. MAO inhibitors

C. Tricyclic antidepressants

D. Mixed serotonin-norepinephrine reuptake inhibitors

3. Which of the following is a primary action of both Glucose 10% and Glucose Gel?

A. Acts as an anti-inflammatory agent

B. Principal energy source for body cells, especially the brain

C. Reduces fever

D. Promotes fat metabolism

4. What can occur at high doses of Atropine?

A. Diarrhea

B. Seizures

C. Hypoglycemia

D. Depression

5. What type of analgesic is methoxyflurane classified as?

A. Inhaled analgesic

B. Topical analgesic

C. Injectable analgesic

D. Oral analgesic

SAMPLE

6. Which type of medications is oxygen commonly used as a drive gas to administer?

A. Oral medications

B. Intravenous medications

C. Nebulised medications

D. Sublingual medications

7. What effect does Naloxone primarily reverse in patients?

A. Increased heart rate

B. Respiratory depression

C. Fever and chills

D. Muscle spasms

8. What is a significant risk associated with oxygen use?

A. Increased blood pressure

B. Impaired oxygenation

C. Increased fire and explosion risk

D. Infection risk

9. What is the primary action of Frusemide?

A. Inhibition of renal sodium reabsorption

B. Promotion of potassium retention

C. Decrease in urine output

D. Stimulation of liver glycogen conversion

10. What is a reported adverse effect of paracetamol?

A. Asthma attacks

B. Nausea

C. Rash

D. Heart palpitations

SAMPLE

Answers

SAMPLE

1. B
2. C
3. B
4. B
5. A
6. C
7. B
8. C
9. A
10. B

SAMPLE

Explanations

SAMPLE

1. What is one of the primary actions of Compound Sodium Lactate?

A. Inhibits blood clotting

B. Distributes throughout the extracellular fluid space

C. Acts as a vasoconstrictor

D. Increases blood pH levels

The primary action of Compound Sodium Lactate, also known as Hartmann's solution, involves its distribution throughout the extracellular fluid space. This intravenous fluid is designed to expand the intravascular volume and is particularly effective in situations such as dehydration, surgery, or trauma where fluid resuscitation is necessary. When administered, Compound Sodium Lactate mimics the composition of extracellular fluid, making it effective for replenishing lost fluids. Its isotonic nature allows it to remain in the extracellular space, aiding in hydration and maintaining electrolyte balance. This distribution is crucial in clinical settings as it helps stabilize blood pressure and ensures adequate tissue perfusion. The other options outline actions that do not accurately represent how Compound Sodium Lactate functions. While it may have secondary effects on acid-base balance, its primary role is not as a direct agent to increase blood pH levels or inhibit blood clotting. It is also not inherently a vasoconstrictor, as its primary mechanism involves fluid replacement rather than affecting vascular tone directly.

2. Which antidepressant class does Sodium Bicarbonate interact with by altering protein binding?

A. SSRIs

B. MAO inhibitors

C. Tricyclic antidepressants

D. Mixed serotonin-norepinephrine reuptake inhibitors

Sodium Bicarbonate notably interacts with tricyclic antidepressants (TCAs) by altering their protein binding, which can affect the medication's efficacy and safety profile. TCAs are known to be highly protein-bound, meaning that a considerable portion of the drug that remains bound is inactive and unable to exert its therapeutic effects. When Sodium Bicarbonate is administered, it can lead to changes in the pH and subsequently affect the protein binding sites, potentially increasing the levels of free, unbound TCAs in circulation. This can enhance the therapeutic effects but also raises the risk of side effects and toxicity, such as cardiovascular complications or central nervous system effects. The other classes of antidepressants, while they may have various interactions with other substances, do not generally have the same significant protein binding alterations when combined with Sodium Bicarbonate. SSRIs, MAO inhibitors, and mixed serotonin-norepinephrine reuptake inhibitors exhibit different pharmacokinetic profiles that are not as influenced by changes in protein binding due to Sodium Bicarbonate as TCAs are. Therefore, understanding the specific interaction of Sodium Bicarbonate with tricyclic antidepressants is crucial for safe and effective pharmacological management of patients requiring these medications.

SAMPLE

3. Which of the following is a primary action of both Glucose 10% and Glucose Gel?

A. Acts as an anti-inflammatory agent

B. Principal energy source for body cells, especially the brain

C. Reduces fever

D. Promotes fat metabolism

The correct answer is that both Glucose 10% and Glucose Gel serve as a principal energy source for body cells, especially the brain. Glucose is a simple sugar that is readily utilized by the body for energy. It is particularly crucial for the brain, which relies heavily on glucose as its primary fuel source due to the organ's high metabolic demand and lack of energy reserves. Administering glucose in either of these forms quickly elevates blood sugar levels, providing immediate energy to cells throughout the body. This is especially important during situations of hypoglycemia, where there is an abnormally low level of glucose in the blood, resulting in symptoms like confusion, weakness, or even loss of consciousness. In contrast, the other options do not align with the primary actions of glucose. Antiinflammatory actions do not pertain to glucose, as its primary role is energy provision rather than modulating inflammation. Similarly, glucose does not have antipyretic (fever-reducing) properties, nor does it play a role in promoting fat metabolism directly; it primarily provides glucose for immediate energy use rather than influencing lipid metabolism significantly.

4. What can occur at high doses of Atropine?

A. Diarrhea

B. Seizures

C. Hypoglycemia

D. Depression

At high doses of Atropine, the central nervous system can be significantly affected, leading to the risk of seizures. Atropine, an anticholinergic drug, primarily works by blocking the action of acetylcholine on muscarinic receptors, resulting in various physiological responses. While its primary use is for treating bradycardia and for other medical purposes, excessive dosages can lead to CNS excitability, manifesting as restlessness, confusion, hallucinations, and can culminate in seizures. The other options listed are not typically associated with high doses of Atropine. Diarrhea is more related to stimulation of the gastrointestinal tract and is not a common effect of Atropine, which actually reduces secretions and motility. Hypoglycemia is not a direct effect of Atropine; it's more relevant to other medications that directly influence insulin secretion. Depression is also not a recognized consequence of Atropine overdose, as the drug does not have the properties that would lead to a depressive state. Therefore, the connection between high doses of Atropine and the likelihood of seizures is well-established in pharmacological literature.

SAMPLE

5. What type of analgesic is methoxyflurane classified as?

A. Inhaled analgesic

B. Topical analgesic

C. Injectable analgesic

D. Oral analgesic

Methoxyflurane is classified as an inhaled analgesic. This classification is based on its method of administration and its pharmacological action. Inhaled analgesics are typically delivered via inhalation, allowing them to quickly enter the bloodstream through the lungs and provide rapid pain relief. Methoxyflurane specifically acts by depressing the central nervous system and modifying the perception of pain, making it particularly effective in pre-hospital settings for managing acute pain. The inhaled route allows for ease of use, particularly in emergency situations, where quick administration and onset are crucial. This is in contrast to topical analgesics, which are applied to the skin; injectable analgesics, which are delivered via injection; and oral analgesics, which are consumed and absorbed through the gastrointestinal tract. Each of these other methods has different indications and uses, but they do not apply to methoxyflurane.

6. Which type of medications is oxygen commonly used as a drive gas to administer?

A. Oral medications

B. Intravenous medications

C. Nebulised medications

D. Sublingual medications

Oxygen is commonly used as a drive gas in the administration of nebulised medications because the nebulisation process converts liquid medication into a fine mist that can be inhaled directly into the lungs. When nebulised, oxygen can help increase the flow and delivery of medication, ensuring that it reaches the lower airways efficiently. This method is particularly beneficial for patients with respiratory conditions, as it allows for rapid absorption and a targeted therapeutic effect. In contrast, oral, intravenous, and sublingual medications do not require the use of oxygen as a drive gas. Oral medications are taken by mouth and absorbed through the digestive system, intravenous medications are delivered directly into the bloodstream through a vein, and sublingual medications are absorbed under the tongue. Each of these routes uses different methods of administration that do not involve the nebulisation process or reliance on oxygen as a drive gas.

SAMPLE

7. What effect does Naloxone primarily reverse in patients?

A. Increased heart rate

B. Respiratory depression

C. Fever and chills

D. Muscle spasms

Naloxone primarily reverses respiratory depression, which is a critical effect of opioid overdose. Opioids bind to specific receptors in the brain that not only provide pain relief but also suppress the respiratory centers. When someone experiences an overdose, their respiratory function can become severely impaired, leading to shallow or halted breathing. Naloxone is an opioid antagonist, meaning it competes with opioids for the same receptor sites and effectively displaces them. By doing so, it restores normal respiratory function. Administering Naloxone can rapidly improve the patient's breathing and oxygenation, which are vital for survival in cases of opioid overdose. This makes it an essential drug in pre-hospital settings where quick intervention can save lives. The other options, while relevant to different medical conditions, are not primarily influenced by Naloxone. For instance, increased heart rate can occur for various reasons not solely related to opioid action. Fever and chills are typically indicative of infections or other systemic responses, and muscle spasms may relate to different neurological or musculoskeletal issues. None of these conditions are reversed through the action of Naloxone as effectively as respiratory depression is addressed.

8. What is a significant risk associated with oxygen use?

A. Increased blood pressure

B. Impaired oxygenation

C. Increased fire and explosion risk

D. Infection risk

Oxygen is a vital therapy in medical settings, particularly for patients experiencing respiratory distress or other conditions that lead to inadequate oxygenation. However, one significant risk associated with its use is the increased fire and explosion risk. Oxygen supports combustion, meaning that it can enhance the intensity and spread of fires. In environments where high concentrations of oxygen are present, even small sparks can ignite materials that would not normally catch fire in atmospheric air. This is why strict safety protocols are followed when administering oxygen. Ensuring proper handling and storage of oxygen cylinders, avoiding open flames, and using non-flammable materials in the vicinity of oxygen are crucial measures to mitigate this risk. Understanding this aspect of oxygen use is essential for safe practice in emergency medical situations.

SAMPLE

9. What is the primary action of Frusemide?

A. Inhibition of renal sodium reabsorption

B. Promotion of potassium retention

C. Decrease in urine output

D. Stimulation of liver glycogen conversion

Frusemide, commonly known as furosemide, primarily functions as a loop diuretic, and its main action is the inhibition of renal sodium reabsorption. It targets the sodium-potassium-chloride (Na-K-2Cl) co-transporter found in the ascending limb of the Loop of Henle in the kidneys. By disrupting this transport mechanism, Frusemide prevents the reabsorption of sodium, chloride, and water, resulting in increased urine production and the excretion of these electrolytes. This mechanism is crucial in clinical practice, particularly for conditions such as heart failure, hypertension, and renal disorders, where excess fluid retention can be problematic. The increased diuresis helps to reduce fluid overload, decrease blood pressure, and improve symptoms related to congestion. The other options do not accurately describe the primary action of Frusemide. While potassium levels may be affected due to increased urinary excretion, the drug does not promote potassium retention; in fact, it can lead to hypokalemia. Additionally, instead of decreasing urine output, Frusemide actually increases urinary excretion. Lastly, there is no significant role of Frusemide in the stimulation of liver glycogen conversion. Understanding Fruse

10. What is a reported adverse effect of paracetamol?

A. Asthma attacks

B. Nausea

C. Rash

D. Heart palpitations

Paracetamol, also known as acetaminophen, is widely used for pain relief and fever reduction. One of the commonly reported adverse effects of paracetamol is nausea. This side effect can be attributed to its effect on the gastrointestinal system, and while it is generally well-tolerated, some individuals may experience discomfort or nausea, particularly if the medication is taken on an empty stomach or in higher-than-recommended dosages. The other options, while they may be associated with various medications or conditions, are not typical adverse effects of paracetamol. Asthma attacks can be triggered by other medications, particularly non-steroidal anti-inflammatory drugs (NSAIDs) in susceptible populations. Rash can occur with some medications as an allergic reaction, but it is not a common side effect of paracetamol. Heart palpitations are generally associated with stimulatory drugs or conditions affecting heart rhythm, rather than paracetamol use.

SAMPLE

Next Steps

Congratulations on reaching the final section of this guide. You've taken a meaningful step toward passing your certification exam and advancing your career.

As you continue preparing, remember that consistent practice, review, and self-reflection are key to success. Make time to revisit difficult topics, simulate exam conditions, and track your progress along the way.

If you need help, have suggestions, or want to share feedback, we'd love to hear from you. Reach out to our team at hello@examzify.com.

Or visit your dedicated course page for more study tools and resources:

https://nswambpharmacology.examzify.com

We wish you the very best on your exam journey. You've got this!

SAMPLE